

[全部](#) [图片](#) [视频](#) [新闻](#) [地图](#) [图书](#)

找到约 1,150 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[Mutations in E2-PePHD, NS5A-PKRBD, NS5A-ISDR, and NS5A-V3 ...](#)jvi.asm.org/content/82/13/6644.full

Sequences within **HCV NS5A** (PKR binding domain [PKRBD], IFN ... The V3 **region** had a very high **genetic** variability, but this was **not** related to SVR. ... **regions** of the **HCV** proteins in correlation with chronic **hepatitis C (CHC)** has been ... to predict **responses** to peg-IFN- α plus RBV **therapy** in **patients** with genotype 1.

[Role of genetic polymorphisms in hepatitis C virus chronic infection](#)<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4568530/>

16 Sep 2015 ... In this review article the effect of old and **new** host **genetics** factors [interleukin 28B, ... **Patients** with chronic **hepatitis C (CHC)** frequently show the increasing ... IL28B and Peg-IFN plus ribavirin **treatment** in **patients** with **non-1 HCV** ... in the **NS5A region** on the **response** to Peg-IFN plus RBV **treatment** in 286 ...

[Amino Acid Polymorphisms Within the Entire HCV NS5A Region in ...](#)<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3867023/>

14 Dec 2013 ... Analysis of **NS5A** polymorphisms in **HCV** genotype 1b pre-**treatment** serum samples from Estonian **patients** and **their** effect on the **treatment response**. ... SVR and non-SVR **patients** in any known **regions** of the **NS5A** protein. ... 1 and 4 are **more** resistant to IFN-based **therapy** than genotypes 2 and 3 (8).

[Treatment regimens for chronic hepatitis C virus genotype 1 ...](#)www.uptodate.com/.../treatment-regimens-for-chronic-hepatitis-c-virus-genotype-1-infection-in-adults

29 Nov 2016 ... **Hepatitis C virus (HCV)** can cause both acute and chronic hepatitis. ... **NS5A** inhibitor failures: **Patients** who did **not respond** to or relapsed after ... However, the cost of these **new** agents prevents universal delivery of antiviral **therapy**. ... regimen selection by **treatment history** (algorithm 1 and algorithm 2).

[\[PDF\] A Protease Inhibitor with Induction Therapy with Natural Interferon ...](#)www.mdpi.com/1422-0067/17/3/350/pdf

9 Mar 2016 ... During IT with nIFN- β , virologic **response** rates in group A ... IT with PI/PR is

Name of Journal: World Journal of Gastroenterology

ESPS Manuscript NO: 32537

Manuscript Type: ORIGINAL ARTICLE

Basic Study

Mutations of the HCV NS5A region in CHC-1 patients non responders to two or more treatments and its relationship with response to new treatment

Paloma Muñoz de Rueda, José Manuel Fuentes Rodríguez, Rosa Quiles Pérez, Ana Gila Medina, Ana Belén Martín Álvarez, Jorge Casado Ruiz, Ángeles Ruiz Extremera, Javier Salmerón

Abstract

Match Overview

1	Internet 171 words crawled on 28-Jun-2016 www.wjgnet.com	4%
2	Crossref 131 words Quiles-Pérez, R., P. Muñoz-de-Rueda, A. Martín-Lagos Maldonado, A. Martín-Álvarez, J. Quer, and J. Salmerón.	3%
3	Crossref 62 words El-Shamy, A., I. Shoji, W. El-Akel, S. E. Bilasy, L. Deng, M. El-Raziky, D.-p. Jiang, G. Esmat, and H. Hotta. "NS5A Se	1%
4	Crossref 34 words P. Munoz de Rueda. "Mutations in E2-PePHD, NS5A-PK RBD, NS5A-ISDR, and NS5A-V3 of Hepatitis C Virus Ge	1%
5	Crossref 30 words "AASLD Abstracts", <i>Hepatology</i> , 2013.	1%
6	Crossref 26 words Ullah, Sajjad, Habib U. Rehman, and Muhammad Idrees. "Mutations in the NS5A gene are associated with resp	1%
7	Crossref 23 words M. Torres-Puente. "Genetic variability in hepatitis C virus and its role in antiviral treatment response : Genetic va ...	1%
8	Crossref 19 words El-Shamy, Ahmed. "Impact of hepatitis C virus heteroge ... eity on interferon sensitivity: An overview", <i>World Journal</i>	<1%
9	Internet 12 words crawled on 19-Oct-2015	<1%

Mutations of the HCV NS5A region in CHC-1 patients non responders to two o



全部 图片 视频 新闻 地图 图书

找到约 620 条结果

时间不限

- 过去 1 小时内
- 过去 24 小时内
- 过去 1 周内
- 过去 1 个月内
- 过去 1 年内

所有结果

精确匹配

[Mutations in E2-PePHD, NS5A-PKRBD, NS5A-ISDR, and NS5A-V3 ...](#)

jvi.asm.org/content/82/13/6644.full

Sequences within HCV NS5A (PKR binding domain [PKRBD], IFN ... regions of the HCV proteins in correlation with chronic hepatitis C (CHC) has been questioned. ... Mutations in this region are believed to influence the response to IFN therapy ... Patients without EVR (non-EVR) were considered nonresponders (NR; n = 8) ...

[Sequencing of E2 and NS5A regions of HCV genotype 3a in ...](#)

www.scielo.br/scielo.php?script=sci_arttext&pid=S0074...

The NS5A and E2 proteins of HCV genotype 1 were reported to inhibit the ... percentage of HCV genotype 3-infected patients do not respond to treatment at all. ... with HCV genotype 3a and their correlation with response to antiviral therapy ... Mutations in the E2-PePHD and NS5A region of hepatitis C virus type 1 and the ...

[Prospective Characterization of Full-Length Hepatitis C Virus NS5A ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC102099/>

In this study, the relationship between NS5A mutations and selection pressures ... In HCV-1a patients, more amino acid changes were observed in isolates from ... of NS5A, that corresponded to the previously described V3 region and a new ... within the first 2 to 6 weeks of therapy in responders but not nonresponders, ...

[Prospective characterization of full-length hepatitis C virus NS5A ...](#)

<https://www.ncbi.nlm.nih.gov/pubmed/10982347>

Nousbaum J(1), Polyak SJ, Ray SC, Sullivan DG, Larson AM, Canithers RL Jr, Gretch DR, ... in the inherent resistance of HCV to interferon (IFN) antiviral therapy in clinical studies. In this study, the relationship between NS5A mutations and selection ..., nonresponders in both HCV-1a- and HCV-1b-infected patients (P<0.05).

[Treatment regimens for chronic hepatitis C virus genotype 1 ...](#)

www.uptodate.com/.../treatment-regimens-for-chronic-hepatitis-c-virus-genotype-1-infection-in-adults

29 Nov 2016 ... (See "Investigational therapies for hepatitis C virus infection".) ... Partial

[全部](#)[新闻](#)[图片](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 257 条结果 (用时 0.83 秒)

[Mutations in E2-PePHD, NS5A-PKRBD, NS5A-ISDR, and NS5A-V3 of ...](#)

jvi.asm.org/content/82/13/6644.full ▾ [翻译此页](#)

作者：PM de Rueda - 2008 - 被引用次数：70 - 相关文章

Sequences within **HCV NS5A** (PKR binding domain [PKRBD], IFN ... those with sustained virologic **responses** (SVR; n = 36) or **nonresponders** (NR; n = 24). ... **regions** of the **HCV** proteins in correlation with chronic **hepatitis C (CHC)** has **Two patients** with one **mutation** had EVR, and later, they were viral RNA positive.

[Prospective Characterization of Full-Length Hepatitis C Virus NS5A ...](#)

<https://www.ncbi.nlm.nih.gov> > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者：JB Nousbaum - 2000 - 被引用次数：151 - 相关文章

In this study, the **relationship** between **NS5A mutations** and selection pressures ... In **HCV-1a patients**, **more** amino acid changes were observed in isolates from ... of **NS5A**, that corresponded to the previously described **V3 region** and a **new** **Two patients** were end-of-treatment **nonresponders** (**patients** 9 and 20), while ...

缺少字词：chc treatments

[Role of genetic polymorphisms in hepatitis C virus chronic infection](#)

<https://www.ncbi.nlm.nih.gov> > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者：N Coppola - 2015 - 被引用次数：3 - 相关文章

2015年9月16日 - **Patients** with chronic **hepatitis C (CHC)** frequently show the increasing severity IL28B and Peg-IFN plus ribavirin **treatment** in **patients** with **non-1 HCV** ... in the **NS5A region** on the **response** to Peg-IFN plus RBV **treatment** in 286 old and **new** combination **treatments** to eradicate **HCV** infection[107,108].

[Treatment regimens for chronic hepatitis C virus genotype 1 infection ...](#)

www.uptodate.com/.../treatment-regimens-for-chronic-hepatitis-c-virus-gen... ▾ [翻译此页](#)