

Couples of patients with ulcerative colitis exhibit a biologically relevant dysbiosis

全部 图片 视频 新闻 地图 图书

找到约 438 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[Microbiota in Inflammatory Bowel Disease Pathogenesis and Therapy](#)

ncp.sagepub.com/content/early/2015/10/09/0884533615606898.full

9 Oct 2015 ... Inflammatory bowel disease (IBD), including ulcerative colitis, Crohn's They also have increased mucosal permeability, as do patients with CD. In both CD and UC, the fecal microbiota (combination of shed Siblings of patients with Crohn's disease exhibit a biologically relevant dysbiosis in mucosal ...

[The Effects of Bowel Preparation on Microbiota-Related Metrics ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4817412/>

11 Feb 2016 ... Bowel preparations affect the composition and diversity of the fecal IBD patients were eligible for participation if the patient had: (1) documented ulcerative colitis (UC) or Crohn's Siblings of patients with Crohn's disease exhibit a biologically relevant dysbiosis in mucosal microbial metacommunities ...

[Intestinal Microbiota and the Innate Immune System – A Crosstalk in ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4585200/>

22 Sep 2015 ... Given that CD patients display changes in their gut microbiota ... affecting the gastrointestinal tract and together with ulcerative colitis (UC) Investigations on fecal microbiota composition in first-degree relatives of patients with IBD Crohn's disease exhibit a biologically relevant dysbiosis in mucosal ...

[Consistent Prebiotic Effect on Gut Microbiota With Altered FODMAP ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4855163/>

14 Apr 2016 ... Five-day fecal samples were collected at the end of each diet and analyzed for ... However, reducing FODMAP intake in 72 IBD patients (including 52 with However, there are survey data showing 57% of patients with ulcerative colitis Crohn's disease exhibit a biologically relevant dysbiosis in mucosal ...

[Preclinical disease and preventive strategies in IBD: perspectives ...](#)

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 33550

Manuscript Type: ORIGINAL ARTICLE

Observational Study

Couples of patients with ulcerative colitis exhibit a biologically relevant dysbiosis in fecal microbial metacommunities

Guang-Lan Chen, Ye Zhang, Wang-Yue Wang, Xue-Liang Ji, Fei Meng,
Pei-Song Xu, Ning-Min Yang, Fu-Qiang Ye, Xiao-Chen Bo

Abstract

AIM

To investigate alterations in the fecal microbiome using 16S rRNA amplicon

Match Overview

1

Internet 14 words
crawled on 23-Oct-2015
www.ecco-ibd.eu

<1%

Couples of patients with ulcerative colitis exhibit a biologically relevant dysbiosis

全部 图片 视频 新闻 地图 图书

找到约 928 条结果

时间不限

过去 1 小时内

过去 24 小时内

过去 1 周内

过去 1 个月内

过去 1 年内

所有结果

精确匹配

[Intestinal Microbiota and the Innate Immune System – A Crosstalk in ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4585200/>

22 Sep 2015 ... Given that CD **patients** display changes in their gut **microbiota** ... Keywords: Crohn's disease, innate immune system, inflammation, **microbiota**, **dysbiosis** ... affecting the gastrointestinal tract and together with **ulcerative colitis** (UC) with Crohn's disease **exhibit a biologically relevant dysbiosis** in mucosal ...

[Involvement of Reduced Microbial Diversity in ... - NCBI - NIH](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5198157/>

15 Dec 2016 ... Sha et al, analyzed the diversity of **fecal microbiota** in IBD **patients** compared with Comparison of the **fecal microbiota** profiles between **ulcerative colitis** and Crohn's disease using Siblings of **patients** with Crohn's disease **exhibit a biologically relevant dysbiosis** in mucosal microbial **metacomunities**.

[Sampling Strategies for Three-Dimensional Spatial Community ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5323387/>

24 Feb 2017 ... Y-axis distribution includes the oral cavity and **feces** (2014) compared IBD **patients** and animal models of colitis with of the intestinal **mucus barrier** in **ulcerative colitis**: a clue to pathogenesis Siblings of **patients** with Crohn's disease **exhibit a biologically relevant dysbiosis** in mucosal **microbial** ...

[Preclinical disease and preventive strategies in IBD: perspectives ...](#)

gut.bmj.com/content/early/2016/05/03/gutjnl-2016-311785

3 May 2016 ... Overall, 31% of **patients** with CD and 0% of controls tested positive for ASCA. Repopulating the gut habitat with a healthy community using **faecal** Intestinal permeability in **patients** with Crohn's disease and **ulcerative colitis** and their first **exhibit a biologically relevant dysbiosis** in mucosal **microbial** ...

[The Effects of Bowel Preparation on Microbiota-Related Metrics ...](#)


[All](#)
[Images](#)
[News](#)
[Videos](#)
[Shopping](#)
[More](#)
[Settings](#)
[Tools](#)

About 488 results (0.83 seconds)

Fecal Microbiota Transplantation for Inflammatory Bowel Disease

<https://www.ncbi.nlm.nih.gov> › NCBI › Literature › PubMed Central (PMC)

by J Lopez - 2016 - Cited by 6 - Related articles

Keywords: **Fecal microbiota** transplantation, **ulcerative colitis**, Crohn's disease, treatment-naive CD demonstrated a marked **dysbiosis** in mucosa-associated ... The mode of administration of FMT may have an **important** effect in UC **patients**.

Missing: couples exhibit biologically metacommunities

Involvement of Reduced Microbial Diversity in Inflammatory Bowel ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5198157/>

by D Gong - 2016 - Cited by 4 - Related articles

15 Dec 2016 - Sha et al. analyzed the diversity of **fecal microbiota** in IBD **patients** compared with Comparison of the **fecal microbiota** profiles between **ulcerative colitis** and Crohn's disease using Siblings of **patients** with Crohn's disease **exhibit a biologically relevant dysbiosis** in mucosal microbial **metacommunities**.

Fecal Microbiota Transplantation for Ulcerative Colitis: A Systematic ...

<https://www.ncbi.nlm.nih.gov> › NCBI › Literature › PubMed Central (PMC)

by Y Shi - 2016 - Cited by 16 - Related articles

13 Jun 2016 - **Fecal microbiota** transplantation (FMT) has been recognized as a novel ... [4] The **dysbiosis** of UC is also characterized by some alterations of ... Additionally, several **relevant** studies were reported after the last systematic review in 2014. Symptoms and stool patterns in **patients with ulcerative colitis**. Gut.

Missing: couples exhibit metacommunities

Preclinical disease and preventive strategies in IBD ... - Gut - The BMJ

gut.bmj.com › Online First

by J Torres - 2016 - Cited by 5 - Related articles

3 May 2016 - Overall, 31% of **patients** with CD and 0% of controls tested positive for ASCA. Repopulating the gut habitat with a healthy community using **faecal** Intestinal permeability in **patients** with Crohn's disease and **ulcerative colitis** and their first ... **exhibit a biologically**