

MALT lymphoma with unusual 18F-FDG hypermetabolism arising at the colorecta



全部 图片 视频 新闻 购物 地图 图书

找到约 1,830 条结果

时间不限

- 过去 1 小时内
- 过去 24 小时内
- 过去 1 周内
- 过去 1 个月内
- 过去 1 年内

所有结果

精确匹配

[Uncommon Colorectal Neoplasms: Colorectal Lymphoma - NCBI](#)

[www.ncbi.nlm.nih.gov/pmc/articles/PMC2780105/](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2780105/)

Most patients with primary colorectal lymphoma present with abdominal pain ... MALT-associated lymphomas are low-grade tumors arising from B cells ... Most often colorectal lymphomas are large, polypoid lesions and mucosal ulceration is unusual. ... in lymphoma and is considered the standard of care by many authors. ...

[Uncommon Colorectal Neoplasms: Colorectal Lymphoma - NCBI](#)

[www.ncbi.nlm.nih.gov/pmc/articles/PMC3311500/](http://www.ncbi.nlm.nih.gov/pmc/articles/PMC3311500/)

Colorectal lymphomas are rare and comprise 10–20% of all gastrointestinal lymphomas ... in the medical literature that follow strict Dawson criteria..... The number of subjects ... The second type of MALT arises in the setting of a chronic inflammatory or .... IBD and lymphoma is an unusual event and IBD associated with GI ...

[API2-MALT1 fusion gene in colorectal lymphoma. - NCBI](#)

[www.ncbi.nlm.nih.gov/pubmed/14681324](http://www.ncbi.nlm.nih.gov/pubmed/14681324)

The API2-MALT1 fusion gene was originally identified from a t(11;18)(q21;q21) ... Gastric MALT lymphomas positive for the API2-MALT1 fusion gene do not ... and clinicopathological behavior of colorectal MALT lymphoma with this genetic ...

[Clinical Applications of PET in Oncology | Radiology - RSNA](#)

[pubs.rsna.org/doi/full/10.1148/radiol.2012021195](http://pubs.rsna.org/doi/full/10.1148/radiol.2012021195)

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 30183

Manuscript Type: CASE REPORT

**MALT lymphoma with unusual  $^{18}\text{F}$ -FDG hypermetabolism arising at the colorectal anastomosis**

Na-Sha Zhang, Fang Shi, Li Kong, Hui Zhu


### Abstract

Mucosa-associated lymphoid tissue (MALT) lymphoma usually originates from the stomach and presents with low  $^{18}\text{F}$ -fluorodeoxyglucose (FDG) avidity with average maximum standard uptake value of 3.6. Colorectal MALT lymphoma is a rare entity that contributes to 1.6% of all MALT lymphomas and < 0.2% of large intestinal malignancies. The case reported herein firstly revealed stage II MALT

### Match Overview

1	<b>Crossref</b> 46 words Carrillo-Cruz, Estrella, Victor A. Marín-Oyaga, Fátima de la Cruz Vicente, Isabel Borrego-Dorado, Marta Ruiz Mercado, I	2%
2	<b>Internet</b> 17 words crawled on 15-Mar-2016 <a href="http://www.wjgnet.com">www.wjgnet.com</a>	1%
3	<b>Publications</b> 15 words Hosein, Peter J.. "The Evolving Role of $^{18}\text{F}$ -FDG PET Scans in Patients with Aggressive non-Hodokin's Lymphoma"	1%
4	<b>Publications</b> 15 words Hosaka, S. Nakamura, N. Akamatsu, T. Fuj. "A case of primary low grade mucosa associated lymphoid tissue (MALT ..."	1%
5	<b>Internet</b> 15 words crawled on 09-Nov-2012 <a href="http://www.uscap.org">www.uscap.org</a>	1%
6	<b>Crossref</b> 13 words Pulmonary Involvement in Patients with Hematological Malignancies, 2011.	1%

MALT lymphoma with unusual 18F-FDG hypermetabolism arising at the colorecta



全部 图片 视频 新闻 地图 图书

找到约 1,850 条结果

时间不限

- 过去 1 小时内
- 过去 24 小时内
- 过去 1 周内
- 过去 1 个月内
- 过去 1 年内

所有结果

精确匹配

[Uncommon Colorectal Neoplasms: Colorectal Lymphoma - NCBI](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2780105/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2780105/>

Most patients with primary colorectal lymphoma present with abdominal pain ... MALT-associated lymphomas are low-grade tumors arising from B cells ... Most often colorectal lymphomas are large, polypoid lesions and mucosal ulceration is unusual. ... in lymphoma and is considered the standard of care by many authors. ...

[Uncommon Colorectal Neoplasms: Colorectal Lymphoma - NCBI](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3311500/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3311500/>

Colorectal lymphomas are rare and comprise 10–20% of all gastrointestinal lymphomas ... in the medical literature that follow strict Dawson criteria..... The number of subjects ... The second type of MALT arises in the setting of a chronic inflammatory or .... IBD and lymphoma is an unusual event and IBD associated with GI ...

[Primary Lymphoma of the Colon - NCBI - National Institutes of Health](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2981851/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2981851/>

Primary lymphoma of the colon is a rare tumor of the gastrointestinal (GI) tract and ... was performed with transverse-sigmoid, end-to-end anastomosis. ... a fleshy growth arising from the cecal wall with lymph nodes in the mesentery. .... 1994;18:291–7. ... of gastrointestinal lymphomas other than low grade MALT lymphomas.

[Clinical Applications of PET in Oncology | Radiology - RSNA](https://pubs.rsna.org/doi/full/10.1148/radiol.2312021185)

[pubs.rsna.org/doi/full/10.1148/radiol.2312021185](https://pubs.rsna.org/doi/full/10.1148/radiol.2312021185)

Because of the similarity of FDG and glucose, glucose metabolic rates can be ... of lung cancer, colorectal cancer, esophageal cancer, head and neck cancer, lymphoma, and melanoma. ... <sup>18</sup>F is by far the most widely used PET radionuclide. .... However, most hypermetabolic pulmonary nodules are malignant, and a lesion ...

[PDF] Book of Abstracts - Human Health Campus - International Atomic ...



全部 图片 新闻 视频 购物 更多

设置 工具

找到约 2,890 条结果 (用时 0.92 秒)

### Uncommon Colorectal Neoplasms: Colorectal Lymphoma - NCBI

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2780105/> - 翻译此页

作者：FJ Quayle - 2006 - 被引用次数：41 - 相关文章

Most patients with primary **colorectal lymphoma** present with abdominal pain, and ... rates of **mucosa-associated lymphoid tissue (MALT) lymphoma** and mantle cell ... are low-grade tumors **arising** from B cells associated with mucosal immunity. ... **lymphomas** are large, polypoid lesions and mucosal ulceration is **unusual**.

缺少字词：fdg hypermetabolism anastomosis

### Colonic Mucosa-Associated Lymphoid Tissue Lymphoma - NCBI

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者：R Akasaka - 2012 - 被引用次数：10 - 相关文章

2012年8月29日 - Although rare, awareness of MALT **lymphoma** of the **colon** is ... **Mucosa-associated lymphoid tissue (MALT)** is found in various parts of the body, .... The patient underwent sigmoidectomy which revealed an **18 mm** tumor (fig.

缺少字词：unusual fdg hypermetabolism arising anastomosis

### Colonic mucosa-associated lymphoid tissue lymphoma identified by ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4277991/> - 翻译此页

作者：SW Seo - 2014 - 被引用次数：2 - 相关文章

2014年12月28日 - Core tip: **Mucosa-associated lymphoid tissue (MALT) lymphoma** in the large ... **Colorectal MALT lymphoma** is a rare disease, which accounts for a .... though **colorectal MALT lymphomas** are rarer than those **arising** from ... Tumor cells of MALT **lymphoma** are also positive for Bcl-2 and negative for Bcl-6[18].

缺少字词：unusual fdg hypermetabolism anastomosis

### Clinical Applications of PET in Oncology | Radiology - RSNA

[pubs.rsna.org/doi/full/10.1148/radiol.2312021185](https://pubs.rsna.org/doi/full/10.1148/radiol.2312021185) ▼ 翻译此页

作者：EM Behren - 2004 - 被引用次数：776 - 相关文章