

[全部](#)[图片](#)[购物](#)[新闻](#)[视频](#)[更多](#)[设置](#)[工具](#)

找到约 15,200 条结果 (用时 0.86 秒)

[PDF] New test meal Refex® 13C-UBT to detect Hp in patients on PPI

media.klimud.org/Klimud.../sitkeaygen_antalya%202015-11-20_infai.pdf ▼ [翻译此页](#)

The Sensitivity and Specificity of the Modified Helicobacter Test INFAL. Using New Test Meal with 13C-UBT in Helicobacter Pylori Positive and. Negative Patients with Dyspepsia Taking PPI. Sitke Aygen. Institut für ... 13C-Urea breath test.

Accuracy of breath tests using low doses of 13C - urea to diagnose ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: L Gatta - 2006 - 被引用次数: 55 - [相关文章](#)

The ¹³C - urea breath test (UBT) for detecting Helicobacter pylori infection is a ... has been extensively modified, including variations in the cut off values, test meal, ... at least 200 more consecutive dyspeptic patients to be tested with this UBT. ... H pylori negative) and one (female H pylori negative) patients, tested with 10 ...

Dyspepsia: When and How to Test for Helicobacter pylori Infection

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: MP Dore - 2016 - 被引用次数: 5 - [相关文章](#)

2016年4月28日 - The most common symptoms in patients with functional dyspepsia in whom dyspepsia remains despite H. pylori eradication, a trial of PPI ... 13C-urea breath tests. ... As with any test, prevalence of the H. pylori infection and the pretest ... To reduce false negative results, the patient should be off antibiotics ...

An optimized 13C-urea breath test for the diagnosis of H pylori infection

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: G Camuzano Maya - 2007 - 被引用次数: 37 - [相关文章](#)

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 32793

Manuscript Type: ORIGINAL ARTICLE

Prospective Study

6
Modified *Helicobacter* test using a new test meal and a ¹³C-urea breath test in *Helicobacter pylori* positive and negative dyspepsia patients on proton pump inhibitors

Tepes B *et al.* New test meal with ¹³C-urea breath test in patients on PPIs

Bojan Tepes, Peter Malfertheiner, Joachim Labenz, Sitke Aygen

Abstract

AIM

To determine the sensitivity and specificity of the ¹³C-urea breath test (UBT) in patients taking proton pump inhibitors (PPIs), using a new test meal Refex.

Match Overview

1	Crossref 58 words A. Agha. "Effect of different organic acids (citric, malic a ... d ascorbic) on intragastric urease activity", <i>Alimentary Ph</i>	2%
2	Crossref 37 words Luigi Gatta. "Effect of Proton Pump Inhibitors and Antacid Therapy on 13C Urea Breath Tests and Stool Test for H ...	1%
3	Crossref 31 words on behalf of the ACTFAST investigators. "Effect of individ ualizing starting doses of a statin according to baseline	1%
4	Crossref 25 words Mana, F.. "Cut-off point, timing and pitfalls of the ¹⁴ C-urea breath test as measured by infrared spectrometry",	1%
5	Crossref 24 words Dino Vaira. "A comparison amongst three rapid urease t ests to diagnose <i>Helicobacter pylori</i> infection in 375 co ...	1%
6	Internet 21 words crawled on 29-Nov-2015 public.ukcrn.org.uk	1%
7	Internet 19 words crawled on 15-Sep-2016 fin.nalgrave-journals.com	1%

[全部](#)[图片](#)[购物](#)[新闻](#)[视频](#)[更多](#)[设置](#)[工具](#)

找到约 15,200 条结果 (用时 0.61 秒)

[PDF] New test meal Refex® 13C-UBT to detect Hp in patients on PPI

media.klimud.org/Klimud.../sitkeaygen_antalya%202015-11-20_infai.pdf - 翻译此页

The Sensitivity and Specificity of the Modified Helicobacter Test INFAI. Using New Test Meal with 13C-UBT in Helicobacter Pylori Positive and. Negative Patients with Dyspepsia Taking PPI. Sitke Aygen. Institut für ... 13C-Urea breath test.

Accuracy of breath tests using low doses of 13C - urea to diagnose ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: L Gatta - 2006 - 被引用次数: 55 - 相关文章

The ¹³C - urea breath test (UBT) for detecting Helicobacter pylori infection is a ... has been extensively modified, including variations in the cut off values, test meal, ... at least 200 more consecutive dyspeptic patients to be tested with this UBT. ... H pylori negative) and one (female H pylori negative) patients, tested with 10 ...

Dyspepsia: When and How to Test for Helicobacter pylori Infection

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: MP Dore - 2016 - 被引用次数: 5 - 相关文章

2016年4月28日 - The most common symptoms in patients with functional dyspepsia in whom dyspepsia remains despite H. pylori eradication, a trial of PPI ... 13C-urea breath tests. ... As with any test, prevalence of the H. pylori infection and the pretest ... To reduce false negative results, the patient should be off antibiotics ...

An optimized 13C-urea breath test for the diagnosis of H pylori infection

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页



Modified Helicobacter test using a new test meal and a 13C-urea breath test in



全部

图片

购物

新闻

视频

更多

设置

工具

找到约 15,300 条结果 (用时 0.79 秒)

Accuracy of breath tests using low doses of 13C-urea to diagnose ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者 : L Gatta - 2006 - 被引用次数 : 55 - 相关文章

The ¹³C-urea breath test (UBT) for detecting *Helicobacter pylori* infection is a ... has been extensively modified, including variations in the cut off values, test meal, ... at least 200 more consecutive dyspeptic patients to be tested with this UBT. ... *H pylori* negative) and one (female *H pylori* negative) patients, tested with 10 ...

Dyspepsia: When and How to Test for Helicobacter pylori Infection

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者 : MP Dore - 2016 - 被引用次数 : 5 - 相关文章

2016年4月28日 - The most common symptoms in patients with functional dyspepsia in whom dyspepsia remains despite *H. pylori* eradication, a trial of PPI ... 13C-urea breath tests. ... As with any test, prevalence of the *H. pylori* infection and the pretest ... To reduce false negative results, the patient should be off antibiotics ...

[PDF] New test meal Refex® 13C-UBT to detect Hp in patients on PPI

media.klimud.org/Klimud.../sitkeaygen_antalya%202015-11-20_infai.pdf ▼ 翻译此页

The Sensitivity and Specificity of the Modified Helicobacter Test INFAI. Using New Test Meal with 13C-UBT in *Helicobacter Pylori* Positive and. Negative Patients with Dyspepsia Taking PPI. Sitke Aygen. Institut für ... 13C-Urea breath test.

[PDF] Urea Breath Tests for Detection of Helicobacter pylori - Numilog

www.numilog.com/package/extraits_pdf/e238349.pdf ▼ 翻译此页

作者 : S WANG - 被引用次数 : 1 - 相关文章

and all patients with confirmed peptic ulcer disease associated with *H. pylori* infec- ... therapy with a proton pump inhibitor or ranitidine bismuth citrate, 13C-urea breath test is considered a standard noninvasive test for both initial diagno- positive and -negative patients is 2 min with endoscopic administration and 6 min