

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 33359

Manuscript Type: ORIGINAL ARTICLE

Basic Study

Circulating inflammatory factors associated with worse long-term prognosis in colorectal cancer

Olsen RS *et al.* Circulating inflammatory factors in CRC

Renate S Olsen, Johnny Nijm, Roland E Andersson, Jan Dimberg, Dick Wågsäter

Abstract

AIM

To investigate association of circulating inflammatory factors at the time of colorectal cancer (CRC) surgery with survival.

METHODS

Match Overview

1	Internet 29 words crawled on 25-Dec-2016 pdfs.semanticscholar.org	1%
2	Internet 23 words www.umdj.edu	1%
3	Crossref 15 words Giuseppe Di Caro, Michele Carvello, Samantha Pesce, Marco Erreni, Federica Marchesi, Jelena Todoric, Matteo Sacch	<1%
4	Crossref 13 words Jan Dimberg "Polymorphisms of Fractalkine receptor CX3CR1 and plasma levels of its ligand CX3CL1 in colorecta ...	<1%
5	Internet 11 words crawled on 18-Jun-2017 www.dtic.mil	<1%
6	Internet 11 words crawled on 29-Jun-2017 boris.unibe.ch	<1%



找到约 922,000 条结果 (用时 0.34 秒)

Google 学术: Circulating inflammatory factors associated with worse long-term prognosis in colorectal cancer

Impact of anastomotic leakage on long-term survival of ... - McArdle - 被引用次数: 335

... leakage is associated with poor long-term outcome in ... - Law - 被引用次数: 268

... lymphocyte ratio as a prognostic predictor after curative ... - Gomez - 被引用次数: 275

Circulating Inflammatory Mediators as Potential Prognostic Markers of ...

<https://www.ncbi.nlm.nih.gov> › [NCBI](#) › [Literature](#) › [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者: G Di Caro - 2016 - 被引用次数: 6 - [相关文章](#)

2016年2月9日 - Novel highly sensitive tools are needed to identify colorectal cancer patients at high risk ... levels were significantly associated with worse survival ($P < 0.001$). ... If the death event was unrelated to CRC disease, time to follow-up was We showed that CXCL-8, VEGF and the long Pentraxin 3 were the best ...

Colorectal cancer tumour markers and biomarkers: Recent therapeutic ...

<https://www.ncbi.nlm.nih.gov> › [NCBI](#) › [Literature](#) › [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者: G Lech - 2016 - 被引用次数: 27 - [相关文章](#)

2016年2月7日 - Colorectal cancer (CRC) is the second most commonly diagnosed ... Some of them are currently found to be more important during treatment and long-term follow-up. ... or circulating tumour cells to identify new CRC screening markers. has been reported to be associated with the worse prognosis[82].


Atrial fibrillation and survival in colorectal cancer - NCBI - NIH

<https://www.ncbi.nlm.nih.gov> › [NCBI](#) › [Literature](#) › [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者: SR Walsh - 2004 - 被引用次数: 6 - [相关文章](#)

2004年11月29日 - Survival in colorectal cancer may correlate with the degree of ... Atrial fibrillation may be regarded as an inflammatory complication. ... or may not [10] be associated with worse survival following surgery for lung cancer. ... Patients with no recurrence were censored at the time of death from any Circulation.

Circulating inflammatory factors associated with worse long-term prognosis in co



全部 图片 视频 新闻 购物 地图 图书

找到约 919,000 条结果

时间不限

- 过去 1 小时内
- 过去 24 小时内
- 过去 1 周内
- 过去 1 个月内
- 过去 1 年内

所有结果

精确匹配

[Google 学术: Circulating inflammatory factors associated with worse long-term prognosis in colorectal cancer](#)

[Impact of anastomotic leakage on long-term survival of ...](#) - McArdle - 被引用次数: 335

[... leakage is associated with poor long-term outcome in ...](#) - Law - 被引用次数: 268

[... lymphocyte ratio as a prognostic predictor after curative ...](#) - Gomez - 被引用次数: 277

[Circulating Inflammatory Mediators as Potential Prognostic Markers ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4747470/>

9 Feb 2016 ... Novel highly sensitive tools are needed to identify colorectal cancer patients at high risk ... levels were significantly associated with worse survival (P<0.001). ... If the death event was unrelated to CRC disease, time to follow-up was We showed that CXCL-8, VEGF and the long Pentraxin 3 were the best ...

[Colorectal cancer tumour markers and biomarkers: Recent ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4724606/>

7 Feb 2016 ... Colorectal cancer (CRC) is the second most commonly diagnosed ... Some of them are currently found to be more important during treatment and long-term follow-up ... or circulating tumour cells to identify new CRC screening markers. has been reported to be associated with the worse prognosis[82].

[Atrial fibrillation and survival in colorectal cancer - NCBI - NIH](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC538254/>

29 Nov 2004 ... Survival in colorectal cancer may correlate with the degree of ... Atrial fibrillation may be regarded as an inflammatory complication. ... or may not [10] be associated with worse survival following surgery for lung cancer. ... Patients with no recurrence were censored at the time of death from any ... Circulation.

[PDF] [Circulating and genetic factors in colorectal cancer Potential ...](#) - DiVA

<https://liu.diva.portal.org/smash/get/diva2.../FULLTEXT01.pdf>



全部 新闻 图片 购物 视频 更多

设置 工具

找到约 922,000 条结果 (用时 0.68 秒)

Google 学术: Circulating inflammatory factors associated with worse long-term prognosis in colorectal cancer

Impact of anastomotic leakage on long-term survival of ... - McArdle - 被引用次数: 335

... leakage is associated with poor long-term outcome in ... - Law - 被引用次数: 268

... lymphocyte ratio as a prognostic predictor after curative ... - Gomez - 被引用次数: 275

Circulating Inflammatory Mediators as Potential Prognostic Markers of ...

<https://www.ncbi.nlm.nih.gov> › NCBI › Literature › PubMed Central (PMC) - 翻译此页

作者: G Di Caro - 2016 - 被引用次数: 6 - 相关文章

2016年2月9日 - Novel highly sensitive tools are needed to identify colorectal cancer patients at high risk ... levels were significantly associated with worse survival ($P < 0.001$). ... If the death event was unrelated to CRC disease, time to follow-up was We showed that CXCL-8, VEGF and the long Pentraxin 3 were the best ...

Atrial fibrillation and survival in colorectal cancer - NCBI - NIH

<https://www.ncbi.nlm.nih.gov> › NCBI › Literature › PubMed Central (PMC) - 翻译此页

作者: SR Walsh - 2004 - 被引用次数: 6 - 相关文章

2004年11月29日 - Survival in colorectal cancer may correlate with the degree of ... Atrial fibrillation may be regarded as an inflammatory complication. ... or may not [10] be associated with worse survival following surgery for lung cancer. ... Patients with no recurrence were censored at the time of death from any ... Circulation.

Colorectal cancer tumour markers and biomarkers: Recent therapeutic ...

<https://www.ncbi.nlm.nih.gov> › NCBI › Literature › PubMed Central (PMC) - 翻译此页

作者: G Lech - 2016 - 被引用次数: 27 - 相关文章

2016年2月7日 - Colorectal cancer (CRC) is the second most commonly diagnosed ... Some of them are currently found to be more important during treatment and long-term follow-up. ... or circulating tumour cells to identify new CRC screening markers. has been reported to be associated with the worse prognosis[82].