

Drainage fluid and serum amylase levels accurately predict development of po



全部 图片 视频 新闻 地图 图书

找到约 1,040,000 条结果

时间不限

- 过去 1 小时内
- 过去 24 小时内
- 过去 1 周内
- 过去 1 个月内
- 过去 1 年内

所有结果

精确匹配

[Google 学术: Drainage fluid and serum amylase levels accurately predict development of postoperative pancreatic fistula](#)

[Postoperative pancreatic fistula: an international study ...](#) - Bassi - 被引用次数: 3145

[Risk prediction for development of pancreatic fistula ...](#) - Pratt - 被引用次数: 228

[... of the International Study Group of Pancreatic Fistula \(...](#) - Pratt - 被引用次数: 262

[The gastric/pancreatic amylase ratio predicts postoperative ...](#) - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/25621676>

The gastric/pancreatic amylase ratio predicts postoperative pancreatic fistula ... Previously identified predictors for POPF often have low predictive accuracy. ... power of the GPAR on POD3 and the levels of serum and ascites amylase. ... The color of the pancreatic drain fluid and high GPAR were independent risk factors.

[Intra-operative amylase in peri-pancreatic fluid independently ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4925798/>

16 Jun 2016 ... Post-operative pancreatic fistula (POPF) is a common and potentially ... Measurement of IOAC allows early and accurate categorization of ... Three millilitres of newly developed intraoperative peri-pancreatic fluid ... Each surgical drain was removed post-operative as soon as amylase level in the drain fluid ...

[Drain data to predict clinically relevant pancreatic fistula - NCBI](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3030756/>

Post-operative pancreatic fistula (POPF) is a common and potentially ... was to evaluate whether drain data accurately predicts clinically significant POPF. ... of fluid with a documented bilirubin level equal to or greater than serum values or of a ... increased drainage with high amylase content signals a developing clinically ...

[Serum amylase on the night of surgery predicts clinically significant ...](#)

www.sciencedirect.com/science/article/pii/S1365182X15316063

16 Dec 2015 ... Overall, 43 (23.2%) patients developed clinically significant POPF. ...

Postoperative day 0 serum amylase of <130 IU/L allows for the early and accurate

Name of Journal: World Journal of Gastroenterology

ESPS Manuscript No: 33466

Manuscript Type: ORIGINAL ARTICLE

Prospective Cohort Study

**Drainage fluid and serum amylase levels accurately predict development of
postoperative pancreatic fistula**

Shuo Jin, Si-Yuan Wang, Xiao-Ju Shi, Ping Zhang, Guo-Yue Lv, Xiao-Hong Du,
Guang-Yi Wang

Abstract

Match Overview

- | Match Number | Source | Words | Similarity |
|--------------|--|----------|------------|
| 1 | Crossref | 12 words | 1% |
| | Yoshitsugu Tajima. "Patient allocation based on preoperative assessment of pancreatic fibrosis to secure pancreatic an | | |
| 2 | Crossref | 11 words | <1% |
| | Jin, Shuo, Xiao-Ju Shi, Xiao-Dong Sun, Ping Zhang, Guo-Yue Lv, Xiao-Hong Du, Si-Yuan Wang, and Guang-Yi Wang. | | |

Drainage fluid and serum amylase levels accurately predict development of po

全部 图片 视频 新闻 地图 图书

找到约 1,040,000 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[Google 学术: Drainage fluid and serum amylase levels accurately predict development of postoperative pancreatic fistula](#)

[Postoperative pancreatic fistula: an international study ...](#) - Bassi - 被引用次数: 3145

[Risk prediction for development of pancreatic fistula ...](#) - Pratt - 被引用次数: 228

[... of the International Study Group of Pancreatic Fistula \(...](#) - Pratt - 被引用次数: 262

[The gastric/pancreatic amylase ratio predicts postoperative ... - NCBI](#)

<https://www.ncbi.nlm.nih.gov/pubmed/25621676>

The gastric/pancreatic amylase ratio predicts postoperative pancreatic fistula ... Previously identified predictors for POPF often have low predictive accuracy. ... power of the GPAR on POD3 and the levels of serum and ascites amylase. ... The color of the pancreatic drain fluid and high GPAR were independent risk factors.

[Intra-operative amylase in peri-pancreatic fluid independently ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4925798/>

16 Jun 2016 ... Post-operative pancreatic fistula (POPF) is a common and potentially ... Measurement of IOAC allows early and accurate categorization of ... Three millilitres of newly developed intraoperative peri-pancreatic fluid ... Each surgical drain was removed post-operative as soon as amylase level in the drain fluid ...

[Drain data to predict clinically relevant pancreatic fistula - NCBI](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3030756/>

Post-operative pancreatic fistula (POPF) is a common and potentially ... was to evaluate whether drain data accurately predicts clinically significant POPF. ... of fluid with a documented bilirubin level equal to or greater than serum values or of a ... increased drainage with high amylase content signals a developing clinically ...

[Serum amylase on the night of surgery predicts clinically significant ...](#)

www.sciencedirect.com/science/article/pii/S1365182X15316063

16 Dec 2015 ... Overall, 43 (23.2%) patients developed clinically significant POPF. ... Postoperative day 0 serum amylase of <130 IU/l allows for the early and accurate categorization of ... Postoperative pancreatic fistula (POPF) persists as the most ... The

找到约 961,000 条结果 (用时 0.54 秒)

Google 学术 : Drainage fluid and serum amylase levels accurately predict development of postoperative pancreatic fistula

Postoperative pancreatic fistula: an international study ... - Bassi - 被引用次数 : 3145

Risk prediction for development of pancreatic fistula ... - Pratt - 被引用次数 : 228

... of the International Study Group of Pancreatic Fistula (... - Pratt - 被引用次数 : 262

The gastric/pancreatic amylase ratio predicts postoperative pancreatic ...

<https://www.ncbi.nlm.nih.gov/pubmed/25621676> - 翻译此页

作者 : S Jin - 2015 - 被引用次数 : 2 - 相关文章

The gastric/pancreatic **amylase** ratio predicts **postoperative pancreatic fistula** with ... power of the GPAR on POD3 and the **levels** of serum and ascites **amylase**. ... The color of the pancreatic **drain fluid and high** GPAR were independent risk factors. Although serum and ascites **amylase** did not **predict** POPF **accurately**, the ...

[PDF] The value of postoperative measurement of amylase in abdominal ...

www.ijhpd.com/archive/2012-archive/.../100009IJHPDRRTB2012-safi.pdf - 翻译此页

作者 : F Safi - 2012 - 相关文章

lipase estimations in the **serum** and abdominal ... **drainage fluid amylase levels** did not appear to ... patients **developed** any clinical signs of **fistula** and or total **amylase** output in **drain fluid** can **predict** **postoperative clinical pancreatic fistula**.

Serum amylase on the night of surgery predicts clinically significant ...

www.sciencedirect.com/science/article/pii/S1365182X15316063 - 翻译此页

作者 : LKP Velu - 2014 - 被引用次数 : 16 - 相关文章

2015年12月16日 - Overall, 43 (23.2%) patients **developed** clinically significant POPF.

Postoperative pancreatic fistula (POPF) persists as the most challenging A **drain fluid amylase** three times the **serum amylase** activity on or after PoD 3 ... of the diagnostic **accuracy** of a **test**: an AUC value of >0.50 indicates the ability of ...