

学术搜索

找到约 3,950 条结果 (用时0.05秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利

☒ 包含引用

☐ 创建快讯

小提示: 只搜索 中文(简体) 结果, 可在 学术搜索设置 指定搜索语言

Long-term prognostic significance of extent of rectal cancer response to preoperative radiation and chemotherapy

L Ruo, S Tickoo, DS Klimstra, BD Minsky, L Saltz... - ... of surgery, 2002 - journals.lww.com
... journal alerts and information, but is committed to maintaining your privacy and will not share ...
as a categorical (grade response) or continuous (percentage response) variable, also did not
have a ... 13,14 Patients with resected rectal cancer with a complete pathologic response to ...
被引用次数: 275 相关文章 所有 8 个版本 引用 保存

[PDF] researchgate.net

Circumferential resection margin involvement: an independent predictor of survival following surgery for oesophageal cancer

SPL Dexter, H Sue-Ling, MJ McMahon, P Quirke... - Gut, 2001 - gut.bmj.com
... For all subsequent survival analyses, cancer specific values will be used. ... Although the study did
not specifically examine patterns of recurrence, locoregional recurrence is common following ...
(1991) Rising incidence of adenocarcinoma of the esophagus and gastric cardia, ...
被引用次数: 150 相关文章 所有 11 个版本 引用 保存

[HTML] bmj.com

Does graded histologic response after neoadjuvant chemotherapy predict survival for completely resected gastric cancer?

JC Mansour, L Tang, M Shah, D Bentrem... - Annals of surgical ..., 2007 - Springer
... Diffuse/entire stomach. ... Our results do not clearly define a role for the use of GHR or
perineural-vascular invasion as a factor ... treatment perineural and vascular invasion (PVI) predicts
survival following preoperative therapy for patients with locally advanced gastric cancer (LAGC ...
被引用次数: 61 相关文章 所有 7 个版本 引用 保存

Lymph node ratio as a prognostic factor in patients with stage III rectal cancer treated with total mesorectal excision followed by chemoradiotherapy

YS Kim, JH Kim, SM Yoon, EK Choi, S Do Ahn... - International Journal of ..., 2009 - Elsevier
PURPOSE: To investigate the prognostic impact of lymph node ratio (LNR) on survival in the
patients with Stage III rectal cancer. METHODS AND MATERIALS: We retrospectively
reviewed the data of 421 consecutive patients who underwent total mesorectal excision ...
被引用次数: 69 相关文章 所有 8 个版本 引用 保存

Histomorphologic tumor regression and lymph node metastases determine

[PDF] researchgate.net

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 30908

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Do pathological variables have prognostic significance in rectal adenocarcinoma treated with neoadjuvant chemoradiotherapy and surgery?

Reggiani Bonetti L *et al.* Prognosis of rectal cancer after CRT

Luca Reggiani Bonetti, Simona Lioni, Federica Domati, Valeria Barresi

Luca Reggiani Bonetti, Federica Domati, Department of Laboratory Integrated Activities, Anatomic Pathology and Legal Medicine, Section of Pathology, University of Modena and Reggio Emilia, 41124 Modena, Italy

Simona Lioni, Valeria Barresi, Department of Pathology in Evolutive Age and Adulthood, "Gaetano Barresi", University of Messina, 98125 Messina, Italy

Match Overview

1	Crossref 110 words Barresi, Valeria, Luca Reggiani Bonetti, Federica Domati, and Luigi Baron. "Prognostic relevance of histopathologica ...	2%
2	Internet 98 words crawled on 22-Nov-2016 pdfs.semanticscholar.org	2%
3	Internet 54 words crawled on 13-May-2016 www.wjgnet.com	1%
4	Crossref 37 words M Kikuchi. "High Ki67, Bax, and thymidylate synthase expression well correlates with response to chemoradiation the ...	1%
5	Internet 21 words crawled on 18-Mar-2014 efiloop.com	<1%
6	Crossref 21 words Valeria Barresi. "Diagnostic value of neutrophil gelatinase-associated lipocalin (NGAL) immunoeexpression in follicul ...	<1%
7	Crossref 20 words Barresi, Valeria, Giovanni Branca, Alessandro Raso, Samantha Mascelli, Maria Caffo, and Giovanni Tuccari. "Atypical t	<1%

学术搜索

找到约 8,200 条结果 (用时0.12秒)

文章

我的图书馆

时间不限

2017以来

2016以来

2013以来

自定义范围...

按相关性排序

按日期排序


搜索所有网页

中文网页

简体中文网页

☒ 包括专利

☒ 包含引用

 创建快讯

小提示：只搜索中文(简体)结果，可在 学术搜索设置 指定搜索语言

What is the role for the circumferential margin in the modern treatment of rectal cancer?

[HTML] ascopubs.org

ID Nagtegaal, P Quirke - Journal of Clinical Oncology, 2008 - jco.ascopubs.org
... of diagnostic imaging allowing the accurate prediction of a potentially involved margin will result in ... They did not observe any influence of timing of surgery (day or night), but there ... Although short-course radiotherapy does not decrease the number of positive margins, in patients ...
被引用次数: 634 相关文章 所有 9 个版本 引用 保存

Rates of circumferential resection margin involvement vary between surgeons and predict outcomes in rectal cancer surgery

[PDF] researchgate.net

KF Birbeck, CP Macklin, NJ Tiffin, W Parsons... - ... of surgery, 2002 - journals.lww.com
... journal alerts and information, but is committed to maintaining your privacy and will not share ...
Many of those who do use different denominators in the expression of their results (eg ... Pathologic data specifically relating to the mode of CRM involvement by tumor were collected by a ...
被引用次数: 546 相关文章 所有 8 个版本 引用 保存

Long-term outcome in patients with a pathological complete response after chemoradiation for rectal cancer: a pooled analysis of individual patient data

[HTML] thelancetnorway.com

M Maas, PJ Nelemans, V Valentini, P Das, C Rödel... - The lancet oncology, 2010 - Elsevier
... Article suggestions will be shown in a dialog on return to ScienceDirect. ... adjuvant treatment for patients who respond well, or to intensify treatment for those who do not respond. ... 47 Patients with pCR had clinical T1 and T2 tumours significantly more often than did those without ...
被引用次数: 618 相关文章 所有 17 个版本 引用 保存

Predictive factors of pathologic complete response after neoadjuvant chemoradiation for rectal cancer

[PDF] researchgate.net

MF Kalady, LF de Campos-Lobato, L Stocchi... - ... of surgery, 2009 - journals.lww.com
... you for journal alerts and information, but is committed to maintaining your privacy and will not share ... regression and Cox proportional hazard ratios to determine factors associated with a pathologic complete response. ... Fifty-eight patients (24%) had a pCR and 184 (76%) did not ...
被引用次数: 209 相关文章 所有 7 个版本 引用 保存

Circumferential margin involvement is the crucial prognostic factor after

[HTML] aacrjournals.org



全部

图片

新闻

视频

购物

更多

设置

工具

找到约 448,000 条结果 (用时 0.89 秒)

Prognostic significance of clinical and pathological stages on locally ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者 : B Wen - 2016 - 被引用次数 : 8 - 相关文章

2015年6月4日 - Keywords: Rectal carcinoma, Neoadjuvant chemoradiotherapy, Clinical stage ... chemoradiotherapy has been demonstrated to improve local tumor ... Studies have reported that pathological TNM stage is of prognostic significance in treatment received post-op adjuvant chemotherapy, 55 patients did not.

Pathologic Assessment of Rectal Carcinoma after Neoadjuvant Radio ...

<https://www.hindawi.com/journals/bmri/2015/574540/> - 翻译此页

作者 : M Hav - 2015 - 被引用次数 : 3 - 相关文章

2015年6月14日 - Neoadjuvant radio(chemo)therapy is increasingly used in rectal cancer ... the many facets of rectal carcinoma patient care after neoadjuvant treatment in rectal carcinoma after neoadjuvant RCT did not have significant impact on of surgical and pathologic variables in rectal cancer: a United States ...

Predictive Response Value of Pre- and Postchemoradiotherapy ...

<https://www.hindawi.com/journals/pri/2016/2164609/> - 翻译此页

作者 : MD Santos - 2016 - 被引用次数 : 1 - 相关文章

2015年9月30日 - Neoadjuvant chemoradiotherapy (CRT) followed by total ... The prognostic relevance of pathological tumor response according to ... Radical surgery consisted mainly in sphincter saving rectal significance of pre- and on-treatment variables as predictive factors of Mandard good responders (TRG1-2).

[PDF] Patients with locally advanced rectal cancer treated with neoadjuvant ...

www.spcoloprocto.org/uploads/colo_n2_vol12_2015_artigooriginal2.pdf ▼ 翻译此页

prognostic factors in locally advanced rectal cancer (LARC), treated with neoadjuvant chemoradiotherapy (nCRT) followed ... Mandard TRG and pathological nodal stage were the main ... Digestive Surgery Service, Hospital de Santo António, Centro Hospitalar do Porto ... to have