

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 38348

Manuscript Type: ORIGINAL ARTICLE

Basic Study

Prognostic value of hypoxia-inducible factor-1 α and prolyl 4-hydroxylase beta polypeptide overexpression in gastric cancer

Jun Zhang, Yue Wu, Yu-Hang Lin, Shuai Guo, Pei-Fang Ning, Zhi-Chao Zheng, Yue Wan, Yan Zhao

Abstract

AIM

To investigate the relationship between hypoxia-inducible factor-1 α (HIF-1 α), prolyl 4-hydroxylase beta (P4HR) expression and clinicopathologic

Match Overview

1	Crossref 75 words Li Chen, Yan Shi, Jing Yuan, Yalin Han, Rui Qin, Qian Wu, Baoqing Jia, Bo Wei, Lixin Wei, Guanghai Dai, Shunch	2%
2	Internet 60 words crawled on 09-Feb-2017 doaj.org	2%
3	Internet 16 words crawled on 19-Aug-2017 www.sigmaaldrich.com	<1%
4	Internet 14 words crawled on 31-Mar-2018 stemcellres.biomedcentral.com	<1%
5	Internet 14 words crawled on 24-Jul-2016 www.scribd.com	<1%
6	Crossref 14 words &NA; "Abstracts", Journal of Thoracic Oncology, 06/2011	<1%

找到约 6,160 条结果 (用时 0.40 秒)

Google 学术 : Prognostic value of hypoxia-inducible factor-1 α and prolyl 4-hydroxylase beta polypeptide overexpression in gastric cancer

Hypoxia-inducible factor-1 α is associated with ... - Bos - 被引用次数 : 200

Regulation of angiogenesis by hypoxia-inducible factor ... - Hirota - 被引用次数 : 551

Rapid degradation of hypoxia-inducible factor-1 α by ... - Choi - 被引用次数 : 43

Oncotarget | The miR-124-Prolyl Hydroxylase P4HA1-MMP1 axis ...

www.oncotarget.com/fulltext/2208 ▼ 翻译此页

2014年8月30日 - In this study, we characterized prolyl 4-hydroxylase, alpha polypeptide I (P4HA1) as overexpressed specifically in aggressive prostate cancer. P4HA1 is a ... Hypoxia also induces hypoxia-stabilized HIF1 α (Hypoxia inducible factor-1 α) protein that promotes tumor growth, angiogenesis, and metastasis [13].

The miR-124-Prolyl Hydroxylase P4HA1-MMP1 axis plays a critical ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4196154/> ▼ 翻译此页

作者 : BVSK Chakravarthi - 2014 - 被引用次数 : 32 - 相关文章

2014年7月12日 - In this study, we characterized prolyl 4-hydroxylase, alpha polypeptide I (P4HA1) as overexpressed specifically in aggressive prostate cancer. P4HA1 is a ... Hypoxia also induces hypoxia-stabilized HIF1 α (Hypoxia inducible factor-1 α) protein that promotes tumor growth, angiogenesis, and metastasis [13].

[全部](#)[图片](#)[新闻](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 13,700 条结果 (用时 0.38 秒)

Google 学术: Prognostic value of hypoxia-inducible factor-1 α and prolyl 4-hydroxylase beta polypeptide overexpression in gastric cancer

Hypoxia-inducible factor-1 α is associated with ... - Bos - 被引用次数: 200

Regulation of angiogenesis by hypoxia-inducible factor ... - Hirota - 被引用次数: 551

Rapid degradation of hypoxia-inducible factor-1 α by ... - Choi - 被引用次数: 43

The miR-124-Prolyl Hydroxylase P4HA1-MMP1 axis plays a critical ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4196154/> - 翻译此页

作者: BVSK Chakravarthi - 2014 - 被引用次数: 32 - 相关文章

2014年7月12日 - In this study, we characterized prolyl 4-hydroxylase, alpha polypeptide I (P4HA1) as overexpressed specifically in aggressive prostate cancer. P4HA1 is a ... Hypoxia also induces hypoxia-stabilized HIF1 α (Hypoxia inducible factor-1 α) protein that promotes tumor growth, angiogenesis, and metastasis [13].

Hypoxia inducible factor pathway inhibitors as anticancer therapeutics

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3871878/> - 翻译此页

作者: SK Burroughs - 2013 - 被引用次数: 57 - 相关文章

The hypoxia inducible factor (HIF) pathway is a major mediator of the biological effects of hypoxia in tissues (Figure 1) [9]. ... Under normoxic conditions, HIF- α subunits are hydroxylated by Fe²⁺-dependent HIF prolyl hydroxylases (PHD), mostly PHD2, at two proline residues (402 and 564 in human HIF-1 α) located in the ...

Prolyl hydroxylase and hypoxia inducible factor: potential targets for ...

www.tandfonline.com/doi/full/10.3109/08923973.2010.545418?scroll... - 翻译此页

跳到Role of HIF-1, HIF-1 α in tumor progression - Therefore HIF- α is stabilized and forms a heterodimer with HIF- β which induces the expression of target genes. HIF-1 α over expression in combination with deficiency or mutation of tumor suppressor genes such as VHL, tumor protein 53, and phosphatase and ...

[PDF] Supplementary materials - Dove Medical Press