



**7**  
**Name of Journal:** *World Journal of Gastroenterology*  
**Manuscript NO:** 40814  
**Manuscript Type:** ORIGINAL ARTICLE

*Retrospective Study*  
**Practical fecal calprotectin cut-off value for Japanese patients with ulcerative colitis**

Jun Urushikubo, Shunichi Yanai, Shotaro Nakamura, Keisuke Kawasaki, Risaburo Akasaka, Kunihiro Sato, Yosuke Toya, Kensuke Asakura, Takahiro Gonai, Tamotsu Sugai, Takayuki Matsumoto

### Match Overview

1	<b>Crossref</b> 48 words Anita Bálint, Klaudia Farkas, Zoltán Szepes, Ferenc Nagy et al. "How disease extent can be included in the endo	1%
2	<b>Crossref</b> 47 words Yamaguchi, Sayaka, Yoshiaki Takeuchi, Katsuhito Arai, Katsuyuki Fukuda, Yuichiro Kuroki, Kunio Asonuma, Hiro	1%
3	<b>Crossref</b> 45 words Klaus Theede, Susanne Holck, Per Ibsen, Steen Ladellund, Inge Nordgaard-Lassen, Anette Mertz Nielsen. "Leve	1%
4	<b>Crossref</b> 17 words R MacRedmond. "Screening for lung cancer using low dose CT scanning", Thorax, 3/1/2004	<1%
5	<b>Internet</b> 16 words crawled on 14-Jun-2017 <a href="http://www.wjgnet.com">www.wjgnet.com</a>	<1%
6	<b>Crossref</b> 14 words Ashish Kumar Jha, Madhur Chaudhary, Vishwa Mohan Dayal, Amarendra Kumar et al. "Optimal cut-off value o ...	<1%
7	<b>Internet</b> 14 words crawled on 05-Oct-2017	<1%



Practical fecal calprotectin cut-off value for Japanese patients with ulcera



全部

新闻

图片

视频

购物

更多

设置

工具

找到约 8,130 条结果 (用时 0.33 秒)

### Fecal Immunochemical Test and Fecal Calprotectin Results Show ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5832338/> - 翻译此页

作者: S Hiraoka - 2018 - 被引用次数: 3 - 相关文章

2017年9月7日 - Among **patients** with mucosal healing (MH) and negative FIT or Fcal results ...

Keywords: **Colitis, ulcerative**, Quantitative fecal immunochemical test, **Fecal calprotectin ... Japan**), where the level of calprotectin in the stool specimens was ... Because the **cutoff values** for Fcal varied widely among the different ...

### Fecal immunochemical test as a biomarker for inflammatory bowel ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4754522/> - 翻译此页

作者: J Kato - 2016 - 被引用次数: 19 - 相关文章

2016年1月25日 - Keywords: **Ulcerative colitis**, Crohn disease, Endoscopy, Fecal immunochemical test, **Fecal calprotectin** ... As a pioneer for fecal surrogate makers, tests for **fecal calprotectin** ... In **Japan**, the prevalence of FIT use in clinical **practice** for IBD may be .... The **cutoff values** of Fcal for the presence or absence of ...

### P256 Correlation of fecal calprotectin levels with endoscopic severity ...

[https://academic.oup.com/ecco-jcc/article/11/suppl\\_1/S208/2961199](https://academic.oup.com/ecco-jcc/article/11/suppl_1/S208/2961199) - 翻译此页

作者: K Matsuoka - 2017

2017年1月26日 - Background: **Calprotectin** is a calcium-binding protein that is abundantly ...

Journal of Crohn's and **Colitis**, Volume 11, Issue suppl\_1, ... the same endoscopic parameters (0–3 for **ulcer** size, **ulcerated/affected** surface, ... This **cut-off value** was also evaluated in 31 **patients** who had no large-bowel disease.

### Endoscopic score vs. fecal biomarkers for predicting relapse in ...

<https://www.nature.com> > ... > original contribution - 翻译此页

作者: T Yamamoto - 2018 - 被引用次数: 1 - 相关文章



全部

新闻

图片

购物

视频

更多

设置

工具

找到约 9,580 条结果 (用时 0.56 秒)

### Endoscopic score vs. fecal biomarkers for predicting relapse in ...

<https://www.nature.com> › ... › original contribution - 翻译此页

作者: T Yamamoto - 2018 - 被引用次数: 2 - 相关文章

跳到 **Patients and methods** - Exclusion criteria were: (1) **patients** with **proctitis** alone; (2) **patients** who ... leucocytapheresis (Adacolumn, JIMRO, Takasaki, **Japan**) in 99 **patients** (60%). ... **Fecal calprotectin** was measured by a Human Calprotectin ... In addition, to find an optimal **cutoff value** to predict future relapse, ...

### Clinical Utility of Fecal Calprotectin Monitoring in Asymptomatic ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5434712/> - 翻译此页

作者: A Heida - 2017 - 被引用次数: 20 - 相关文章

2018年6月1日 - We aimed to determine the usefulness of repeated **fecal calprotectin** (FC) ... **Patients** with repeated normal FC **values** had a 67–94% probability to remain in ... (IBD), consisting of Crohn's disease (CD) and **ulcerative colitis** (UC), is a ..... A **practical cut-off** range could be as follows: levels below 250 µg/g as ...

### Fecal immunochemical test as a biomarker for inflammatory bowel ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4754522/> - 翻译此页

作者: J Kato - 2016 - 被引用次数: 19 - 相关文章

2016年1月25日 - Keywords: **Ulcerative colitis**, Crohn disease, Endoscopy, Fecal immunochemical test, **Fecal calprotectin** ... As a pioneer for fecal surrogate makers, tests for **fecal calprotectin** ... In **Japan**, the prevalence of FIT use in clinical **practice** for IBD may be .... The **cutoff values** of Fcal for the presence or absence of ...

### Usefulness of fecal calprotectin for the early prediction of short-term ...

找到约 9,810 条结果 (用时 0.47 秒)

### [Clinical Utility of Fecal Calprotectin Monitoring in Asymptomatic ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5434712/> - 翻译此页

作者: A Heida - 2017 - 被引用次数: 22 - 相关文章

2018年6月1日 - We aimed to determine the usefulness of repeated **fecal calprotectin (FC)** ... **Patients** with repeated normal **FC values** had a 67–94% probability to remain in ... (IBD), consisting of Crohn's disease (CD) and **ulcerative colitis (UC)**, is a ..... A **practical cut-off** range could be as follows: levels below 250 µg/g as ...

### [Fecal Immunochemical Test and Fecal Calprotectin Results Show ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5832338/> - 翻译此页

作者: S Hiraoka - 2018 - 被引用次数: 3 - 相关文章

2017年9月7日 - A total of 110 colonoscopy intervals of **UC patients** were assessed, and ... Keywords: **Colitis, ulcerative**, Quantitative fecal immunochemical test, **Fecal calprotectin ... Japan**), where the level of calprotectin in the stool specimens was ... Because the **cutoff values** for Fcal varied widely among the different ...

### [Endoscopic score vs. fecal biomarkers for predicting relapse in ...](#)

<https://www.nature.com> > ... > **original contribution** - 翻译此页

作者: T Yamamoto - 2018 - 被引用次数: 2 - 相关文章

跳到**Patients and methods** - Exclusion criteria were: (1) **patients** with **proctitis** alone; (2) **patients** who ... leucocytapheresis (Adacolumn, JIMRO, Takasaki, **Japan**) in 99 **patients** (60%). ... **Fecal calprotectin** was measured by a Human Calprotectin ... In addition, to find an optimal **cutoff value** to predict future relapse, ...

### [Usefulness of fecal calprotectin for the early prediction of short-term ...](#)

<https://journals.plos.org/plosone/article?id=10.1371/journal.pone...> ▼ 翻译此页

作者: T Toyonaga - 2017 - 被引用次数: 2 - 相关文章

2017年9月21日 - Background **Fecal calprotectin (FC)** is well accepted as a ... Editor: Emiko Mizoguchi, Kurume University School of Medicine, **JAPAN** ..... to be useful to detect MH, although there is no consensus on the **cut-off value** [15, 16]. .... variability of **fecal calprotectin** levels in **patients** with active **ulcerative colitis**: what ...