

## Match Overview

1	<b>Publications</b> 13 words "Studies from Nagasaki University in the Area of Intercellular Signaling Peptides and Proteins Descri", Health & Medicine	<1%
2	<b>Crossref</b> 13 words Wang, S-C, Y-H Li, H-L Piao, X-W Hong, D Zhang, Y-Y Xu, Y Tao, Y Wang, M-M Yuan, D-J Li, and M-R Du. "PD-1 and Tim	<1%
3	<b>Internet</b> 13 words crawled on 25-May-2016 <a href="#">media.proquest.com</a>	<1%
4	<b>Internet</b> 12 words crawled on 15-Apr-2018 <a href="#">www.nature.com</a>	<1%
5	<b>Internet</b> 12 words crawled on 08-Sep-2017 <a href="#">preview-virologyj.biomedcentral.com</a>	<1%

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 40991

Manuscript Type: ORIGINAL ARTICLE

### *Case Control Study*

Mode of delivery by an ulcerative colitis mother in a case of twins: Immunological differences in cord blood and placenta

Dunsmore G *et al.* Mode of delivery and impacts on the newborn

Garett Dunsmore, Petya Koleva, Reed Taylor Sutton, Lindsay Ambrosio, Vivian Huang, Shokrollah Elahi

### Abstract

#### AIM

To understand the effects of delivery mode on the immune cells frequency and function in cord blood and placenta.

### METHODS



Articles

About 1,370 results (0.10 sec)

Any time

Since 2018

Since 2017

Since 2014

Custom range...

Sort by relevance

Sort by date

☒ include patents☒ include citations☒ Create alert

## Placental transfer of anti-tumor necrosis factor agents in pregnant patients with inflammatory bowel disease

U Mahadevan, DC Wolf, M Dubinsky, A Cortot... - Clinical ..., 2013 - Elsevier

... Breast milk samples were not analyzed by the NMR **method** because it was not validated for quantification ... CZP (Table 4), the last dose was given a median of 19 days before **delivery** (range, 5 ... 2800 g (range, 1956–4110 g), with 5 of the infants defined as low **birth** weight (<2700 ...

☆ Cited by 271 Related articles All 17 versions

## Non-placental causes of intrauterine growth restriction

N Hendrix, V Berghella - Seminars in perinatology, 2008 - Elsevier

... Nucleated red **blood** cells suggest fetal anemia and chronic fetal hypoxia as an important **mechanism** of FGR ... 37. Gastrointestinal diseases, such as Crohn's disease, **ulcerative colitis**, and gastrointestinal bypass surgery, can cause lower **birth** weight because of ...

☆ Cited by 149 Related articles All 7 versions

## Gender aspects of preterm birth

I Ingemarsson - BJOG: An International Journal of Obstetrics & ..., 2003 - Wiley Online Library

... Sex Ratio and **Mode of Delivery** ... The release of catecholamines during labour is an important defence **mechanism** by the hypoxic fetus ... We analysed 15,840 acid–base balances in **cord** artery **blood** at term **delivery** with long-term follow-up of acidotic infants22 ...

☆ Cited by 287 Related articles All 9 versions

## Intrauterine exposure and pharmacology of conventional thiopurine therapy in pregnant patients with inflammatory bowel disease

B Jharap, NKH de Boer, P Stokkers, DW Hommes... - Gut, 2013 - gut.bmj.com

... IBD is associated with an increased risk of lower **birth** weight or preterm **birth**.11 ,12 ... article, these converted metabolite levels (and thus comparable with data generated by the **method** of Lennard ... the end of the first and second trimesters, at **delivery**, and 3 months after **delivery** ...

☆ Cited by 71 Related articles All 13 versions

[HTM

[HTM

[PDF

[全部](#)[图片](#)[视频](#)[购物](#)[新闻](#)[更多](#)[设置](#)[工具](#)

找到约 75,100 条结果 (用时 0.55 秒)

### Placental umbilical cord blood transfusion: a new method of treatment ...

<https://www.ncbi.nlm.nih.gov/pubmed/17089581> ▾ [翻译此页](#)

作者: N Bhattacharya - 2006 - 被引用次数: 17 - [相关文章](#)

**Cord blood**, because of its rich mix of fetal and adult hemoglobin, high platelet ... choice for **cases** of diabetes with severe anemia necessitating blood transfusion. ... after lower uterine cesarean section (LUCS) from consenting **mothers** (56 ml- 138, ... implicated as the cause of scleroderma in **mothers delivering** male babies .

缺少字词: ulcerative colitis twins

### Safety of TNF- $\alpha$ inhibitors during IBD pregnancy: a systematic review

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3734216/> ▾ [翻译此页](#)

作者: OH Nielsen - 2013 - 被引用次数: 80 - [相关文章](#)

2013年7月31日 - However, most presentations were **case** reports or **case** series supplied ... Crohn's disease, IBD, Lactation, Pregnancy, **Ulcerative colitis** ... However, CZP may **differ** from the other tumor necrosis factor ... of pregnancy, Study type, **Mode of delivery**, Outcome of pregnancies .... **Twins**, normal, healthy, full term ...

### Umbilical Cord Complications Publications | PubFacts

<https://www.pubfacts.com/search/Umbilical+Cord+Complications> ▾ [翻译此页](#)

Fetal microchimerism by **mode** of **delivery**: a prospective cohort study. ... Plasma and **Cord Blood** Levels in **Mothers** with Spontaneous Preterm and Term **Delivery**. .... Methods: 20 **cases** of full-term **placenta** without pregnancy complications and .... clinical remission of steroid-dependent or -resistant **ulcerative colitis** patients.

### Can I bank cord blood for twins? - Parent's Guide to Cord Blood

<https://parentsguidecordblood.org/en/faqs/can-i-bank-cord-blood-twins> ▾ [翻译此页](#)

2018年5月4日 - You can preserve **cord blood** from **twins** in a family bank, but not in a public bank. ... However obstetricians are trained to keep track of which baby is which in a multiple **birth**. ... less **cord blood** in the umbilical cord and the **placenta**, and in public ... In the **case** of identical **twins**, the babies have identical DNA.



全部

图片

视频

购物

新闻

更多

设置

工具

找到约 88,900 条结果 (用时 0.61 秒)

## Innate immune function in placenta and cord blood of hepatitis C ...

<https://www.ncbi.nlm.nih.gov/pubmed/20814429> - 翻译此页

作者: CW Hurtado - 2010 - 被引用次数: 41 - 相关文章

2010年8月30日 - Innate immune function in placenta and cord blood of hepatitis ... 80% of cases following exposure, the rate of mother-to-child transmission (2-6%) is strikingly low. ... We hypothesized that innate immune responses would differ across ... In keeping with a model where the maternal-fetal interface provides ...

缺少字词: delivery ulcerative colitis twins

## Placental umbilical cord blood transfusion: a new method of treatment ...

<https://www.ncbi.nlm.nih.gov/pubmed/17089581> - 翻译此页

作者: N Bhattacharya - 2006 - 被引用次数: 17 - 相关文章

Cord blood, because of its rich mix of fetal and adult hemoglobin, high platelet choice for cases of diabetes with severe anemia necessitating blood transfusion. after lower uterine cesarean section (LUCS) from consenting mothers (56 ml-138, implicated as the cause of scleroderma in mothers delivering male babies.

缺少字词: ulcerative colitis twins

## Diseases Treated - Parent's Guide to Cord Blood

<https://parentsguidecordblood.org/en/diseases> - 翻译此页

2018年8月10日 - This website was the first in 2003 to sort diseases treated with cord blood ... it is also important to distinguish between two modes of treatment: ... Leukemia is a cancer of the blood immune system, where the cells .... are defined as follows (in other countries phases may be different): ...

Ulcerative Colitis, ✓ ...

缺少字词: ease twins

## Can I bank cord blood for twins? - Parent's Guide to Cord Blood

<https://parentsguidecordblood.org/en/faqs/can-i-bank-cord-blood-twins> - 翻译此页

2018年5月4日 - You can preserve cord blood from twins in a family bank, but not in a public bank. ...

However obstetricians are trained to keep track of which baby is which in a multiple birth. ... less cord blood in the umbilical cord and the placenta, and in public. In the case of identical twins, the babies