

<sup>9</sup>  
Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 41855

Manuscript Type: ORIGINAL ARTICLE

*Retrospective Study*

Clinicopathological parameters predicting recurrence of pT1N0 esophageal squamous cell carcinoma

Li-Yan Xue, Xiu-Min Qin, Yong Liu, Jun Liang, Hua Lin, Xue-Min Xue, Shuang-Mei Zou, Mo-Yan Zhang, Bai-Hua Zhang, Zhou-Guang Hui, Zi-Tong Zhao, Li-Qun Ren, Yue-Ming Zhang, Xiu-Yun Liu, Yan-Ling Yuan, Jian-Ming Ying, Shu-Geng Gao, Yong-Mei Song, Gui-Qi Wang, Sanford M Dawsey, Ning Lu

### Match Overview

1	Internet 49 words crawled on 31-Jul-2016 <a href="http://www.nature.com">www.nature.com</a>	1%
2	Crossref 38 words Liyang Xue. "Parameters predicting lymph node metastasis in patients with superficial esophageal squamous cell carcinoma"	1%
3	Crossref 37 words Hao-Xian Yang, Xue Hou, Qian-Wen Liu, Lan-Jun Zhang, Jin-Geng Liu, Peng Lin, Jian-Hua Fu. "Tumor Location"	1%
4	Crossref 32 words Shin Saito. "A clinicopathological study of basaloid squamous carcinoma of the esophagus", <i>Esophagus</i> , 09/200	<1%
5	Internet 19 words crawled on 18-May-2018 <a href="http://www.wjgnet.com">www.wjgnet.com</a>	<1%
6	Crossref 15 words Antonio Palumbo, Nathalia Meireles Da Costa, Francesco Esposito, Marco De Martino et al. "HMGA2 overexpression"	<1%
7	Publications 15 words Thakur, B. Zhang, Chun Meng, Xian Bhakta. "Eight-year"	<1%

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 34,500 条结果 (用时 0.57 秒)

CEP55 overexpression predicts poor prognosis in patients with locally ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5244839/> - 翻译此页

作者: W Jiang - 2017 - 被引用次数: 6 - 相关文章

2016年11月22日 - Development of **esophageal squamous cell carcinoma** (ESCC) involves ... clinicopathological staging ( $P=0.002$ ) and tumor **recurrence** ( $P=0.021$ ) in ... **Clinicopathological characteristics** of the 110 patients are listed in Table I.

缺少字词: pttt0

ADAMTS-6 is a predictor of poor prognosis in patients with ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/29475036> - 翻译此页

作者: L Liu - 2018 - 相关文章

2018年2月21日 - ... of poor prognosis in patients with **esophageal squamous cell carcinoma**. ... 0.042), lymph node metastasis ( $P = 0.014$ ) and **recurrence** ( $P = 0.033$ ). ... and other **clinicopathological parameters** including age, sex, tumor size, ...

缺少字词: pttt0

FOXM1: A potential indicator to predict lymphatic metastatic ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6068428/> - 翻译此页

作者: Z Xiao - 2018 - 相关文章

2018年6月6日 - FOXM1 could be a **predictor** of lymphatic metastatic **recurrence** in stage IIA ...  
Keywords: **Esophageal squamous cell carcinoma**, FOXM1, lymphatic ... between FOXM1 expression and **clinicopathological parameters**, and the ...

缺少字词: pttt0

Reiko Masunaga's research works | Shimane University, Matsue ...

[https://www.researchgate.net/scientific.../5032694\\_Reiko\\_Masunaga](https://www.researchgate.net/scientific.../5032694_Reiko_Masunaga) - 翻译此页



全部

图片

新闻

视频

购物

更多

设置

工具

找到约 62,700 条结果 (用时 0.71 秒)

## ADAMTS-6 is a predictor of poor prognosis in patients with ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/29475036> - 翻译此页

作者: L Liu - 2018 - 被引用次数: 1 - 相关文章

2018年2月21日 - ... of poor prognosis in patients with **esophageal squamous cell carcinoma**. ...  
0.042), lymph node metastasis ( $P = 0.014$ ) and **recurrence** ( $P = 0.033$ ). ... and other  
**clinicopathological parameters** including age, sex, tumor size, ...

缺少字词: pT1N0

## Pathological tumor regression grade of metastatic tumors in lymph ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5989742/> - 翻译此页

作者: T Kadota - 2018 - 相关文章

2018年5月3日 - Keywords: chemotherapy-effective rate, **esophageal squamous cell carcinoma**, ...  
primary and metastatic tumors has been implicated in **predicting** prognosis., No residual tumor in ....  
**Recurrence**-free survival curves were estimated using the Kaplan-Meier ... Patients and  
**clinicopathological characteristics**.

缺少字词: pT1N0

## Parameters predicting lymph node metastasis in patients with ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3505024/> - 翻译此页

作者: L Xue - 2012 - 被引用次数: 16 - 相关文章

2012年5月25日 - The relationships between **clinicopathological parameters** and ... Patients with  
superficial **esophageal squamous cell carcinoma** 1) thinner than ...

缺少字词: recurrence pT1N0



找到约 28,200 条结果 (用时 0.49 秒)

### FOXM1: A potential indicator to predict lymphatic metastatic ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6068428/> - 翻译此页

作者: Z Xiao - 2018 - 被引用次数: 1 - 相关文章

2018年6月6日 - FOXM1 could be a predictor of lymphatic metastatic recurrence in stage IIA ...

Keywords: Esophageal squamous cell carcinoma, FOXM1, lymphatic ... between FOXM1 expression and clinicopathological parameters, and the ...

缺少字词: pt1n0

### The crucial role of blood VEGF kinetics in patients with locally ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6102846/> - 翻译此页

作者: YH Chen - 2018

2018年8月20日 - Esophageal squamous cell carcinoma (ESCC) is one of the most fatal human ... Local recurrence, metastatic spread, and resistance to ... Therefore, identifying a robust predictive biomarker or biomarkers to guide .... The clinicopathological parameters of the patients are shown in Table Table11.

缺少字词: pt1n0

### Impact of pretreatment plasma D-dimer levels and its perioperative ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5668066/> - 翻译此页

作者: J Li - 2017 - 被引用次数: 2 - 相关文章

2017年6月16日 - Keywords: esophageal squamous cell carcinoma, D-dimer, surgery, prognosis. Go to: ... In ESCC, plasma D-dimer levels are useful for predicting lymph node metastasis [16]. ...

Relationship between clinicopathological parameters, plasma ... while 38 patients (12.9%) experienced local-regional recurrence.

缺少字词: pt1n0

### Clinicopathological and Prognostic Significance of EGFR Over ...

[https://www.researchgate.net/.../51208803\\_Clinicopathological\\_and\\_Progno...](https://www.researchgate.net/.../51208803_Clinicopathological_and_Progno...) - 翻译此页

2018年5月14日 - ... of EGFR Over-expression in Esophageal Squamous Cell Carcinoma: A ... effects

of EGFR over-expression on clinicopathological parameters ... and it might be useful as a predictive biomarker in clinical practice, ..... clinicopathological features, local recurrence, metastasis status and