

6

Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 51451

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Transumbilical enterostomy for Hirschsprung's disease in two-stage laparoscopy-assisted pull-through

Xu PP *et al.* Transumbilical enterostomy for Hirschsprung's disease

Abstract

BACKGROUND

Since 1998, one-stage transanal endorectal pull-through for treatment of rectosigmoid Hirschsprung's disease (HD) without complications was widely

Match Overview

Match Number	Source	Words	Similarity
1	Crossref	37 words	1%
	Yang, Li, Shao-tao Tang, Shuai Li, T.H. Aubdoollah, Guo-qin g Cao, Hai-yan Lei, and Xin-xing Wang. "Two-stage laparosc		
2	Internet	23 words	1%
	crawled on 27-Dec-2016 office.wjgnet.com		
3	Crossref	17 words	1%
	Aubdoollah, Tajammool Hussein. "Clinical outcomes and e... onomics analysis of three laparoscopic techniques for Hirsch		
4	Crossref	17 words	1%
	M.-M. Garcia-Barcelo. "Is there a role for the IHH gene in H... schsprung's disease?", Neurogastroenterology and Motility.		
5	Crossref	14 words	<1%
	Xi Zhang, Guo-qing Cao, Shao-tao Tang, Xiao-pan Chang, Sh uai Li, Li Yang, Kang Li, Ying Zhou, De-hua Yang. "Laparosc		
6	Internet	14 words	<1%
	crawled on 28-Aug-2019 f6publishing.blob.core.windows.net		
7	Internet	12 words	<1%
	crawled on 15-Jun-2019 csd2018.meetingchina.org		



国内版 国际版

Transumbilical enterostomy for Hirschsprung's disease in two-



Chat with Bing

Sign in



All Images Videos

关闭取词

Add Bing Firefox extension

1,920 Results Any time

Laparoscopic-Assisted Pull-Through for Hirschsprung's ...

<https://obgynkey.com/laparoscopic-assisted-pull-through-for-hirschsprungs-disease>

Oct 25, 2017 · Fig. 36.1. Contrast enema post evacuation phase demonstrating the transition zone (arrows). The aganglionic Hirschsprung's bowel is distal to the transition zone (TZ—transition zone), which is less than 5 cm in length in most cases. Treatment Traditionally, surgical treatment for Hirschsprung's disease was performed in two or even three stages.

Primary Laparoscopic-Assisted Endorectal Colon Pull ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1420812>

To describe the surgical technique and early clinical results after a one-stage laparoscopic-assisted endorectal colon pull-through for Hirschsprung's disease. Summary Background Data Recent trends in surgery for Hirschsprung's disease have been toward earlier repair and fewer surgical stages.

Cited by: 273 Author: Keith E. Georgeson, Roger D. Cohen, An...

Publish Year: 1999

One-stage laparoscopy-assisted endorectal pull-through for ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4643354>

Children with late-presenting Hirschsprung's disease (HD) are classically treated by a staged operation with enterostomy. An alternative may be one-stage laparoscopy-assisted endorectal pull-through, which has cosmetic advantages. This case-series report describes the outcomes of children with ...

Author: So Hyun Nam, Min Jeong Cho, Dae Y... Publish Year: 2015

One-stage laparoscopy-assisted endorectal pull-through for ...

europepmc.org/articles/PMC4643354

Children with late-presenting Hirschsprung's disease (HD) are classically treated by a staged operation with enterostomy. An alternative may be one-stage laparoscopy-assisted endorectal pull-through, which has cosmetic advantages. This case-series report describes the outcomes of children with late-presenting HD who underwent this procedure.

Author: So Hyun Nam, Min Jeong Cho, Dae Y... Publish Year: 2015

One-stage laparoscopy-assisted endorectal pull-through for ...

<https://www.sciencedirect.com/science/article/pii/S2210261215003958>

The operation of Hirschsprung's disease in children older than 3 years is traditionally staged operation with enterostomy because of severe dilated bowel. • We did one stage laparoscopy-assisted endorectal

3,610 Results Any time

Laparoscopic-Assisted Pull-Through for Hirschsprung's Disease

<https://obgynkey.com/laparoscopic-assisted-pull-through-for-hirschsprungs-disease>

Oct 25, 2017 · Fig. 36.1. Contrast enema post evacuation phase demonstrating the transition zone (arrows). The aganglionic Hirschsprung's bowel is distal to the transition zone (TZ—transition zone), which is less than 5 cm in length in most cases. Treatment Traditionally, surgical treatment for Hirschsprung's disease was performed in two or even three stages.

Primary Laparoscopic-Assisted Endorectal Colon Pull ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1420812>

To describe the surgical technique and early clinical results after a one-stage laparoscopic-assisted endorectal colon pull-through for Hirschsprung's disease. Summary Background Data Recent trends in surgery for Hirschsprung's disease have been toward earlier repair and fewer surgical stages.

Cited by: 275 Author: Keith E. Georgeson, Roger D. Cohen, An...

Publish Year: 1999

One-stage laparoscopy-assisted endorectal pull-through for ...

<https://www.sciencedirect.com/science/article/pii/S2210261215003958>

The operation of Hirschsprung's disease in children older than 3 years is traditionally staged operation with enterostomy because of severe dilated bowel. • We did one stage laparoscopy-assisted endorectal pull-through with short segment Hirschsprung's disease for late presented children. •

Author: So Hyun Nam, Min Jeong Cho, Dae Y... Publish Year: 2015

Laparoscopic-Assisted Pull-Through for Hirschsprung's Disease

https://www.researchgate.net/publication/11056174_Laparoscopic-Assisted_Pull-Through...

The purpose of this report is to evaluate the cost-effectiveness of a single-stage laparoscopic pull-through for Hirschsprung's disease compared to the traditional two-stage Duhamel procedure.

Author: Keith E Georgeson

Duhamel versus transanal endorectal pull through (TERPT ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5040736>

Sep 14, 2016 · For the surgical treatment of Hirschsprung's disease, several surgical techniques are used to resect the distal aganglionic colon. Two frequently used techniques are the Duhamel procedure and the transanal endorectal pull-through procedure. During the '8th Pediatric Colorectal Course' in ...

Cited by: 3 Author: E. Arts, S. M. B. I. Botden, M. Lacher, P...



Get quick access to Bing with the Bing extension for Firefox

MAYBE LATER

YES



4,450 Results Any time ▾

Laparoscopic-Assisted Pull-Through for Hirschsprung's Disease

<https://obgynkey.com/laparoscopic-assisted-pull-through-for-hirschsprungs-disease> ▾

Oct 25, 2017 · Fig. 36.1. Contrast enema post evacuation phase demonstrating the transition zone (arrows). The aganglionic **Hirschsprung's** bowel is distal to the transition zone (TZ—transition zone), which is less than 5 cm in length in most cases. Treatment Traditionally, surgical treatment for **Hirschsprung's disease** was performed in two or even three stages.

One-stage laparoscopy-assisted endorectal pull-through for ...

<https://www.sciencedirect.com/science/article/pii/S2210261215003958>

The operation of **Hirschsprung's disease** in children older than 3 years is traditionally staged operation with **enterostomy** because of severe dilated bowel. • We did one stage laparoscopy-assisted endorectal pull-through with short segment **Hirschsprung's disease** for late presented children. •

Author: So Hyun Nam, Min Jeong Cho, Dae Yeo... **Publish Year:** 2015

Analysis of the Costs of Surgery for Hirschsprung's ...

<https://journals.sagepub.com/doi/abs/10.1177/000992289903801004?related-urls=yes138/10...>

Oct 01, 1999 · The purpose of this report is to evaluate the cost-effectiveness of a single-stage laparoscopic pull-through for **Hirschsprung's disease** compared to the traditional **two-stage** Duhamel procedure. In this series of 33 children, the length of hospitalization (2.5 ± 3.5 vs 10.6 ± 3.9 , $p < 0.01$), cost ($19,088 \pm 13,075$ vs $34,110 \pm 19,443$, $p < 0.05$), and complications were all significantly less with ...

Cited by: 32 **Author:** Anthony J. Bufo, Mike K. Chen, Rasik Shah,...

Publish Year: 1999

One-stage laparoscopy-assisted endorectal pull-through for ...

europepmc.org/articles/PMC4643354

Children with late-presenting **Hirschsprung's disease** (HD) are classically treated by a staged operation with **enterostomy**. An alternative may be one-stage laparoscopy-assisted endorectal pull-through, which has cosmetic advantages. This case-series report describes the outcomes of children with late-presenting HD who underwent this procedure.

Author: So Hyun Nam, Min Jeong Cho, Dae Yeo... **Publish Year:** 2015

Laparoscopic-Assisted Pull-Through for Hirschsprung's ...

https://www.researchgate.net/publication/311944213_Laparoscopic-Assisted_Pull-Through...

Request PDF | Laparoscopic-Assisted Pull-Through for **Hirschsprung's Disease** | **Hirschsprung's disease** (congenital megacolon) is a congenital disorder of the intestine. It is caused by ...