

ALL IMAGES VIDEOS

9,410 Results Any time

Special Issue "Endoscopic Diagnosis and Treatment of Early ...
<https://www.mdpi.com/journal/cancers/special...>

Dear Colleagues, The most common form of gastric cancer develops through a well-established precancerous inflammatory cascade pathway. The first stage, active chronic inflammation, progresses from non-atrophic gastritis to multifocal atrophic gastritis, gastric intestinal metaplasia, dysplasia, and eventually early and finally invasive cancer.

Endoscopic Diagnosis and Management of Gastrointestinal ...
<https://www.cancerjournal.com/view/endoscopic...>

Oct 01, 1995 - Tada et al reported a 5-year survival rate of 83.9% in 82 patients with early gastric cancer who were treated by strip biopsy, as compared with a rate of 89.3% in 27 patients who underwent surgical treatment [37]. Endoscopic treatment may be appropriate for selected patients with superficial gastric cancer who are unable or unwilling to undergo surgery, for focal areas of severe dysplasia, and for the rare gastric sessile villous adenoma.

PEOPLE ALSO ASK

- How is endoscopy used to diagnose gastric cancer? ▾
- When to have an endoscopy for low grade dysplasia? ▾
- How is endoscopic imaging used to diagnose GC? ▾
- What do you need to know about gastric metaplasia? ▾


See results for

Barrett's Esophagus (Medical Condition)

Abnormal change of the cells present in the lower portion of the esophagus (food pipe) due to acid reflux, which causes the lining to thicken and infla...

Search Tools

Turn off Hover Translation (关闭悬停)



Name of Journal: *World Journal of Gastroenterology*

Manuscript NO: 66627

Manuscript Type: REVIEW

Endoscopic diagnosis and treatment of gastric dysplasia and early cancer: current evidence and what the future may hold

Endoscopy in gastric neoplasia

Edward Young, Hamish Philpott, Rajvinder Singh

Match Overview

1	Crossref 18 words Soo Juy Phne, Nagashwar Reddy, Philip W Y Chiu, Pradeep Rabata et al. "Robot-Assisted Endoscopic Submucosal Dis...	<1%
2	Internet 18 words crawled on 16-Jul-2020 link.springer.com	<1%
3	Internet 13 words crawled on 18-Feb-2019 eprints.whiterose.ac.uk	<1%
4	Crossref 12 words "Endoscopy in Early Gastrointestinal Cancers, Volume 2", Springer Science and Business Media LLC, 2021	<1%
5	Crossref 12 words Hyun Ju Kim, Hyunsoo Chung, Da Hyun Jung, Jun Chul Park, Sung Kwan Shin, Sang Kil Lee, Yong Chan Lee. "Clinical outc...	<1%
6	Internet 12 words crawled on 21-May-2020 www.epg.pt	<1%

国内版

国际版

Endoscopic diagnosis and treatment of gastric dysplasia and early



ALL

IMAGES

VIDEOS

9,470 Results

Any time ▾

Special Issue "Endoscopic Diagnosis and Treatment of Early ...

<https://www.mdpi.com/journal/cancers/special...> ▾

Dear Colleagues, The most common form of gastric cancer develops through a well-established precancerous inflammatory cascade pathway. The first stage, active chronic inflammation, progresses from non-atrophic gastritis to multifocal atrophic gastritis, gastric intestinal metaplasia, dysplasia, and eventually early and finally invasive cancer.

Endoscopic Diagnosis and Management of Gastrointestinal ...

<https://www.cancernetwork.com/view/endoscopic...> ▾

Oct 01, 1995 · Tada et al reported a 5-year survival rate of 83.9% in 82 patients with early gastric cancer who were treated by strip biopsy, as compared with a rate of 89.3% in 27 patients who underwent surgical treatment [37]. Endoscopic treatment may be appropriate for selected patients with superficial gastric cancer who are unable or unwilling to undergo surgery, for focal areas of severe dysplasia, and for the rare gastric sessile villous adenoma.

PEOPLE ALSO ASK

How is endoscopy used to diagnose gastric cancer? ▾

When to have an endoscopy for low grade dysplasia? ▾

How is endoscopic imaging used to diagnose GC? ▾

What do you need to know about gastric metaplasia? ▾

Feedback

AGA Clinical Practice Update on Endoscopic Treatment of ...

[https://www.gastrojournal.org/article/S0016-5085\(19\)41537-0/fulltext](https://www.gastrojournal.org/article/S0016-5085(19)41537-0/fulltext)

Nov 12, 2019 · After complete eradication (endoscopic and histologic) of intestinal metaplasia has been achieved with BET, surveillance endoscopy with biopsies should be performed at the following intervals: for baseline diagnosis of HGD/esophageal adenocarcinoma: at 3, 6, and 12 months and annually thereafter; and baseline diagnosis of low-grade dysplasia: at 1 and 3 years.

Cited by: 61

Author: Prateek Sharma, Prateek Sharma, Nichol...