

国内版国际版

Microsoft Bing

Relationship between Non Alcoholic Fatty liver disease and chronic

Chat with Bing

Sign in

ALLIMAGESVIDEOS

68,200 ResultsAny time


Relationship between non-alcoholic fatty liver disease and ...
<https://pubmed.ncbi.nlm.nih.gov/23432982>
Non-alcoholic fatty liver disease (NAFLD) is now regarded as hepatic component of the metabolic syndrome. In addition, NAFLD has emerged as a growing public health problem worldwide and an important challenge for health authorities. NAFLD is associated with insulin resistance and hyperlipidemia and ...
Cited by: 21Author: Asma A. Hamad, Abf A. Khalil, Vincent C...
Publish Year: 2012

Relationship between non-alcoholic fatty liver disease and ...
<https://www.sciencedirect.com/science/article/pii/S1687197912000858>
Dec 01, 2012. The possible link between non-alcoholic fatty liver disease (NAFLD) and kidney disease is subject for considerable research interest. NAFLD is emerging as an important public health problem across the globe with an estimated prevalence of 20–30% in Western communities and 90% in morbidly obese [2], [3].
Cited by: 21Author: Asma A. Hamad, Abf A. Khalil, Vincent C...
Publish Year: 2012

Non-alcoholic fatty liver and chronic kidney disease ...
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8108029>
May 07, 2021. INTRODUCTION: Non-alcoholic fatty liver disease (NAFLD) encompasses a spectrum of chronic liver disease ranging from steatosis on one end to fibrosis and cirrhosis on the other end[1]. NAFLD and non-alcoholic steatohepatitis (NASH) are the hepatic manifestations of metabolic syndrome (MetS), which is a disease form for a multitude of comorbidities, such as insulin resistance,

Non-Alcoholic Fatty Liver Disease

Medical Condition



A condition in which fat accumulates in the liver in people who drink little or no alcohol.

Very common (More than 3 million cases per year in US)

Requires lab test or imaging

Treatments can help manage condition, no known cure

Can last several years or be lifelong

国内版

国际版

Relationship between non-alcoholic fatty liver disease and chronic



ALL

IMAGES

VIDEOS

35,800 Results

Any time ▾

CKD and Nonalcoholic Fatty Liver Disease | Request PDF

<https://www.researchgate.net/publication/264430796...>

Background: Non-alcoholic fatty liver disease (NAFLD) is a risk factor for the development of chronic kidney disease (CKD), which is early marked by kidney glomerular hyperfiltration.

Guidelines on the management of ascites in cirrhosis

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1860002>

Apr 28, 2006 · Mortality from cirrhosis has increased from 6 per 100 000 population in 1993 to 12.7 per 100 000 population in 2000. 7 Approximately 4% of the general population have abnormal liver function or liver disease 8 and approximately 10–20% of those with one of the three most common chronic liver diseases (non-alcoholic fatty liver disease ...

Cited by: 601

Author: K P Moore, G P Aithal

Publish Year: 2006

PEOPLE ALSO ASK

How is non alcoholic fatty liver disease different from alcohol?



How are diabetes and obesity related to fatty liver disease?



Is there a link between Nash and fatty liver disease?



What is the diet for nonalcoholic fatty liver disease?



Feedback

Association between Nonalcoholic Fatty Liver Disease and ...

<https://www.hindawi.com/journals/cjgh/2021/6678142> ▾

Jan 11, 2021 · Nonalcoholic fatty liver disease (NAFLD) has a rising prevalence worldwide. Its potential for evolution towards liver cirrhosis and hepatocellular carcinoma, as well as associations with extrahepatic manifestations, represents a double burden for patients and physicians alike. Recently, there has been increasing evidence of the association between NAFLD and a number of endocrinopathies, ...

Author: Ana-Maria Singeap, Carol Stanciu, La...

Publish Year: 2021