

69470\_Auto\_EditedC.docx

**Name of Journal:** *World Journal of Gastroenterology*

**Manuscript NO:** 69470

**Manuscript Type:** REVIEW

## **Epidemiology of stomach cancer**

Ilic M *et al.* Stomach cancer

Milena Ilic, Irena Ilic

### **Abstract**

Despite a decline in incidence and mortality during the last decades, stomach cancer is one of the main health challenges worldwide. According to the GLOBOCAN 2020 estimates, stomach cancer caused about 800000 deaths (accounting for 7.7% of all cancer deaths), ranking as the fourth leading cause of cancer deaths in both genders combined. About 1.1 million new cases of stomach cancer were diagnosed in 2020 (accounting for 5.6% of all cancer cases). About 75% of all new cases and all deaths from stomach cancer are reported in Asia. Stomach cancer is one of the most lethal malignant tumours, with the five-year survival rate of about 20%. There are some well-established risk factors for stomach cancer: *Helicobacter pylori* infection, dietary factors, tobacco, obesity, radiation. Until today, the most important way for prevention of stomach cancer is reduction of exposure to risk factors, as well as screening and early detection. Further research of risk factors can help identify various opportunities for a more effective prevention. Screening program for stomach cancer has been implemented in a few countries, either as a national or opportunistic screening of high-risk individuals only. Generally, due to high aggressiveness and heterogeneity, stomach cancer still remains a severe global health problem.

**Key Words:** Stomach cancer; Epidemiology; Incidence; Mortality; Survival; Predictive factors; Prevention.

<sup>1</sup> Ilic M, Ilic I. Epidemiology of stomach cancer. *World J Gastroenterol* 2022; In press

**Core Tip:** Despite the <sup>3</sup> decline in incidence and mortality during the last decades, stomach cancer is one of main health challenges worldwide. According to the GLOBOCAN 2020 estimates, stomach cancer caused about 800000 <sup>1</sup> deaths, ranking as the fourth <sup>9</sup> leading cause of deaths from cancer in both genders combined. Around 1.1 million new stomach cancer cases were diagnosed in 2020. There are some well-established <sup>1</sup> risk factors for stomach cancer: *Helicobacter pylori* infection, dietary factors, tobacco, obesity, radiation. Until today, the most important way for prevention of stomach cancer is reduction of exposure to risk factors, as well as screening and early detection.

## INTRODUCTION

Stomach cancer was the fifth most common malignant tumor in the world in 2020 with approximately 1.1 million new cases, and is the fourth leading cause of cancer death, with about 800000 deaths<sup>[1,2]</sup>. Over 85 percent of stomach cancer cases are registered in countries with high and very high Human Developing Index (590000 and 360000 cases, respectively)<sup>[1]</sup>. The highest number of cases of stomach cancer (almost 820000 new cases and 580000 deaths) was registered in Asia (mainly in China)<sup>[1,2-4]</sup>. The estimated five-year survival rate is lower than 20%<sup>[2,6-8]</sup>.

Worldwide, stomach cancer incidence and mortality correlate with increasing age and are relatively rare in persons of both gender younger than 45<sup>[2,4,7]</sup>. Frequency of stomach cancer in men is approximately double to that in women<sup>[1-3]</sup>. In men, stomach cancer was the most commonly diagnosed cancer in 2020 in seven countries (all countries were in Asia: Iran, Afghanistan, Turkmenistan, Uzbekistan, Tajikistan, Kyrgyzstan, and Bhutan) and the leading cause of death from cancer in ten countries (Iran, Afghanistan, Tajikistan, Kyrgyzstan, and Bhutan in Asia, Mali and Cape Verde in Africa, Colombia and Peru in South America, Costa Rica in Central America)<sup>[1,2]</sup>. Although stomach cancer was not the most diagnosed cancer in women in any country, stomach cancer was the leading cause of death from cancer among females in three countries (Tajikistan, Bhutan, and Peru)<sup>[1,2]</sup>. Incidence and mortality rates from stomach cancer were generally low in Northern America and Northern Europe in 2020 and equivalent to rates registered across most of the African regions<sup>[1,2]</sup>.

In the first half of the 20<sup>th</sup> century, gastric cancer was the leading cause of death from malignant tumours in the United States and Europe<sup>[9,10]</sup>. Over the past decades incidence and mortality of stomach cancer have substantially declined in many countries<sup>[1,2]</sup>.

Stomach cancer is a multifactorial disease<sup>[9-12]</sup>, including both lifestyle and environmental risk factors [*Helicobacter pylori* (*H. pylori*) infection, low socioeconomic status, dietary factors, such as high intake of salty and smoked food and low consumption of fruits and vegetables, fiber intake, tobacco smoking, alcohol use, low

physical activity, obesity, radiation], gastroesophageal reflux disease, positive family history and inherited predisposition. Besides, the etiology of stomach cancer has not yet been sufficiently elucidated.

Topographically, stomach cancer is classified into two subsites: the cardia stomach cancer (arising from the upper stomach) and noncardia stomach cancer (arising from the other parts of the stomach), which differ in epidemiologic patterns and etiology<sup>[13]</sup>. Majority of all stomach cancers (approximately 90%) are adenocarcinomas, while other types (including lymphoma, sarcoma, neuroendocrine tumors) are rare<sup>[12,14]</sup>. Two major histologic types of stomach cancer adenocarcinomas are diffuse and intestinal, which differ in epidemiological peculiarities, such as age at diagnosis, gender ratio, *etc.*<sup>[15,16]</sup>.

Despite the strong declining trends in incidence and mortality, stomach cancer remains an important part of the global burden of cancer. Many of the risk factors remain insufficiently understood and need to be the focus of further research in order to achieve more specific, targeted measures of prevention.

## **INCIDENCE**

Worldwide, there is a considerable geographic variation in stomach cancer incidence. Stomach cancer incidence rates in 2020 were the highest in Eastern Asia (22.4 per 100000 people), followed by Central and Eastern Europe (11.3 per 100000 people), and South America, Polynesia and Western Asia (equally about 8.6 per 100000 people) (Figure 1A)<sup>[2]</sup>. The lowest rate (3.3 per 100000 people) was registered in Southern Africa.

More than three quarters (75.3%; 819944) of all stomach cancer cases are residents of Asia<sup>[2]</sup>. Most (86.7%; 944591 cases) stomach cancer cases were residents of more developed regions. The least number of stomach cancer cases were recorded in Micronesia/Polynesia.

For incidence of stomach cancer in 2020, as well as for mortality, notable international variations existed around the world (Figures 1-4)<sup>[2]</sup>. The highest incidence rates were recorded in countries of eastern Asia (Mongolia, Japan, Republic of Korea), while the highest death rates were observed in countries of western Asia (Tajikistan, Kyrgyzstan,

Iran). The lowest rates of incidence and mortality of stomach cancer were recorded in Northern America and Northern Europe, Australia/New Zealand and some African countries. Patterns in females were broadly similar to those observed in males, but large differences were observed between sexes and throughout different countries/regions (Figures 1-4)<sup>[2]</sup>. Globally, the incidence rate of stomach cancer in males was 15.8 per 100000 in 2020, and in females 7.0 per 100000 (Figure 1B)<sup>[2]</sup>. The gastric cancer incidence rates were about 2 to 3 times higher in males than in females (ranging from 32.5 per 100000 in Eastern Asia to 4.5 per 100000 in Middle Africa for men, and in women ranging from 13.2 in Eastern Asia to 2.4 in Southern Africa) (Figure 1B)<sup>[2]</sup>. By countries, the differences were fiftyfold: the incidence rates of gastric cancer in men ranged from 48.1 per 100000 in Japan to 1.0 per 100000 in Mozambique in 2020 (Figure 2A). Also, similar differences were observed by regions: the highest incidence rates were reported in Eastern Asia (Japan: 48.1, Mongolia: 47.2, Republic of Korea: 39.7), while the lowest rates were recorded in South Africa (Mozambique: 1.0, Lesotho: 2.1). The incidence rates of stomach cancer in women ranged from 20.7 per 100000 inhabitants in Mongolia (followed by Tajikistan: 18.7, Republic of Korea: 17.6 and Japan: 17.3) to about 0.5 in Indonesia and Mozambique in 2020 (Figure 2B).

However, the distribution of stomach cancer did not have a clear geographical pattern: namely, even though the highest risk populations in the world are in Asian countries (e.g. Japan, Mongolia, Republic of Korea), some other countries in Asia register relatively low rates (such as Sri Lanka, Indonesia, Thailand) (Figure 2A and B)<sup>[2]</sup>. On the other hand, in some low-risk populations, there are some high-risk groups for stomach cancer, such as Koreans and Japanese who live in the United States<sup>[17,18]</sup>.

Also, rates varied across races. The stomach cancer incidence in men in the United States was the highest in blacks, followed by Asians/Pacific Islanders, Hispanics, and American Indian/Alaska natives<sup>[7]</sup>. In women, the highest rates were registered in Hispanics, followed by blacks and Asian/Pacific Islanders, and American Indian/Alaska natives. In the United States, for both sexes, the lowest rates were recorded in whites.



Also, incidence and mortality of gastric cancer in all indigenous groups exceeded the frequency among their non-indigenous counterparts: the highest gastric cancer rates were registered in Indigenous Siberians, Mapuche in Chile and among Alaskan Inuit<sup>[19]</sup>. Additionally, increasing incidence trends were observed in some indigenous groups, especially in Inuit residing in the circumpolar region and in Maori in New Zealand.

Although differences in stomach cancer incidence in different parts of the world are still not fully clear, most of the variation in stomach cancer incidence worldwide is due to variations in exposure to environment or lifestyle related risk factors<sup>[20-22]</sup>. Additionally, migrant studies<sup>[23]</sup> and secular trends of gastric cancer rates also indicate that environmental factors have an important role in etiology of gastric cancer<sup>[24]</sup>. The most important established risk factor for gastric cancer is infection with *H. pylori*<sup>[25]</sup>. Internationally, variations in *H. pylori* infection prevalence show similarities with variations in stomach cancer prevalence; in developing countries, *H. pylori* infection prevalence in adults is 76% vs 58% in developed countries<sup>[26]</sup>. The prevalence was estimated to be 77.6% in South Africa, 55.8% in China, 52.2% in Mexico, 24.6% in Australia and 22.1% in Denmark<sup>[27]</sup>. In the United States of America, prevalence in non-Hispanic blacks was 53%, in Mexican Americans - 62%, but 26% among non-Hispanic whites<sup>[28]</sup>. In part, geographical variation of *H. pylori* infection rates correlate with the frequency of stomach cancer across populations. On the other hand, certain highly infected populations (e.g. in Africa and South Asia), unlike the East Asian countries, do not have a high incidence of stomach cancer, which can be explained, at least in part, by the differences in prevalence of genotypes of *H. pylori* (in East Asian the vacA m1 genotype is predominant, whereas the m2 genotype predominated in Africa, South Asia, and Europe)<sup>[29]</sup>.

Additionally, several other environmental factors are also considered as contributors to gastric cancer occurrence<sup>[21,22,30]</sup>. Differences between sexes and international variations could likely be due to tobacco smoking<sup>[31]</sup>. Based on the Global Burden of Disease study<sup>[22]</sup>, the drop in burden of stomach cancer was associated with improved

Socio-demographic Index, then to high-sodium diet in both genders combined, as well as to smoking in males, in particular in east Asian populations.

In addition, some research points to the role of aging<sup>[24]</sup> and hereditary and genetic factors<sup>[6]</sup> in stomach cancer burden. Incidence differences by sex have never been fully explained, but some theories have suggested a protective role of female sex-specific hormones<sup>[15,32]</sup>. A higher stomach cancer incidence in males than in females may be due to differences in the incidence of different subtypes of adenocarcinoma according to histology (intestinal or diffuse) and location (proximal or distal)<sup>[12,13,33]</sup>. Diffuse adenocarcinoma is more common in younger and female patients, whereas intestinal adenocarcinoma is more common in males and the elderly<sup>[15]</sup>. Intestinal adenocarcinoma dominates high-risk areas and is considered responsible for much of the international variation in incidence. The observed differences in stomach cancer incidence worldwide could be due to diagnostic capacity and changes in the quality of registries, which coverage, completeness and accuracy vary by country<sup>[34]</sup>.

## **MORTALITY**

Nearly three quarters of stomach cancer deaths (74.8%; 575206 deaths) were registered in Asia<sup>[2]</sup>. Most (83.7%, 643609 deaths) of all deceased from stomach cancer were residents of more developed regions. The least number of deaths were recorded in Micronesia/Polynesia.

The mortality of stomach cancer varies greatly across populations and regions. Mortality rates for stomach cancer in 2020 in both genders were the highest in Eastern Asia region (14.6 per 100000 people), followed by South America, Polynesia, Western Asia and Central and Eastern Europe (equally about 8.5 per 100000 people) (Figure 1A)<sup>[2]</sup>. The lowest mortality rates (about 2.0 per 100000 people) were registered in Northern Africa and Australia. The differences in mortality rates were thirtyfold between the population with the highest rate (Mongolia - 24.6), and the one with the lowest rate (Mozambique - 0.7).



Stomach cancer mortality by gender shows significant geographic variations<sup>[1,2,6-8]</sup>. Globally, the mortality rate of stomach cancer in males in 2020 was 11.0 per 100000, and in females 4.9 per 100000 (Figure 1C)<sup>[2]</sup>. The region with the highest mortality rates of stomach cancer in 2020 in both genders was Eastern Asia (21.3 and 8.8 per 100000, respectively) (Figure 1C)<sup>[2]</sup>. The lowest rates of stomach cancer mortality in both sexes were in North America (2.3 and 1.3 per 100000, respectively). In men, the risk of dying from stomach cancer was the highest in Mongolia (36.5), then in Kyrgyzstan, Tajikistan and China (approximately 25.0 per 100000) (Figure 2C). Contrary to that, the risk of death from stomach cancer was the lowest in men in Mozambique (1.0) and Indonesia (1.9). Women living in Tajikistan and Mongolia had the greatest risk (approximately 15.0 per 100000) of death from stomach cancer, while the risk for women in Indonesia and Mozambique was the lowest (less than 1.0 per 100000) (Figure 2D).

Mortality rates of gastric cancer begin to rise in middle-aged persons, with the highest rates observed in the elderly (aged 75 years and older) age group for both males and females (Figure 4B).

Stomach cancer mortality showed apparent geographical variability. Generally, the large differences for mortality rates come up between developing and developed countries. Considering developed countries, this mortality pattern could be explained by the increased hygiene standards, dissemination of food refrigeration, better conservation of food, high intake of fresh fruits and vegetables and eradication of *H. pylori*<sup>[11,22,35]</sup>. In the second decade of the 21<sup>st</sup> century in Japan, mortality of stomach cancer reached the levels of Western countries (Figure 4B), which could be attributed to introduction of screening for gastric cancer screening and to changes in lifestyle, such as the reduction in salt use and an increase in the use of fresh fruits and vegetables, improvement of food storage, smoking reduction and prevention of infection with *H. pylori*<sup>[22,36,37]</sup>. But, reasons for the significant international variations in stomach cancer mortality rates are not fully clear. Diffuse adenocarcinoma is more common in females, while intestinal adenocarcinoma is dominant in males, the subtype being responsible for most of the international variations<sup>[15]</sup>.

There is a wide variation in the relative contribution of cardia and noncardia cancers to the overall number of stomach cancer cases, with a higher proportion of cardia cancers in countries with lower stomach cancer incidence and mortality rates (such as the United States, Canada and Denmark)<sup>[38]</sup>. In males in Europe, the proportion of cardia and noncardia stomach cancers ranged between 11.6% (Belarus) and 72.0% (Finland), with higher proportions observed in Northern Europe and lower in Eastern Europe. Among other countries worldwide, the proportion of cardia stomach cancers ranged between 5.8% (Republic of Korea) and 64.8% (Iran). In females, a similar geographic pattern was observed, although rates were lower: in Europe, the proportion of cardia and noncardia stomach cancers ranged between 10.6% (Italy) and 44.5% (United Kingdom), while worldwide it ranged from 4.3% (Republic of Korea) to 31.5% (Australia).

Low incidence rates of stomach cancer, which notably became a rare diagnosis among the white United States population, are attributed to the “unplanned triumph” of prevention, which involves a decreased *H. pylori* prevalence and improved food storage and preservation<sup>[9,39]</sup>.

Cancer mortality data are influenced by data on incidence, as well as the success of treatment. Although World Health Organization estimates present detailed and high-quality information on the incidence and mortality of stomach cancer recorded by cancer registries (regional or national) around the world, these estimates should be interpreted with considerable caution, due to the limited quality and coverage of cancer data worldwide, especially in low- and middle-income countries, due to issues of local data quality, registry coverage, and analytical capacity<sup>[2,40,41]</sup>. The effect of coronavirus disease 2019 pandemic on cancer burden is not clear yet, particularly taking into consideration the geographical variations and evolution of the pandemic across countries (because of the lockdown, possible delays in cancer diagnoses, etc.)<sup>[2]</sup>

The differences in availability of improvements in stomach cancer diagnosis and treatment may have had some role in the observed variations in mortality rates worldwide, but this contribution remains open to further quantification<sup>[42]</sup>. Screening

programs and early detection of stomach cancer which have been implemented in Japan<sup>[43]</sup> and in Korea<sup>[44]</sup> can partly explain the differences in mortality rates. Also, in Japan advancements were made in surgical treatment of early disease, resulting in a better survival rate compared to other countries<sup>[45]</sup>. But, stomach cancer survival remains unsatisfactory low in most areas of the world<sup>[46,47]</sup>.

High prevalence of *H. pylori* infection is widely recognized as the key contributor to high rates of stomach cancer mortality<sup>[48]</sup>. There is abundant evidence that exposure to other risk factors (tobacco, diet, alcohol use, *etc.*) may have contributed to the apparent international differences in mortality rates of gastric cancer<sup>[49-51]</sup>. Also, disparities in socio-economic status could have an influence on stomach cancer mortality rates, mediated by varying exposures to infection, environmental factors, as well as barriers in accessing medical care<sup>[22,52]</sup>.

### TEMPORAL TRENDS

Declining gastric cancer incidence rates are the dominant epidemiological pattern globally<sup>[1,2]</sup>. Figures 3A and B present information, for males and females, on stomach cancer incidence secular trends for selected populations. In both sexes the underlying pattern was a rapid decline in incidence rates over the whole considered time period, regardless of the background stomach cancer risk. There were two exceptions to this pattern. First exception was seen in the Japanese population (Miyagi prefecture) where, particularly in males, very high rates were observed until the 1990s, and then declined but remained high. Second exception was for the United States population where over the entire time period the rates were constantly very low. The exact reason for the decrease in the incidence of gastric cancer in the last few decades is not completely known, but it most likely includes improvements in diet, food storage and declining prevalence of infection with *H. pylori* due to a general improvement in sanitation and increased use of antibiotics<sup>[53]</sup>. Eradication of *H. pylori* can be achieved with antibiotic therapy; but, the treatment of asymptomatic carriers is not practical because many countries have a very high infection burden (*e.g.*, over 75% of adult persons living in

sub-Saharan Africa have *H. pylori* infection) and reinfection is relatively easy<sup>[54,55]</sup>. Figure 3C and D represent secular trends for stomach cancer mortality, for males and females, in selected countries over the period 1961 to 2016<sup>[2]</sup>. Also, downward trends for stomach cancer mortality rates show a very similar pattern as well as incidence trends. In men, the steep decreasing trends for stomach cancer mortality were observed in all selected countries continuously over the observed period. The two exceptions to mortality pattern were seen. First exception was for Slovakia where, particularly in women, the rates showed a slower downward trend up to the 2000s, with a flattening of the mortality trend from 2010s to onwards. Second exception was for the United States population where mortality rates remained constantly very low over the entire time period. Stomach cancer mortality in both women and men has shown a significant declining trend in most developed countries over the past 50 years<sup>[1,2]</sup>. A similar trend, although starting later, has been seen in some countries in Asia, such as Japan and China<sup>[37,50]</sup>. Factors that led to a decline in mortality involve increased availability of fresh fruits and vegetables, reduced use of salt, reduced incidence of *H. pylori* infection due to improved hygiene and use of antibiotics, and implementation of screening programs<sup>[56,57]</sup>.

By age, incidence and mortality patterns for stomach cancer in women were broadly similar to those in men, regardless of the background stomach cancer risk being high or low (Figure 4). In selected countries, stomach cancer was predominantly a disease of the elderly, and almost 90% of all cases were diagnosed after the age of 55 years (Figure 4A). For both sexes, stomach cancer mortality continuously increases with age, and is two times higher in older than 70 years (Figure 4B).

The favorable trend of stomach cancer incidence in developed countries could largely be attributed to a decrease in *H. pylori* prevalence: this is reflected by the “birth cohort effect” where in some countries (including Korea, Japan, the United States) rates of *H. pylori* have been declining in younger generations<sup>[24,34,38]</sup>. Intestinal adenocarcinoma dominates high-risk areas and is considered responsible for much of the variation in incidence. Recent studies indicate an increase in gastric cancer incidence (cardia and



noncardia stomach cancers combined) in persons under the age of 50 in both low- and high-risk countries (such as the United Kingdom, the United States, Canada, Belarus, Chile)<sup>[58]</sup>. Besides, the increasing prevalence of autoimmune gastritis and dysbiosis of the stomach microbiome could have contributed to the increase of stomach cancer incidence among younger generations<sup>[59]</sup>.

<sup>18</sup> In both men and women, trends in prevalence of cigarette smoking are related to trends in incidence and mortality of stomach cancer with a lag of roughly several decades<sup>[22,60]</sup>. Besides, stomach cancer mortality trends were minimally influenced by changes in the coding of this disease in the second half of the twentieth century<sup>[61]</sup>.

## SURVIVAL

In general, survival for patients with stomach cancer is poor<sup>[5,62]</sup>. Besides, global variation in stomach cancer survival was very wide<sup>[34,47]</sup>. Worldwide, except for Japan and Korea, most areas have an overall 5-year relative survival of stomach cancer of about 20%-30%<sup>[63]</sup>. <sup>4</sup> A 5-year relative stomach cancer survival rate of about 20% is observed in Western developed countries and in developing countries according to an international comparison of data from population based cancer registries<sup>[64-66]</sup>. In contrast to North America and Europe<sup>[66]</sup>, stomach cancer survival is higher in East Asia: *e.g.*, 5-year survival rate is 67% in Korea and 69% in Japan<sup>[67,68]</sup>, followed by Jordan (56%) and Costa Rica (46%) in 2010-2014<sup>[69]</sup>. Also, a notable increase in stomach cancer survival was seen in <sup>4</sup> China (from 30.2% to 35.9%) in recent years<sup>[69]</sup>. These differences are in part explained by the early stomach cancer detection due to the screening programs implemented in East Asia<sup>[63]</sup>. The relatively high overall survival for stomach cancer in Japan is a result of a high proportion of patients being diagnosed in the early stage of the disease: in 1995-2000, 53% of stomach cancers were diagnosed at an early stage in Japan<sup>[70]</sup> in contrast to about 27% in the United States<sup>[71]</sup>. <sup>3</sup> Differences in tumor biology and stomach cancer subtype (with the East/Central Asia and Eastern Europe having a larger proportion of non-cardia stomach cancers than the North America and Western Europe) may also contribute to survival differences<sup>[8,46,72,73]</sup>. Worldwide, the

proportion of cardia stomach cancers, indicating worse prognosis, ranged from 6% in South Korea to 72% in Finland for men, and for women it ranged from 4% in South Korea to 52% in Serbia<sup>[38]</sup>. Similarly, cases with the intestinal subtype had a higher survival rate than patients with diffuse tumors<sup>[70]</sup>. Many factors influence the survival of stomach cancer, including the type of cancer, stage at diagnosis, age, sex, race, overall health, lifestyle<sup>[74-76]</sup>. Generally, due to high aggressiveness and heterogeneity, stomach cancer still remains a severe global health problem<sup>[47,77]</sup>.

## ETIOLOGY AND RISK FACTORS

Stomach cancer is a multifactorial disease. The notable international variation, time trends, and the migratory effect on stomach cancer frequency suggest that environmental and lifestyle factors are very important in the development of this disease.

In 1994, the International Agency for Research on Cancer has classified *H. pylori* infection as a carcinogen in humans due to the evidence which links *H. pylori* infection and risk of gastric cancer<sup>[48]</sup>. *H. pylori* as a carcinogen most likely acts indirectly, by causing gastritis, which is a precursor to stomach atrophy, metaplasia, and dysplasia. While the risk of stomach cancer correlated with the duration of *H. pylori* infection, no association was found for the histologic subtype of stomach cancer (intestinal or diffuse), nor sex. Based on a meta-analysis of cohort studies, the risk of stomach cancer in people with *H. pylori* infection was 2.36<sup>[78]</sup>. Chronic or recurrent *H. pylori* infection is a major cause of stomach cancer; the relative risk is estimated to be 2.7-3.8 for cancer of cardia, and 1.1-11.1 for noncardia stomach cancer<sup>[20]</sup>. *H. pylori* infection attributed to 592000 (63.4%) of all cases of stomach cancer globally<sup>[26]</sup>. *H. pylori* is a stomach colonizing bacterium; how *H. pylori* is transmitted has not been elucidated definitely, but the person-to-person pathway is most likely a contact pathway<sup>[48]</sup>. *H. pylori* infection is acquired in childhood, and population prevalence is associated with socioeconomic status<sup>[79]</sup>. High prevalence of *H. pylori* infection, and little international variation, suggest that other factors are important in the etiology of gastric cancer<sup>[53]</sup>. The main



1 risk factors for noncardia stomach cancer are *H. pylori* infection, tobacco smoking and dietary factors, while gastroesophageal reflux disease, obesity and possibly tobacco smoking play an important role in the development of cardia gastric cancer<sup>[6,9-13,22]</sup>.

17 The main cofactors responsible for the development of stomach cancer are smoking and diet<sup>[30,31,80-82]</sup>. After adjusting for alcohol intake or the presence of chronic *H. pylori* infection in the stomach, an independent association with smoking was confirmed<sup>[83,84]</sup>. 8 Over 45 case-control studies and 27 cohort studies confirmed the association of tobacco with stomach cancer, with the average relative risk being  $RR = 1.5-2.0$ <sup>[85]</sup>. One recent 9 meta-analysis of prospective observational studies suggests that the summary relative risk was higher in men (1.63) than in women (1.30)<sup>[31]</sup>. The risk of stomach cancer increases significantly with cigarette smoking (40 percent for smokers and 82 percent for heavy smokers) and alcohol consumption<sup>[86]</sup>. It is estimated that in developing countries the gastric cancer risk attributable to smoking is 11% in men and 4% in women, while in developed countries the risk is 17% in men and 11% in women<sup>[85,87]</sup>.

While some authors believe that diet has no role in the etiology of gastric cancer, the American Cancer Society states that smoked foods, salted fish and meat, and pickled vegetables represent risk factors for gastric cancer<sup>[88]</sup>. Some bacteria, like *H. pylori*, can convert nitrates and nitrites (commonly found in meat products) to substances which are shown to cause gastric cancer in animals<sup>[48,89]</sup>. It is also known that adherence to the Mediterranean diet is significantly inversely correlated with gastric cancer<sup>[89,90]</sup>.

The correlation between salt intake (high in salt, smoked foods, salted fish and meat) and the stomach cancer risk has been indicated in several epidemiological studies<sup>[91-94]</sup>. Sodium chloride is known to increase gastroduodenal carcinogenesis using N-methyl-N-nitro-N-nitrosoguanidine in a rat experiment<sup>[95]</sup>, as well as in a study in humans<sup>[96]</sup>. The mucin layer which covers and protects the stomach epithelium is damaged by high doses of salt, which also cause high osmotic pressure that further damages the epithelial cells. 8 Prolonged damage to the mucous membrane leads to chronic atrophic gastritis and intestinal metaplasia, which are precursors for stomach cancer.

Higher consumption of fruits and vegetables has been associated with a lower risk of malignant tumors in a number of epidemiological studies (over 200 case-control and cohort studies)<sup>[97,98]</sup>, while results are particularly numerous and consistent for stomach cancer<sup>[99]</sup>. The intake of fresh fruits and vegetables, which contain antioxidant vitamins (e.g. vitamins A and C), reduces gastric cancer risk. In a cohort of 900000 adults (404576 men and 495477 women) who were not diagnosed with malignancy at the time of enrollment, 57145 people died after 16 years, with the highest weight subjects having a higher mortality rate from malignant tumors in general: men had a 52 percent higher rate, and women a 62 percent higher rate, compared to people with normal body weight<sup>[100]</sup>. Higher mortality was found for esophageal, colon, liver, gallbladder, pancreatic and kidney cancer, but also for non-Hodgkin's lymphoma and multiple myeloma.

Patients with gastroesophageal reflux disease (GERD), especially with the long-standing forms, had shown a significantly increased risk for cardia stomach cancer and the majority of studies noted a 2-4-fold increase in risk<sup>[10,101,102]</sup>, although not all<sup>[103]</sup>. One of the explanations for the association between GERD and cardia gastric cancer is that GERD may cause metaplasia with potential progression to adenocarcinoma<sup>[104]</sup>. On the other hand, a lack of association between GERD and noncardia stomach cancer might be explained, at least in part, by the association with atrophic gastritis which might be associated with a decrease in gastric acid secretion and lower risk of GERD<sup>[105]</sup>.

Of the demographic factors, socio-economic status, older age and male gender play an important role<sup>[9-11]</sup>. Socio-economic deprivation, within any population, is consistently linked with increased gastric cancer risk<sup>[106,107]</sup>. The risk of developing stomach cancer increases with age<sup>[7,34]</sup>. Stomach cancer rarely develops before the age of 40, more than 80% of stomach cancers occur between 60 and 80 years of age. Stomach cancer affects men more than women<sup>[1-3]</sup>. The consistency of risk difference by sex has never been adequately explained although possible explanations included differences in environmental exposures and lifestyle factors, as well as the theory regarding the potentially protective role of female sex-specific hormones<sup>[108]</sup>. According to the results

of studies conducted in the United States, stomach cancer occurs more often in African Americans compared to the white population<sup>[7]</sup>. Possible reasons for this increased risk include socioeconomic factors, prevalence of *H. pylori* infection, cigarette smoking, obesity<sup>[7,22,109,110]</sup>.

It is estimated that around 10% of stomach cancer cases aggregate in families, and only 1%-3% are hereditary<sup>[10,13,111]</sup>. A positive family history of stomach cancer in a first-degree relative is a risk factor for stomach cancer, but the magnitude of risk varies with different ethnic groups and geographic regions, ranging from 2 to 10<sup>[112,113]</sup>. Although familial aggregation could be a risk factor because of shared genetic factors, the influence of shared environment cannot be ruled out, *e.g.*, passage of *H. pylori* infection from parents to children, the same dietary factors, *etc.* Although migrant studies indicate a significant reduction in the risk of gastric cancer in Japanese immigrants, results of many studies point out that exposure to environmental factors in childhood is important for determining the gastric cancer risk<sup>[48,53,79]</sup>. Namely, migrant studies show that exposure in childhood is important in stomach cancer etiology: *e.g.* infection with *H. pylori* often occurs before the age of 10, *i.e.* often before the migration, also children born in the immigrant country are likely to acquire the infection from family members who have migrated from their native country<sup>[114]</sup>.

Gastric cancer risk is increased in many genetic disorders, such as hereditary diffuse gastric cancer, Peutz-Jeghers syndrome, Familial adenomatous polyposis<sup>[10,112]</sup>. Persons who have mutations or deletion in genes such as *p53*, *BRCA2*, *MSH2*, *MLH1*, have an increased stomach cancer risk<sup>[10,111]</sup>.

Some studies found a link between stomach cancer and antioxidant use, non-steroidal anti-inflammatory drugs, statins, physical activity, radiation<sup>[6,9-13]</sup>. Some researchers have suggested a correlation between excess or deficit of iodine, goiter, and stomach cancer, as well as a decrease in stomach cancer mortality after performing effective iodine prophylaxis<sup>[115]</sup>.

Other potential risk factors in relation to stomach cancer include poor oral hygiene and tooth loss<sup>[116]</sup>, hookah and opium use<sup>[117]</sup>, Epstein-Barr virus infection<sup>[118]</sup>, and

consumption of pickled vegetables<sup>[119]</sup>, but the results are not convincing, at least as of yet. In addition, in many persons with stomach cancer there is no one specific stomach cancer risk factor.

## **PREVENTION**

During the past century, Western developed countries experienced a major reduction in stomach cancer incidence and mortality, without the introduction of specific primary and secondary prevention measures. Generally, favorable trends in the frequency of stomach cancer are thought to be in important part a consequence of changes such as the reduction in the use of salt and an increase in the consumption of fruit and fresh vegetables due to improvements in food storage (refrigerators, freezers). This phenomenon has been dubbed the “unplanned triumph” of prevention<sup>[9,39]</sup>.

Primary and secondary prevention strategies are the focus of stomach cancer prevention.

Primary prevention measures involve improvements in environment and lifestyle habits such as tobacco control/smoking cessation, reducing salt intake, increasing fruit and vegetable intake, developing other healthy behaviors (such as Mediterranean diet, higher intake of fibers, physical activity), *H. pylori* eradication, other medications (intake of non-steroidal anti-inflammatory drugs, statins), refraining from high alcoholic beverages, sanitation and hygiene improvements. The WHO has set a global goal of reducing the intake of salt to less than 5 g (2000 mg of sodium) per person per day by the year 2025<sup>[120]</sup>. A meta-analysis of randomized trials (all trials were performed in areas with high incidence of stomach cancer, mostly in Asia), in a total of 6695 participants followed from 4 to 10 years has showed that the risk of stomach cancer can be reduced by 35% with the treatment of *H. pylori*<sup>[121]</sup>. In addition to endoscopic and histological surveillance, the American and European guidelines recommend eradication of *H. pylori* in all persons who have atrophy and/or intestinal metaplasia and all persons who are first-degree relatives of stomach cancer patients<sup>[122,123]</sup>. According to the Asian Pacific Gastric Cancer Consensus, population-based screening



and treatment of *H. pylori* infection is recommended in regions which have an annual stomach cancer incidence of more than 20/100000<sup>[124]</sup>. Eradication of *H. pylori* can be achieved with antibiotic therapy; but, the treatment of asymptomatic carriers is not practical because many countries have a very high infection burden (e.g., over 75% of adult persons living in sub-Saharan Africa have *H. pylori* infection) and reinfection is relatively easy<sup>[54]</sup>.

Japan has had a national endoscopic surveillance program since the early 1970s because of the high stomach cancer risk<sup>[125]</sup>. It is recommended that all people older than 40 years undergo screening with a double-contrast barium X-ray radiography and endoscopy every year<sup>[126]</sup>. A study in China demonstrated that a preventive intervention which included eradication of *H. pylori*, nutritional supplements, and screening (with double-contrast radiography and endoscopy) resulted in a 49% reduction in relative risk for overall mortality in a high-risk group of individuals<sup>[127]</sup>.

Upper gastrointestinal endoscopy is the gold standard for stomach cancer diagnosis and due to its high detection rate it is used for stomach cancer screening in high-risk areas (such as Japan, Korea, Venezuela and other areas), but the available evidence shows that endoscopic surveillance of premalignant gastric lesions showed conflicting results<sup>[128]</sup>. Besides, the procedure is expensive, not pleasant for the patient and carries risk of haemorrhage and perforation<sup>[129,130]</sup>.

Stomach cancer screening might be possible via detection of potential markers of gastric atrophy (a stomach cancer precursor lesion)<sup>[125,131,132]</sup>, including serum pepsinogens, serum ghrelin, *H. pylori* serum antibodies, gastrin-17, or antigastric parietal cell antibodies, but the results are not convincing, at least as of yet.

## CONCLUSION

Worldwide, stomach cancer incidence and mortality have declined significantly during the past five decades. However, stomach cancer remains a global health problem as the fifth leading cancer and fourth most common cause of cancer-related deaths in the world. A further illumination of risk factors can help identify various opportunities for

prevention. Primary and secondary prevention strategies with more effectiveness are needed in order to reduce stomach cancer incidence and mortality, particularly in populations with high burden of stomach cancer.



20%

SIMILARITY INDEX

---

PRIMARY SOURCES

---

- |   |  |                |
|---|--|----------------|
| <div style="background-color: red; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">1</div>      | <a href="http://www.wjgnet.com" style="color: red;">www.wjgnet.com</a><br><small>Internet</small>  | 238 words — 4% |
| <div style="background-color: magenta; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">2</div>  | <a href="http://mafiadoc.com" style="color: magenta;">mafiadoc.com</a><br><small>Internet</small>  | 112 words — 2% |
| <div style="background-color: purple; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">3</div>   | <a href="http://link.springer.com" style="color: purple;">link.springer.com</a><br><small>Internet</small>   | 104 words — 2% |
| <div style="background-color: teal; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">4</div>     | <a href="http://www.ncbi.nlm.nih.gov" style="color: teal;">www.ncbi.nlm.nih.gov</a><br><small>Internet</small>   | 88 words — 1%  |
| <div style="background-color: green; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">5</div>    | <a href="#" style="color: green;">Ana Ferro, Bárbara Peleteiro, Matteo Malvezzi, Cristina Bosetti et al. "Worldwide trends in gastric cancer mortality (1980–2011), with predictions to 2015, and incidence by subtype", European Journal of Cancer, 2014</a><br><small>Crossref</small> | 87 words — 1%  |
| <div style="background-color: brown; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">6</div>    | <a href="http://emedicine.medscape.com" style="color: brown;">emedicine.medscape.com</a><br><small>Internet</small>  | 66 words — 1%  |
| <div style="background-color: darkred; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">7</div>  | <a href="#" style="color: darkred;">Nagini, Siddavaram. "Carcinoma of the stomach: A review of epidemiology, pathogenesis, molecular genetics and chemoprevention", World Journal of Gastrointestinal Oncology, 2012.</a><br><small>Crossref</small>                                     | 55 words — 1%  |
| <div style="background-color: darkblue; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">8</div> | <a href="#" style="color: darkblue;">Helicobacter pylori, 2016.</a><br><small>Crossref</small>   |                |

52 words — 1%

---

9 Jacques Ferlay, Hai-Rim Shin, Freddie Bray, David Forman, Colin Mathers, Donald Maxwell Parkin. "Estimates of worldwide burden of cancer in 2008: GLOBOCAN 2008", International Journal of Cancer, 2010  
Crossref 38 words — 1%

---

10 Kai Zhang, Jian Yin, Huiyao Huang, Le Wang, Lanwei Guo, Jufang Shi, Min Dai. "Expenditure and Financial Burden for Stomach Cancer Diagnosis and Treatment in China: A Multicenter Study", Frontiers in Public Health, 2020  
Crossref 36 words — 1%

---

11 Kwong Ming Fock. "Asia-Pacific consensus guidelines on gastric cancer prevention", Journal of Gastroenterology and Hepatology, 3/2008  
Crossref 30 words — < 1%

---

12 Parisa Karimi, Farhad Islami, Sharmila Anandasabapathy, Neal D. Freedman, Farin Kamangar. "Gastric Cancer: Descriptive Epidemiology, Risk Factors, Screening, and Prevention", Cancer Epidemiology Biomarkers & Prevention, 2014  
Crossref 30 words — < 1%

---

13 *Helicobacter pylori*, 2000.  
Crossref 27 words — < 1%

---

14 [ijmbs.org](http://ijmbs.org)  
Internet 26 words — < 1%

---

15 D COGGON, E D ACHESON. "THE GEOGRAPHY OF CANCER OF THE STOMACH", British Medical Bulletin, 1984  
Crossref 24 words — < 1%

- 
- 16 [www.spandidos-publications.com](http://www.spandidos-publications.com) 23 words — < 1%  
Internet
- 
- 17 Crew, Katherine D, and Alfred I Neugut. 22 words — < 1%  
"Epidemiology of gastric cancer", *World Journal of Gastroenterology*, 2006.  
Crossref
- 
- 18 Elham Goodarzi, Ali Hassanpour Dehkordi, Reza Beiranvand, Hasan Naemi, Zaher Khazaei. 21 words — < 1%  
"Epidemiology of the Incidence and Mortality of Pancreas Cancer and its Relationship with the Human Development Index (HDI) in the World: An Ecological Study in 2018", *Current Pharmaceutical Design*, 2020  
Crossref
- 
- 19 O. Andrade. "Chemoprevention of Precancerous Gastric Lesions With Antioxidant Vitamin Supplementation: A Randomized Trial in a High-Risk Population", *JNCI Journal of the National Cancer Institute*, 01/17/2007 21 words — < 1%  
Crossref
- 
- 20 [www.cancer.gov](http://www.cancer.gov) 19 words — < 1%  
Internet
- 
- 21 Masoud Amiri. "The decline in stomach cancer mortality: exploration of future trends in seven European countries", *European Journal of Epidemiology*, 11/18/2010 13 words — < 1%  
Crossref
- 
- 22 [journals.physiology.org](http://journals.physiology.org) 13 words — < 1%  
Internet
- 
- 23 Haejin In, Marisa Langdon-Embry, Lauren Gordon, Clyde B. Schechter et al. "Can a gastric 10 words — < 1%

cancer risk survey identify high-risk patients for endoscopic screening? A pilot study", Journal of Surgical Research, 2018

Crossref

24 Chun-Dong Zhang, Hiroharu Yamashita, Yasuyuki Seto. "Gastric cancer surgery: historical background and perspective in Western countries versus Japan", Annals of Translational Medicine, 2019

9 words — < 1%

Crossref

25 E Fernandez, J.R González, J.M Borràs, V Moreno, V Sánchez, M Peris. "Recent decline in cancer mortality in Catalonia (Spain). A joinpoint regression analysis", European Journal of Cancer, 2001

9 words — < 1%

Crossref

26 Linghui Zhou, Qin Yu, Guoqing Wei, Linqin Wang, Yue Huang, Kejia Hu, Yongxian Hu, He Huang. "Global, Regional, and National Burden of Multiple Myeloma from 1990 to 2019: Estimates from the 2019 Global Burden of Disease study", Research Square, 2020

9 words — < 1%

Crossref Posted Content

27 Patrick Andersen, Anja Mizdrak, Nick Wilson, Anna Davies, Laxman Bablani, Tony Blakely. "Disaggregating proportional multistate lifetables by population heterogeneity to estimate intervention impacts on inequalities", Cold Spring Harbor Laboratory, 2021

9 words — < 1%

Crossref Posted Content

28 docksci.com

Internet

9 words — < 1%

29 de Moura Gallo, C.V.. "TP53 mutations as biomarkers for cancer epidemiology in Latin America: Current knowledge and perspectives", Mutation Research-Reviews in Mutation Research, 200505

8 words — < 1%

Crossref

31 Elham Kharazmi, Masoud Babaei, Mahdi Fallah,  
Tianhui Chen, Kristina Sundquist, Kari Hemminki.

7 words — < 1%

"Importance of tumor location and histology in familial risk of  
upper gastrointestinal cancers: a nationwide cohort study",  
Clinical Epidemiology, 2018

Crossref

32 L.-N. Mu, W. Cao, Z.-F. Zhang, S.-Z. Yu et al.

7 words — < 1%

"Polymorphisms of 5,10-methylenetetrahydrofolate  
reductase (MTHFR), fruit and vegetable intake, and the risk of  
stomach cancer", Biomarkers, 2008

Crossref

33 Melissa A. Jim, Paulo S. Pinheiro, Helena Carreira,  
David K. Espey, Charles L. Wiggins, Hannah K.

7 words — < 1%

Weir. "Stomach cancer survival in the United States by race and  
stage (2001-2009): Findings from the CONCORD-2 study",  
Cancer, 2017

Crossref

EXCLUDE QUOTES OFF

EXCLUDE MATCHES OFF

EXCLUDE BIBLIOGRAPHY OFF