

77462\_Auto\_Edited.docx

**Name of Journal:** *World Journal of Gastroenterology*

**Manuscript NO:** 77462

**Manuscript Type:** LETTER TO THE EDITOR

**Comments on** <sup>3</sup> "Development and validation of a prediction model for moderately severe and severe acute pancreatitis in pregnancy"

Comments on a prediction model

Qun-Ying Yang, Jian-Wen Hu

## Abstract

We reviewed a study addressing the development and validation of a prediction model for moderately severe and severe acute pancreatitis in pregnancy. We identified some statistical deficiencies in this article. In addition, we believe that the role of cholesterol as a predictor should be described in more detail.

**Key Words:** Acute pancreatitis; Prediction model; Statistical analyses; Cholesterol

Yang QY, Hu JW. Comments on "Development and validation of a prediction model for moderately severe and severe acute pancreatitis in pregnancy". *World J Gastroenterol* 2022; In press

**Core Tip:** This is a comment on a study involving the development and validation of a prediction model for moderately severe and severe acute pancreatitis in pregnancy. We believe that the role of cholesterol as a predictor should be more clearly described.

## <sup>2</sup> TO THE EDITOR

We were pleased to read the high-level article published by Yang <sup>1</sup>. In this article, the authors developed and validated a nomogram with good accordance for the prediction of moderately severe and severe acute pancreatitis in pregnancy (MSIP). The authors reported a nomogram that incorporated numerous blood indices for albumin, lactate dehydrogenase, triglyceride, and cholesterol levels, thus facilitating the early individualized prediction of the severity of acute pancreatitis in pregnancy (APIP). This study is of great significance for the clinical management of APIP. However, in our opinion, this article has some problems that need to be discussed further.

First, we found that some data that were not suitable for this article. In the result sections, we noticed that 134 patients were classified as having mild acute pancreatitis in pregnancy (MAIP) and 56 patients were classified as having MSIP. However, Figure 1 (<https://www.wjgnet.com/1007-9327/full/v28/i15/WJG-28-1588->

g001.htm) and Table 1 (<https://www.wjgnet.com/1007-9327/full/v28/i15/1588-T1.htm>) showed that the numbers of patients with MAIP and MSIP were 136 and 54, respectively. This inconsistency should be addressed.

In addition, there are problems associated with the statistical analyses in that the methods used for statistical analysis should be described in more detail. In Table 1 (<https://www.wjgnet.com/1007-9327/full/v28/i15/1588-T1.htm>) and Table 2 (<https://www.wjgnet.com/1007-9327/full/v28/i15/1588-T2.htm>) of the results section, the authors should provide more accurate statistical values, including Student's t-values or c2 values, instead of just providing P values. Most of the variables mentioned by the authors in Table 1 (<https://www.wjgnet.com/1007-9327/full/v28/i15/1588-T1.htm>) and Table 2 (<https://www.wjgnet.com/1007-9327/full/v28/i15/1588-T2.htm>), such as cholesterol and platelets, are not labeled with units of measurement. Moreover, P values were not listed in the statistical results for "trimester of pregnancy on admission".

In the discussion sections, the authors mentioned that hypercholesterolemia is a known risk factor for cardiovascular diseases. In fact, high-density lipoprotein (HDL) and non-HDL cholesterol have opposite associations with cardiovascular diseases<sup>[2]</sup>, plasma HDL-cholesterol concentrations correlate negatively with the risk of cardiovascular diseases<sup>[3]</sup>. The authors mentioned that cholesterol is a predictive factor of MSIP, the cholesterol levels of patients with MAIP and MSIP were  $7.34 \pm 5.63$  and  $12.80 \pm 6.64$  respectively in Table 1 (<https://www.wjgnet.com/1007-9327/full/v28/i15/1588-T1.htm>). The authors mentioned some previous studies in their discussion section. We took a close look at these studies and found that only one recent study<sup>[4]</sup> showed that low levels of total cholesterol (TC) and high TC within 24h of admission were independently associated with an increased risk of SAP. Other studies<sup>[5-7]</sup> have suggested that serum levels of HDL-cholesterol are inversely correlated with disease severity in patients with predicted severe acute pancreatitis. However, this study showed that cholesterol is a predictive factor for MSIP but not HDL. This is a confusing

statement. We believe that the present study is inconsistent with previous studies and that the role of cholesterol as a predictor should be more clearly described.

21%

SIMILARITY INDEX

PRIMARY SOURCES

- 1 Mitsushige Sugimoto, Hideo Yasuda, Akira Andoh. " Nutrition status and infection in patients receiving hemodialysis ", World Journal of Gastroenterology, 2018 36 words — 5%

Crossref
- 2 Wei Wang, Chen-Chen Pan, Wen-Ying Zhao, Jin-Yu Sheng, Qi-Qi Wu, Si-Si Chen. "Comments on "Effect of type 2 diabetes mellitus in the prognosis of acute-on-chronic liver failure patients in China"", World Journal of Gastroenterology, 2022 32 words — 5%

Crossref
- 3 [www.wjgnet.com](http://www.wjgnet.com) 28 words — 4%

Internet
- 4 [www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) 22 words — 3%

Internet
- 5 Wei Wang, Zhi-Peng Li, Juan Zhang, Zi-Jun Lin, Mei-Lin Ma, Si-Si Chen. " Comment "Asymptomatic small intestinal ulcerative lesions: Obesity and are likely to be risk factors" ", World Journal of Gastroenterology, 2022 14 words — 2%

Crossref
- 6 [jyx.jyu.fi](http://jyx.jyu.fi) 12 words — 2%

Internet

---

EXCLUDE QUOTES            ON

EXCLUDE BIBLIOGRAPHY   ON

EXCLUDE SOURCES           < 12 WORDS

EXCLUDE MATCHES           < 12 WORDS