

83714\_Auto\_Edited.docx

***Basic Study*****Dihydroergotamine ameliorates liver fibrosis by targeting transforming growth factor  $\beta$  type II receptor**

Dihydroergotamine ameliorates liver fibrosis

Ke-Xin Zheng, Shou-Li Yuan, Meng Dong, Han-Lin Zhang, Xiao-Xiao Jiang, Chun-Long Yan, Rong-Cai Ye, Hui-Qiao Zhou, Li Chen, Rui Jiang, Zi-Yu Cheng, Zhi Zhang, Qi Wang, Wan-Zhu Jin, Wen Xie

**Abstract****BACKGROUND**

The transforming growth factor  $\beta$  (TGF $\beta$ ) signaling pathway plays a crucial role in the development of liver fibrosis by activating TGF $\beta$  type II receptor (TGF $\beta$ R2), followed by the recruitment of TGF $\beta$ R1 finally triggering downstream signaling pathway.

**AIM**

To find drugs targeting TGF $\beta$ R2 that inhibit TGF $\beta$ R1/TGF $\beta$ R2 complex formation, theoretically inhibit TGF $\beta$  signaling pathway, and thereby ameliorate liver fibrosis.

**METHODS**

Food and Drug Administration (FDA)-approved drugs were screened for binding affinity with TGF $\beta$ R2 by virtual molecular docking. We identified 6 candidates and further explored their potential by Cell Counting Kit-8 (CCK-8) cell cytotoxic experiment to validate toxicity and titrated the best cellular working concentrations.

Next, we further demonstrated the detailed molecular working mechanisms using mutagenesis analysis. Finally, we used a mouse model to investigate its potential anti-liver fibrosis effect.

## RESULTS

We identified 6 drug candidates. Among these 6 drugs, dihydroergotamine (DHE) shows great ability in reducing fibrotic gene expressions such as collagen, p-SMAD3, and  $\alpha$ -SMA in TGF $\beta$  induced cellular model of liver fibrosis in LX-2 cells. Furthermore, we demonstrated that DHE binds to TGF $\beta$ R2. Moreover, mutation of Leu27, Phe30, Thr51, Ser52, Ile53, and Glu55 of TGF $\beta$ R2 disrupted the binding of TGF $\beta$ R2 with DHE. In addition, DHE significantly improved liver fibrosis, as evidenced by Masson's trichrome staining of liver sections. This is further supported by the width and the velocity of the portal vein, and serum markers of liver function. In line with those observations, DHE also decreased macrophage infiltration and extracellular matrix deposition in the liver.

## CONCLUSION

DHE alleviates liver fibrosis by binding to TGF $\beta$ R2 thereby suppressing TGF $\beta$  signaling pathway. We show here that as far as drug repurposing, DHE has great potential to treat liver fibrosis.

**Key Words:** Liver fibrosis; transforming growth factor  $\beta$  (TGF $\beta$ ) signaling pathway; TGF $\beta$  type II receptor (TGF $\beta$ R2); Virtual screening; Drug-repurposing; Dihydroergotamine (DHE)

Zheng KX, Yuan SL, Dong M, Zhang HL, Jiang XX, Yan CL, Ye RC, Zhou HQ, Chen L, Jiang R, Cheng ZY, Zhang Z, Wang Q, Jin WZ, Xie W. Dihydroergotamine ameliorates liver fibrosis by targeting transforming growth factor  $\beta$  type II receptor. *World J Gastroenterol* 2023; In press

**Core Tip:** An effective and safe drug for treating liver fibrosis is urgently needed in current clinical practice. Here, we investigated and discovered that dihydroergotamine (DHE) could alleviate liver fibrosis by specific binding of transforming growth factor  $\beta$  type II receptor (TGF $\beta$ R2) to disrupt the binding of TGF $\beta$ R2 with TGF $\beta$ 1, and ultimately suppressing its downstream TGF $\beta$  signaling pathway. DHE may be an effective anti-liver fibrosis drug, which could be employed in liver cirrhotic patients.

## **INTRODUCTION**

Liver fibrosis is the consequence of various chronic pathogenic factors<sup>[1]</sup>, it is a dynamic process that is characterized by an excessive accumulation of extracellular matrix<sup>[2]</sup>. Early liver fibrosis can be reversed to a normal architecture by removal of underlying causes<sup>[2]</sup>, but liver fibrosis could further develop into cirrhosis without effective treatment<sup>[3]</sup>. Liver cirrhosis can be complicated by variceal bleeding, hepatic encephalopathy, ascites, bacterial peritonitis, and hepatocellular carcinoma, which has high mortality<sup>[4]</sup>. Liver cirrhosis can regress to early stage of disease, but it cannot be reversed to a normal liver<sup>[5]</sup>. Therefore, it is very important to control the disease progression in the early reversible stage of liver fibrosis.

Etiological treatment of liver fibrosis is most important and effective, such as antivirals, quitting alcohol consumption, and weight loss<sup>[1]</sup>. However, effects of etiological treatment are limited and insufficient, and difficult to prevent the development of liver fibrosis into cirrhosis. At present, liver transplantation is a radical cure for cirrhosis but is associated with a high cost, organ shortages, and the risk of immune rejection<sup>[6]</sup>. In addition, almost all current clinical trials targeting fibrosis are focused on non-alcoholic steatohepatitis, specifically focusing on hepatic stellate cell (HSC) activation and / or fibrogenesis<sup>[7]</sup>. However, there are still no approved antifibrotic therapies for liver fibrosis<sup>[7]</sup>. Therefore, it is urgent to develop new effective drugs.

Developing new drugs is a difficult, high-cost, and extremely low success rate procedure. A good strategy to address this problem is to investigate new indications of

old drugs, a process called “drug repurposing”<sup>[8]</sup>. Scientists have repurposed many old drugs such as propranolol<sup>[9]</sup>, cimetidine<sup>[10]</sup>, sildenafil<sup>[11]</sup>, and thalidomide<sup>[12]</sup>. Thus, drug repurposing is an attractive approach and has been widely employed. Drugs that have been approved by Food and Drug Administration (FDA) have passed preliminary clinical trials and are considered extremely safe. Therefore, FDA-approved drugs may be good candidates for developing new indications. Molecular docking, a fast, efficient, and widely used technique in drug repurposing, is a computational strategy to predict binding sites between ligands and targets based on their structures<sup>[8, 13]</sup>.

The activation of HSCs is considered the central effector of liver fibrosis<sup>[14]</sup>. There are many associated signaling molecules, including transforming growth factor  $\beta$  (TGF $\beta$ ), platelet-derived growth factor, and connective tissue growth factor<sup>[15]</sup>. The TGF $\beta$  signaling pathway plays a crucial role in the development of liver fibrosis<sup>[16]</sup>. A review paper illustrated that TGF $\beta$ 1 activates TGF $\beta$  type II receptor (TGF $\beta$ R2), followed by the recruitment of TGF $\beta$ R1. Afterward, TGF $\beta$ R2 phosphorylates TGF $\beta$ R1 thereby triggering down-stream signaling pathway to regulate the expression of collagens and extracellular matrix (ECM)<sup>[17]</sup>. Therefore, TGF $\beta$ R2 is considered an important target for developing drugs against liver fibrosis. Consistently, our group and others demonstrated that both inhibiting the expression of TGF $\beta$ R2 and exogenous extracellular domain of TGF $\beta$ R2 supplement effectively alleviated liver fibrosis<sup>[18, 19]</sup>.

In the current study, FDA-approved drugs were screened for binding affinity with the TGF $\beta$ R2 by virtual molecular docking. We used cellular and mouse models to investigate its potential anti-liver fibrosis effect. In addition, by using mutagenesis analysis we further demonstrated detailed molecular working mechanism.

## **MATERIALS AND METHODS**

Molecular docking of FDA-approved drugs and TGF $\beta$ R2  
The structures of FDA-approved drugs were downloaded from the ZINC database (<https://zinc12.docking.org/>). Then, the files of each small molecular structures were generated using Open Babel GUI (3.1.1). The TGF $\beta$ R2-TGF $\beta$ 1 complex (PDB: 3KFD) was

downloaded from the Protein Data Bank (<https://www.rcsb.org/>). The TGF $\beta$ R2 structure was derived from the TGF $\beta$ R2-TGF $\beta$ 1 complex using AutoDock Vina software (<https://vina.scripps.edu/>). AutoDock Vina software was employed to screen for the lowest energy complex among complexes of the extracellular TGF $\beta$ R2 domain and the FDA-approved drugs.

Cell culture and viability experiments

The human HSC line LX-2 was purchased from the BeNa Culture Collection (Beijing, China) and cultured in Dulbecco's modified Eagle medium (DMEM, Gibco, USA) with 10% fetal bovine serum (FBS, Gibco, USA) at 37°C in a 5% CO<sub>2</sub> atmosphere. LX-2 was activated by human recombinant protein TGF $\beta$ 1 (5 ng/mL) for 24 h, followed by drug treatment for 24 h. The expression of fibrosis-related genes was analyzed. Cell Counting Kit-8 (CCK-8, Beyotime Biotechnology, Shanghai, China) test was used to test the cytotoxicity of a series of working concentrations of the candidate drugs on LX-2 cell. After 24 h of treatment, the CCK-8 solution was added to each well for an additional 1 h of incubation (37°C in a 5% CO<sub>2</sub> atmosphere). A microplate analyzer was used to measure the optical density of each well at 450 nm. Cell vitality was expressed as a percentage of optical density between the treatment wells and the negative control cells.

Determination of K<sub>d</sub> value between TGF $\beta$ R2 and dihydroergotamine

The K<sub>d</sub> value of the small molecules and the TGF $\beta$ R2 protein was measured using a microscale thermophoresis (MST)-Nanotemper instrument (Nanotemper, Germany). Micro thermophoresis is the directional movement of particles in the micro temperature gradient. The affinity is determined by measuring the change of micro thermophoresis caused by the change of hydration layer (usually caused by the change of biomolecular structure / conformation). First, 100  $\mu$ L each of 10  $\mu$ M TGF $\beta$ R2 and different concentrations of the small molecules (diluted from 400  $\mu$ M stock) were prepared. Second, TGF $\beta$ R2 was labeled with a fluorescent dye and mixed with the small molecules. Third, fluorescence was measured to assess the binding behavior of the small molecule-TGF $\beta$ R2 complex.

## Molecular dynamics simulation

Molecular dynamics simulations were performed using Gromacs 2020.1, in which a charm<sup>36-mar2019</sup> force field was chosen. The TGFβR2 or TGFβR2-small molecule complex was solved with TIP3P water and immersed in a dodecahedron box extending to at least 1 nm of the solvent on all sides. The system was neutralized with Na<sup>+</sup> and Cl<sup>-</sup> by adding 0.15 M NaCl. It was energy-minimized using the steepest descent algorithm for 5,000 steps and creating a maximum force of <1,000 kJ/mol/nm. After energy minimization, the system was equilibrated with a constrained “number of particles, volume, and temperature” (NVT) and “number of particles, pressure, and temperature” (NPT) running for 100 ps. Through NVT and NPT equilibration, the system was well-equilibrated at 300 K and 1 bar. Finally, MD simulations of the TGFβR2 or complex were carried out for 200 ns; trajectories were saved every 10 ps for analysis. The Verlet cut-off scheme and a Leap-frog integrator with a step size of 2 fs were applied. The modified Berendsen thermostat was used for temperature coupling, the Parrinello-Rahman barostat was used for pressure coupling, and the Particle Mesh Ewald method was used to determine long-range electrostatic interactions. The root-mean-square displacement (RMSD) and root-mean-square fluctuation (RMSF) of TGFβR2 and the complex were calculated using GROMACS 2020.1. The content of the secondary structure was calculated using DSSP software<sup>[20, 21]</sup>. The last MD simulation frame of the complex was extracted using GROMACS 2020.1. LigPlot+ software (<https://www.ebi.ac.uk/thornton-srv/software/LigPlus/>) was used to analyze the detailed interactions between the extracellular TGFβR2 domain and small molecule<sup>[22]</sup>. PyMOL (<http://www.pymol.org/>) was used to prepare the structural images.

Production and purification of site-directed mutagenesis of TGFβR2

The binding sites of TGFβR2 and small molecule were established using the molecular docking results. The amino acids at the binding sites were mutated to alanine whose nucleotide sequence was GCG. The sequences of mutated extracellular TGFβR2 and a 6X His-tag were inserted into a pMAL-c5x plasmid that was synthesized by GenScript



Biotech

Corporation.

The mutated plasmid was transformed into BL21(DE3) bacteria and induced with 0.8 mmol/L/L IPTG for 16 h at 28°C. Then, bacteria were collected and resuspended in PBS. The mutated TGFβR2 protein was purified by HisTrap FF affinity chromatography and eluted with PBS containing gradient concentrations of imidazole. Finally, the eluted proteins were concentrated and the imidazole was removed by dialysis.

CCl<sub>4</sub>

injury

mouse

model

Six-week-old male C57BL/6N mice were purchased from Vital River Laboratory Animal Technology Co. Ltd. to induce a liver fibrosis model. The animal protocol was designed to minimize pain or discomfort to the animals. The animals were acclimatized to laboratory conditions (23 °C, 12 h/12 h light/dark, 50% humidity, ad libitum access to food and water) for 2 wk prior to experimentation. After being transferred to our institute, the animals were randomly and evenly divided into four groups (eight mice per group) according to their body weight: corn oil, carbon tetrachloride (CCl<sub>4</sub>), low-concentration treatment, and high-concentration treatment groups. The same volume of CCl<sub>4</sub> (0.5 μL/g of body weight, Sigma-Aldrich, St. Louis, MO, USA) and corn oil were intraperitoneally injected three times a week to induce a liver fibrosis model and in control animals for four weeks, respectively. Intragastric gavage administration was carried out with conscious animals, using straight gavage needles appropriate for the animal size. Then, the small molecule aqueous solution and water was administered orally once a day as treatment and control for eight weeks, respectively. All animals were euthanized for tissue collection. All animal studies were performed following the National Institutes of Health's Guide for the Care and Use of Laboratory Animals and conducted with the approval of the Institutional Animal Care and Use Committee of the Institute of Zoology, Chinese Academy of Sciences.

B-mode

ultrasonography

Small-animal B-ultrasound was used to inspect the portal veins. A high-resolution ultrasound imaging system (Vevo LAZR, VisualSonics, Canada) was used to measure the width and velocity of the portal vein. Mice were fasted for 12 h and shaved before



the ultrasonic examination. Then, the mice were fixed on the platform and examined after being anesthetized with Avertin. After coating the mouse's abdomen with the coupling gel, the probe was used to inspect the mouse's portal vein.

#### Masson's trichrome staining

Mouse liver tissues were fixed in 4% paraformaldehyde, embedded in paraffin, and sectioned at 5- $\mu$ m thickness. Then, the liver paraffin-embedded tissue sections were stained with a Masson's trichrome kit (G1281, Solarbio, Beijing, China) and observed under a 10x objective lens. Masson's trichrome staining is used to distinguish collagen fibrosis from muscle fibers. Muscle fibers were stained red and collagen fibrosis was stained green or blue. The collagen area was quantified using ImageJ 1.52a software. And one liver section was taken from each mouse for analysis.

#### Western blot analysis

Total proteins were extracted from the LX-2 cells or liver tissues using radioimmunoprecipitation assay lysis buffer (50 mmol/L Tris-HCl, pH 7.4; 1% NP-40; 0.25% sodium deoxycholate; 150 mmol/L NaCl; and 1 mmol/L EDTA) containing a protease and phosphatase inhibitor mixture (Roche Diagnostics). The proteins were separated on 10% SDS polyacrylamide gels and transferred onto polyvinylidene fluoride membranes (Millipore). After blocking with 5% skim milk in Tris-buffered saline-Tween 20 (0.02 M Tris base, 0.1% Tween 20, 0.14 M NaCl, pH 7.4) for 1 h, the membranes were incubated with primary antibodies overnight at 4°C. The membranes were then incubated with HRP-conjugated secondary antibodies for 1 h at room temperature. Table S1 Lists the antibodies.

#### Real-time quantitative PCR (RT-PCR) analysis

Total RNA was extracted from the LX-2 cells or liver tissues using TRIzol™ Reagent (Thermo Fisher Scientific, USA) according to the manufacturer's instructions. Complementary DNA (cDNA) was obtained from the reverse-transcribed RNA using a high-capacity cDNA reverse-transcription kit (Promega, USA). The relative expression of genes was analyzed by RT-PCR (Light Cycler 480, Roche, Sweden) with SYBR Green Master Mix (Promega, USA). Table S2 Lists the primer sequences.

Isolation of liver mononuclear cells (MNCs) and flow cytometry analysis

Liver mononuclear cells (MNCs), including Kupffer cells, were isolated from mouse liver tissues. Liver samples were collected from mice under deep anesthesia. The liver samples were cut into pieces, transferred into 5 mL of enzyme mix (RPMI 1640 containing 1 mg/mL collagen IV [Sigma] and 20 U/mL DNase I [Roche, Burgess Hill, UK]), and digested in a water bath at 37°C for 30 min. RPMI 1640 containing 10% FBS was added to arrest the digestion. The digested mixture was filtered through a 70  $\mu$ m strainer, centrifuged at 4°C for 10 min at 1,000  $\times$  g, and the pellet was washed in PBS twice. Liver immune cells were subsequently isolated using a 33% Percoll cell separation solution after centrifuging for 25 min at 2,200  $\times$  g at room temperature. The red blood cells were removed using a red blood cell lysis buffer (YESEN, Shanghai, China). Finally, liver immune cells were counted and stained with anti-F4/80 antibody eFluor® 450 (eBioscience, San Diego, CA), brilliant violet 510™-conjugated anti-CD45 antibody (Biolegend, San Diego, CA), and percp-cyanine5.5-conjugated anti-CD11b antibody (eBioscience, San Diego, CA), which were then filtered into flow tubes through a 0.45  $\mu$ m strainer. The result of flow cytometry was analyzed using a BD Fortessa instrument (BD, New York, USA).

Plasma biochemical markers

Mouse blood samples were collected from the tail vein, collected into EDTA-containing tubes, and gently shaken. Plasma was collected after centrifuging at 3,000  $\times$  g for 15 min at 4°C. Plasma alanine aminotransferase (ALT) and aspartate aminotransferase (AST) levels were measured using a biochemistry analyzer (Cobas c 501, Roche, Sweden).

Statistical analyses

Data are expressed as mean  $\pm$  standard error of the mean (SEM). The statistical significance between groups was analyzed using one-way ANOVA test. Statistical significance was set at  $p < 0.05$  and  $p < 0.05$ ,  $p < 0.01$ ,  $p < 0.001$ , and  $p < 0.0001$  were denoted as \*, \*\*, \*\*\*, and \*\*\*\*, respectively. GraphPad Prism software was used to perform all statistical analyses.

## **RESULTS**

The docking analysis of FDA-approved small molecule drugs and TGF $\beta$ R2 AutoDock Vina software was employed to dock drugs with the extracellular domain of TGF $\beta$ R2 and output the complex structures. The complex and the binding sites of TGF $\beta$ R2 and TGF $\beta$ 1 were shown (Figure 1A), and the structure of TGF $\beta$ R2 was split from the complex (Figure 1B). The binding affinity of each FDA-approved drug was evaluated (Figure 1C). Among those, Darifenacin, cyproheptadine, lifitegrast, difenoxin, phenytoin, dihydroergotamine (DHE), naldemedine, and irinotecan showed higher scores out of 1615 drugs (Figure 1C). Darifenacin is a competitive muscarinic M receptor antagonist used to treat urinary frequency, urgency, and incontinence caused by bladder hyperstimulation<sup>[21]</sup>. Cyproheptadine is an antihistamine used to treat allergies<sup>[22]</sup>. Lifitegrast is a small molecule integrin inhibitor primarily used to treat symptoms and signs of dry eye<sup>[23]</sup>. Difenoxin is a human metabolite of diphenoxylate, which is a derivative of pethidine and can be used to treat functional diarrhea and chronic enteritis<sup>[24]</sup>. Phenytoin is an effective voltage-gated Na<sup>+</sup> channel blocker used to treat epilepsy, neuralgia, and arrhythmia<sup>[25]</sup>. DHE is an adrenergic receptor antagonist used to treat severe orthostatic hypotension, migraine, and headache<sup>[26]</sup>, which can bind with various receptors. <sup>2</sup> DHE is also an agonist at 5-HT1B, 5-HT1D, and 5-HT1F receptors, but it also binds to 5-HT1A and 5-HT2A receptors. Naldemedine is an opioid receptor antagonist used to treat non-cancerous pain and opioid-induced constipation<sup>[27]</sup>. Irinotecan is a semi-synthesis of water-soluble camptothecin derivatives used to treat advanced colorectal cancer and postoperative adjuvant chemotherapy<sup>[28]</sup>. (Table 1). All the above drugs were purchased from Selleck Chemicals, except for difenoxin and naldemedine which are banned from purchase as they are under management control. Therefore, we performed the following experiments using the rest six small molecule drugs.

Cytotoxicity of candidate drugs

To investigate the cytotoxicity of these drugs, LX-2 cells were treated with 20  $\mu$ M of each candidate for 24 h. The results demonstrated that irinotecan and cyproheptadine

were cytotoxic to LX-2 cells at this concentration (Figure 1D). Therefore, irinotecan and cyproheptadine were regarded as cytotoxic drugs and excluded from subsequent experiments. Next, to identify the best range of concentrations that would not influence the viability of these cells, LX-2 cells were treated with different concentrations of the remaining drugs for 24 h. The results demonstrated that neither lifitegrast nor phenytoin was cytotoxic to LX-2 cells when concentrations were up to 100  $\mu$ M and 150  $\mu$ M, respectively (Figure 1E and F). Darifenacin and DHE were cytotoxic to LX-2 cells when concentrations were beyond 50  $\mu$ M and 20  $\mu$ M, respectively (Figure 1G and H). Therefore, to avoid interference on fibrosis-related gene expression due to the cytotoxicity of candidate drugs, lifitegrast and phenytoin concentrations below 100  $\mu$ M and darifenacin and DHE concentrations below 20  $\mu$ M were used to treat LX-2 cells.

Anti-fibrotic effect of candidate drugs in cellular model. Increased expression of collagen and  $\alpha$ -smooth muscle actin ( $\alpha$ -SMA) are principal markers of HSC activation<sup>[29]</sup>. LX-2 cells were treated with different concentrations of drugs for 24 h after TGF $\beta$ 1 stimulation. The results demonstrated that lifitegrast, phenytoin and darifenacin did not decrease the protein levels of collagen III and  $\alpha$ -SMA after TGF $\beta$ 1 stimulation (Figure 2A and B). Moreover, darifenacin did not decrease the protein levels of collagen III, p-SMAD3, and  $\alpha$ -SMA as much as DHE (Figure 2C and D). Consistently, DHE significantly decreased the mRNA expression of collagen I  $\alpha$ 1 (COL1A1), collagen I  $\alpha$ 2 (COL1A2), collagen III  $\alpha$ 1 (COL3A1), and  $\alpha$ -SMA compared with darifenacin (Figure 2E and F). Taken together, DHE (PubChem CID: 10531) was the most effective small molecule drug that suppressed the TGF $\beta$ 1 induced LX-2 activation.

The binding affinity of DHE to TGF $\beta$ 2. We further verified the molecular docking result by MST-Nanotemper. The results demonstrated that the fluorescence intensity of TGF $\beta$ 2 changed gradually in proportion to the DHE concentration (Figure 3A). The affinity is determined by measuring the change of micro thermophoresis caused by the change of hydration

layer, and results showed the binding affinity of DHE to TGFβR2 with a K<sub>d</sub> value of 17.64 μM.

Molecular dynamics simulation of DHE and TGFβR2 To identify the specific binding mode of DHE and TGFβR2, we used the complex structure obtained from AutoDock vina to perform the molecular dynamics simulation with GROMACS 2020.1 software for 200 ns. The results demonstrated that the RMSD of the backbone atoms of TGFβR2 and TGFβR2–DHE in the simulation system reached equilibrium after 100 ns (Figure 3B). The RMSFs of the TGFβR2 skeleton carbon atoms in the two simulated systems were almost identical (Figure 3C). The secondary structural elements of TGFβR2 in the complex simulation were only slightly changed (Figure 3D and Table 2). The coil and 3-helix components of TGFβR2 decreased in the complex simulation system (Table 2). Finally, we extracted the simulated complex structure and analyzed the amino acid sites of TGFβR2 bound to DHE by Ligplot+ software (Figure 3E and F). In conclusion, our molecular dynamics simulations showed that the binding of DHE to TGFβR2 has little effect on the structure of TGFβR2. We also obtained the binding sites of TGFβR2 to DHE by stimulations.

Leu27, Phe30, Thr51, Ser52, Ile53, and Glu55 of TGFβR2 bind with DHE Molecule docking demonstrated that Leu27, Phe30, Thr51, Ser52, Ile53, and Glu55 of TGFβR2 were binding sites for DHE (Figure 3F). To verify these binding sites, we mutated the above-mentioned amino acids to alanine. The DNA sequences of mutated extracellular TGFβR2 were synthesized (Table 3) and the binding affinity was measured. The results demonstrated that the mutated TGFβR2 no longer bound to DHE (Figure 3G). Thus, the above-mentioned binding sites of TGFβR2 predicted by molecule docking were the exact binding sites of TGFβR2 and DHE, and this further verified the binding of TGFβR2 and DHE. Therefore, we inferred that the binding of TGFβ1 and TGFβR2 was blocked by the binding of DHE and TGFβR2, further preventing the TGFβ signaling cascade.

DHE alleviated liver fibrosis in mouse To explore whether DHE could alleviate fibrosis in vivo, we used two different

concentrations of DHE to treat a CCl<sub>4</sub>-induced mouse fibrosis model. Six-week-old C57BL/6N mice were intraperitoneally injected three-time per week with CCl<sub>4</sub> and corn oil (control group) during the whole experimental periods (Figure S1). After four weeks, the mice intraperitoneally injected with CCl<sub>4</sub> were randomly divided into three groups: CCl<sub>4</sub> and the 2 and 5 mg/kg DHE treatment groups. The aqueous solution of DHE was orally gavaged to the mice in the treatment groups. For vehicle treatment, water was provided by oral gavage to the mice in the corn oil and CCl<sub>4</sub> group (disease group).

It is well known that more serious the degree of liver fibrosis, the wider the portal vein and the slower its velocity. <sup>3</sup> The results demonstrated that the portal vein in the CCl<sub>4</sub> group was significantly wider than that in the corn oil group. Interestingly, the width of the portal vein in the 2 mg/kg DHE treatment group was significantly decreased (Figure 4A). Furthermore, the velocity of the portal vein in the treatment groups was also improved (Figure 4B). After eight weeks of treatment, <sup>5</sup> there were no significant differences in body weight, liver weight, spleen weight, or the ratio of liver weight to body weight among the four groups (Figure 4C and S2A). Gross liver specimens of the CCl<sub>4</sub> group were paler and had a rough appearance. However, it was back to normal appearance by DHE treatment compared to the CCl<sub>4</sub> group (Figure 4D). The level of plasma ALT of the CCl<sub>4</sub> group mice was significantly higher than corn oil group. Whereas, DHE treatment significantly decreased the level of ALT compared with CCl<sub>4</sub> group mice. In addition, the level of plasma AST was significantly decreased after 2mg/kg DHE treatment (Figure 4E). Collagen area were visualized by Masson's trichrome and quantified by using ImageJ 1.52a software. DHE treatment in both dosages significantly reduced collagen accumulation compared with those of the CCl<sub>4</sub> group (Figure 4F and G). Taken together, these results demonstrated that DHE significantly protected from liver fibrosis in CCl<sub>4</sub>-induced mouse fibrosis model.

DHE decreased the liver macrophages infiltration and ECM deposition. The infiltration of macrophages reflects the severity of liver inflammation. To investigate the degree of macrophages infiltration, the <sup>1</sup> liver immune cells were



extracted and labeled with CD45+, F4/80+, and CD11b+ antibodies followed by flow-cytometric analysis. The results demonstrated that the DHE treatment at both dosages significantly reduced proportion of CD11b+ cells (Figure 5A, B, and S2B–F). Consistent with the result of flow cytometry, the mRNA levels of TNF $\alpha$  and F4/80 which are two important indicators of liver inflammation significantly decreased upon DHE treatment (Figure 5C). On the other hand, the accumulation of ECM is another important sign of liver fibrosis. The results demonstrated that the mRNA levels of COL1A1, COL1A2, and  $\alpha$ -SMA significantly decreased in the DHE treatment group more than those of the CCl4 group (Figure 5D). The mRNA levels of COL3A1 also decreased in the DHE treatment groups (Figure 5D). Meanwhile, the protein levels of  $\alpha$ -SMA and p-SMAD3 were significantly decreased in the DHE treatment group compared to those of the CCl4 group (Figure 5E and F). Thus, we demonstrated here that, DHE significantly decreased the liver macrophages infiltration and ECM deposition.

## **DISCUSSION**

TGF $\beta$ 1 is regarded as the most potent fibrogenic cytokine, which activates TGF $\beta$ 2, followed by the recruitment of TGF $\beta$ 1, therefore triggers HSC activation<sup>[30]</sup>. In the current study, we aimed to screen drugs targeting TGF $\beta$ 2 and further blocking TGF $\beta$  down-stream signaling pathway from FDA-approved small molecule library. Among these six candidate drugs, DHE significantly decreased the protein and mRNA expression of fibrotic-related genes in LX-2 cells. The results of the affinity experiment demonstrated that DHE binds with TGF $\beta$ 2 at Leu27, Phe30, Thr51, Ser52, Ile53, and Glu55. DHE also significantly alleviated liver fibrosis by decreasing macrophages infiltration and ECM accumulation in CCl4-induced mouse fibrosis model. Thus, we demonstrate here first time repurposing of DHE, an anti-headache drug, to treat liver fibrosis.

Developing a new drug needs more than a decade and significant investment<sup>[8]</sup>. Drug repurposing is a potential tool to accelerate the drug discovery process, which has been employed to develop therapies for COVID-19<sup>[31]</sup>, antimicrobials<sup>[32]</sup>, and rare diseases<sup>[33]</sup>.



The chemical structure of DHE is similar to that of many natural neurotransmitters, including epinephrine, norepinephrine, dopamine, and serotonin<sup>[26]</sup>. It can modulate noradrenergic, serotonergic, and dopaminergic neurotransmission<sup>[34]</sup>. DHE is an adrenergic receptor antagonist used to treat severe orthostatic hypotension, migraine, and headache<sup>[26]</sup>, which can bind with various receptors. DHE is also an agonist of 5-HT1B, 5-HT1D, and 5-HT1F receptors, but it also binds to 5-HT1A and 5-HT2A receptors<sup>[26]</sup>. In the current study, surprisingly, we found that DHE also bind with TGFβR2 at reasonable affinity. Furthermore, we also identified specific binding sites by mutagenesis analysis. These results imply that DHE not only specifically targets 5-HT, but also TGFβR2. It further highlights wide-spread function of DHE in physiology / pathophysiology.

Continuous local and systemic inflammation aggravates liver injury, which is a critical factor in liver fibrosis. The innate immune system plays a pivotal role from the onset to the end stage of chronic liver disease<sup>[35]</sup>. Hepatic macrophages are considered as the first line of defense against pathogens, are a key cellular determinant in the process of fibrosis<sup>[36]</sup>. Bone marrow monocyte-derived macrophages and Kupffer cells are two distinct subsets of macrophages in the liver, which have been identified as key regulators of liver inflammation and key to the progression or regression of liver fibrosis. Kupffer cells, which are resident macrophages in liver tissue, exert anti-inflammatory effects<sup>[37]</sup>. Activated macrophages produce large amounts of TGFβ, which activates HSC into myofibroblast-like cells and synthesizes ECM<sup>[38]</sup>. This inflammatory response is also evident in the CCl<sub>4</sub>-induced mouse model, especially in the increased number of macrophages in liver. Consistent with previous results, we also found that the macrophage infiltration was significantly increased after CCl<sub>4</sub> treatment. We speculate that, upon binding to TGFβR2, DHE prevents recruitment of TGFβR1 to TGFβR2, thereby inhibiting TGFβ down-stream signaling pathway. Consistently, F4/80 and TNFα mRNA expression also significantly decreased upon DHE treatment. Taken together, DHE might reduce macrophage infiltration and finally prevent liver inflammation.

Activated HSCs secreting ECM represent a critical event in development of liver fibrosis<sup>[37]</sup>. In the fibrotic liver, type I and III collagens are deposited instead of laminins, type IV collagen, and proteoglycans in the normal liver<sup>[17]</sup>. Various mechanisms of HSC activation have been postulated, including TGF $\beta$ /SMAD pathway, Notch, Wnt/ $\beta$ -catenin, Hedgehog, and Hippo signaling<sup>[16]</sup>. In the current study, we revealed that DHE significantly reduced TGF $\beta$  induced HSCs activation in LX2 cellular model through specific blocking of TGF $\beta$  signaling pathway. It explained clearly the significant reduction of ECM by DHE treatment. Taken together, our data show that DHE alleviated liver fibrosis by binding to TGF $\beta$ R2, preventing the binding of TGF $\beta$ 1 and TGF $\beta$ R2, and blocking TGF $\beta$ 1 signaling to reduce liver inflammation.

## **CONCLUSION**

Our study demonstrated that DHE alleviated liver fibrosis while decreasing inflammation-related gene expression and HSC activation. The mechanism of its action is likely to be associated with decreased macrophages infiltration and ECM accumulation. Considering that liver fibrosis can be reversed, DHE might open-up new avenue to treat liver fibrosis in the future.

## **ARTICLE HIGHLIGHTS**

### ***Research background***

7 The transforming growth factor  $\beta$  (TGF $\beta$ ) signaling pathway plays a crucial role in the development of liver fibrosis by activating TGF $\beta$  type II receptor (TGF $\beta$ R2), followed by the recruitment of TGF $\beta$ R1 finally triggering downstream signaling pathway.

### ***Research motivation***

TGF $\beta$ R2 is considered an important target for developing drugs against liver fibrosis. Previous studies demonstrated that both inhibiting the expression of TGF $\beta$ R2 and exogenous extracellular domain of TGF $\beta$ R2 supplement effectively alleviated liver fibrosis.

### ***Research objectives***

To find drugs targeting TGF $\beta$ R2 that inhibit TGF $\beta$ R1/TGF $\beta$ R2 complex formation, theoretically inhibit its downstream TGF $\beta$  signaling pathway, and thereby ameliorate liver fibrosis, we screened drugs approved by the Food and Drug Administration (FDA) to identify potential TGF $\beta$ R2 blockers.

### ***Research methods***

FDA-approved drugs were screened for binding affinity with TGF $\beta$ R2 by virtual molecular docking. We identified 6 candidates and further explored their potential by Cell Counting Kit-8 (CCK-8) cell cytotoxic experiment to validate toxicity and titrated the best cellular working concentrations. Next, we further demonstrated the detailed molecular working mechanisms using mutagenesis analysis. Finally, we used a mouse model to investigate its potential anti-liver fibrosis effect.

### ***Research results***

Dihydroergotamine (DHE) shows great ability in reducing fibrotic gene expressions such as collagen, p-SMAD3, and  $\alpha$ -SMA in TGF $\beta$  induced cellular model of liver fibrosis in LX-2 cells. Furthermore, we demonstrated that DHE binds to TGF $\beta$ R2 with a Kd value of 17.64  $\mu$ M. In addition, DHE significantly improved liver fibrosis, as evidenced by Masson's trichrome staining of liver sections, the width and the velocity of the portal vein, and serum markers of liver function. In line with those observations, DHE also decreased macrophage infiltration and extracellular matrix deposition in the liver.

### ***Research conclusions***

DHE could alleviate liver fibrosis by binding to TGF $\beta$ R2 thereby suppressing its downstream TGF $\beta$  signaling pathway. We show here that as far as drug repurposing, DHE has great potential to treat liver fibrosis.

### *Research perspectives*

Considering that liver fibrosis can be reversed, DHE might open-up new avenue to treat liver fibrosis in the future.

# 4%

SIMILARITY INDEX

### PRIMARY SOURCES

|   |   |                 |
|---|---|-----------------|
| 1 | <a href="https://link.springer.com">link.springer.com</a><br>Internet   | 40 words — 1%   |
| 2 | <a href="https://www.ahmjournal.com">www.ahmjournal.com</a><br>Internet   | 39 words — 1%   |
| 3 | <a href="https://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a><br>Internet   | 38 words — 1%   |
| 4 | <a href="https://www.mdpi.com">www.mdpi.com</a><br>Internet   | 26 words — < 1% |
| 5 | Y Kishimoto. "Sulindac and a cyclooxygenase-2 inhibitor, etodolac, increase APC mRNA in the colon of rats treated with azoxymethane", Gut, 2000<br>Crossref | 20 words — < 1% |
| 6 | <a href="https://id.nii.ac.jp">id.nii.ac.jp</a><br>Internet   | 15 words — < 1% |
| 7 | <a href="https://www.scilit.net">www.scilit.net</a><br>Internet   | 15 words — < 1% |
| 8 | <a href="https://ir.library.louisville.edu">ir.library.louisville.edu</a><br>Internet   | 14 words — < 1% |
| 9 | <a href="https://diposit.ub.edu">diposit.ub.edu</a><br>Internet   | 13 words — < 1% |

---

EXCLUDE QUOTES            ON

EXCLUDE BIBLIOGRAPHY   ON

EXCLUDE SOURCES           < 12 WORDS

EXCLUDE MATCHES           < 12 WORDS