



All

Images

Videos

翻译成中文

关闭取词

42,100 Results

Any time ▼

## Rates and risk of hospitalisation among patients with type ...

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) › ... › Wiley-Blackwell Online Open

To investigate the rates and **risk** of hospitalisations **in patients with type 2 diabetes** (T2D) mellitus in England. Methods This **retrospective** population-based cohort **study** used computerised records from the General Practice Research Database linked to Hospital Episode Statistics data in England.

Cited by: 31

Author: J. M. Khalid, M. Raluy-Callado, B. H. Cur...

Publish Year: 2014

## Risk of disseminated intravascular coagulation in patients ...

[bmjopen.bmj.com](http://bmjopen.bmj.com) › Archive › Volume 7, Issue 1 ▼

Jan 01, 2017 · **Diabetes mellitus** (hereafter, diabetes) is characterised by fasting hyperglycaemia and is an established risk factor of atherothrombotic **disorders**. These affect the coronary, cerebral and peripheral **arterial** trees. Patients with diabetes have increased thrombotic tendency due to **platelet hyper-reactivity**.

Cited by: 1

Author: Kenichiro Nogami, Kenichiro Nogami, Isa...

Publish Year: 2017

## Risk Factors, Mortality, and Cardiovascular Outcomes in ...

<https://www.nejm.org/doi/full/10.1056/NEJMoa1800256>

**Risk Factors** and Cardiovascular Outcomes in **Type 2 Diabetes** In an observational **study**, **patients with type 2 diabetes** who had glycated hemoglobin, LDL cholesterol, albuminuria, and blood pressure ...

Cited by: 16

Author: Aidin Rawshani, Araz Rawshani, Stefan F...

Publish Year: 2018

## Increased Risk of Subsequent Myocardial ... - Diabetes Care

[care.diabetesjournals.org/content/37/5/1329](http://care.diabetesjournals.org/content/37/5/1329) ▼

## Match Overview

1 Internet 39 words  
crawled on 01-Jul-2016  
[www.medsci.org](http://www.medsci.org)

2%

**Name of Journal:** *World Journal of Diabetes*

**Manuscript NO:** 46616

**Manuscript Type:** ORIGINAL ARTICLE

*Retrospective Study*

**Risk factors in patients with type 2 diabetes in Bengaluru: A retrospective study**

**Running title:** Aravinda J, Risk Factors in Patients with T2DM in Bengaluru

Aravinda Jagadeesha

**Abstract**

**BACKGROUND**

Risk factors such as hereditary, ecological, and metabolic are interrelated and contribute to the development of type 2 diabetes mellitus. Family history of diabetes mellitus, age,



国内版

国际版

Risk factors in patients with type 2 diabetes in Bengaluru: A retrospective study Ri



All

Images

Videos

翻译成中文

关闭取词

748 Results

Any time ▾

### (PDF) Effectiveness and prescription pattern of lipid ...

[https://www.researchgate.net/publication/330462243\\_Effectiveness...](https://www.researchgate.net/publication/330462243_Effectiveness...)

Use of statin therapy in patients with type 2 diabetes mellitus (T2DM) has been recommended by most clinical guidelines. Cardiovascular disease (CVD) is a leading cause of morbidity and mortality ...

### MedWorm: Diabetic Ulcers Research

<https://medworm.com/research/diabetic-ulcers.xml>

The objectives of this study were to examine the effectiveness of different types of exercise on risk factors of diabetic foot ulcers, including glycated hemoglobin, peripheral arterial disease, and diabetic peripheral neuropathy, in people with type 2 diabetes mellitus.

### (PDF) Type 2 diabetes mellitus might be a risk factor for ...

[https://www.researchgate.net/publication/308753597\\_Type\\_2\\_diabetes...](https://www.researchgate.net/publication/308753597_Type_2_diabetes...)

Type 2 diabetes mellitus might be a risk factor for mild cognitive impairment progressing to Alzheimer's disease ... study of incidence and risk factors ... of type 2 diabetes mellitus (T2DM ...

### Journal of Education and Health Promotion: Browse articles

[www.jehp.net/browse.asp?date=0-2018](http://www.jehp.net/browse.asp?date=0-2018) ▾

Original Article: Periodontal therapy as an adjunctive modality for HbA1c reduction in type-2 diabetic patients Chalini Sundar, Sundar Ramalingam, Viswanathan Mohan, Rajendra Pradeepa, MJ ...

### CLINICAL VIGNETTES - - 2004 - Journal of General Internal ...

<https://onlinelibrary.wiley.com/doi/full/10.1111/j.1525-1497.2004...>

A retrospective review of all exposures reported to a poison control center revealed that 0.34% were hydrogen peroxide related. ... T-scores below -1.5 is the recommended level for therapy in patients with risk factors, and therapy should begin at T-scores below -2.0 in the absence of risk factors. ... are