

80473\_Auto\_Edited-check.docx

38

**Name of Journal:** *World Journal of Diabetes*

**Manuscript NO:** 80473

**Manuscript Type:** REVIEW

47

**Mesenchymal stem cell-derived exosomes: The dawn of diabetic wound healing**

Wu J *et al.* MSC-Exos for diabetic wounds

Jing Wu, Li-Hong Chen, Shi-Yi Sun, Yan Li, Xing-Wu Ran

### **Abstract**

Chronic wound healing has long been an unmet medical need in the field of wound repair, with diabetes being one of the major etiologies. Diabetic chronic wounds (DCWs), especially diabetic foot ulcers, are one of the most threatening chronic complications of diabetes. Although the treatment strategies, drugs, and dressings for diabetic chronic wounds have made great progress, they remain ineffective in some patients with refractory wounds. Stem cell-based therapies have achieved specific efficacy in various fields, with mesenchymal stem cells (MSCs) being the most widely used. Although MSCs have achieved good feedback in preclinical studies and clinical trials in the treatment of cutaneous wounds or other situations, the potential safety concerns associated with allogeneic/autologous stem cells and unknown long-term health effects need further attention and supervision. Recent studies have reported that stem cells mainly exert their trauma repair effects through paracrine secretion, and exosomes play an important role in intercellular communication as their main bioactive component. MSC-derived exosomes inherit the powerful inflammation and immune modulation, angiogenesis, cell proliferation, and migration promotion, oxidative stress alleviation, collagen remodeling imbalances regulation of their parental cells, and can avoid the potential risks of direct stem cell transplantation to a large extent, thus demonstrating promising performance as novel "cell-free" therapies in chronic wounds.

This review aimed to elucidate the potential mechanism and update the progress of MSC-derived exosomes in diabetic chronic wound healing, thereby providing new therapeutic directions for diabetic chronic wounds that are difficult to be cured using conventional therapy.

**Key Words:** Diabetic wounds; Wound and injuries; Mesenchymal stem cells; Exosomes; Pre-conditioning; Preclinical translation

Wu J, Chen LH, Sun SY, Li Y, Ran XW. Mesenchymal stem cell-derived exosomes: The dawn of diabetic wound healing. *World J Diabetes* 2022; In press

**Core Tip:** Diabetic chronic wounds are one of the most serious chronic complications of diabetes, and the efficacy of stem cell therapy for refractory chronic wounds has been studied previously. Stem cell-derived exosomes are one of the important active components of stem cell paracrine secretion, which inherit the wound repair capacity of parental cells as parts of novel cell-free therapies in addition to cell-based ones. Herein we discuss the mechanism and latest progress of mesenchymal stem cell-derived exosomes in promoting diabetic chronic wound healing.

## INTRODUCTION

Wound healing after skin tissue injury relies on a dynamic chain of physiological reactions including hemostasis, inflammation, cell proliferation, and tissue remodeling<sup>[1]</sup>. Any step out of balance, such as excessive inflammation, impaired fibroblast migration and proliferation, abnormal collagen formation and deposition, and hindered re-epithelialization, ultimately leads to delayed wound healing and formation of chronic wounds. Chronic wounds are those that have failed to proceed through an orderly and timely reparative process to produce anatomical and functional integrity of the injured site<sup>[2]</sup>. They refer to wounds caused by multiple factors that have not healed or have not demonstrated a tendency to heal after a certain period clinically,

with a chronic duration ranging from 4 to 12 wk<sup>[3,4]</sup>. Various pathological states result in chronic wound development, including diabetes, pressure injuries, infections, and arterial/venous insufficiency of which reports are similar in China and developed Western countries<sup>[4-6]</sup>, which have the most complicated pathogenesis and therapeutic strategies being diabetic chronic wounds (DCWs).

<sup>10</sup> Diabetes mellitus (DM) is a metabolic disease characterized by elevated blood glucose levels, of which DCWs are among the most threatening complications. The combination of a high-glucose environment and several biological <sup>37</sup> factors, including ischemia and hypoxia, abnormal inflammatory response, excessive oxidative stress, and peripheral neuropathy, contributes to wound formation<sup>[7-9]</sup>. Such wounds have problems of protracted healing, long treatment time, difficulties in management, high cost, repeated attacks, and high disability/mortality rates, resulting in heavy physical, psychological and economic burdens<sup>[10,11]</sup>. The intervention of DCWs cannot be underestimated based on what is mentioned above. Hence, solving persistent inflammation, impaired <sup>66</sup> cell proliferation and migration, decreased angiogenesis, and remodeling of the extracellular matrix (ECM) is important. Innovative wound repair methods, such as local negative pressure, growth factors, and autologous platelet-rich gels, have remarkable effects on healing DCWs<sup>[12-15]</sup>. However, more specific treatment options are required for refractory and contraindicated wounds.

With the rapid development of tissue engineering, cell therapies have gradually become widely used in various disciplines. <sup>51</sup> Stem cells can be used in regenerative <sup>60</sup> medicine and play an indispensable role in wound repair<sup>[16]</sup>, of which mesenchymal stem cells (MSCs) are the most commonly used. MSCs have self-renewal abilities and multi-directional differentiation potential, participating in damage repair through intercellular communication and bioactive factor secretion, finally achieving the effect of promoting wound healing<sup>[17]</sup>. Clinical trials of MSCs for treating various types of cutaneous wounds are currently in full swing, <sup>63</sup> and their efficacy and safety in promoting wound regeneration have been initially demonstrated. As clinical trials continue to progress, further attention and supervision need to be paid to their potential



safety issues of proliferative lesion formation, abnormal organ reaction and unknown long-term health effects after transplantation<sup>[18-20]</sup>.

Studies have revealed stem cells promote repair and regeneration mainly through paracrine signaling, whereas exosomes are one of their important paracrine active components<sup>[21]</sup>. MSC-derived exosomes (MSC-Exos) carry genetic information, functional RNAs, and proteins from parental cells, demonstrating wound healing effects *via* intercellular communication after these biologically active substances are acquired by recipient cells<sup>[22-24]</sup>. Thus, MSC-Exos have broad application prospects in diabetic wound repair<sup>[25]</sup>; however, they have not yet been carried out in clinical practice. The important <sup>42</sup>role of MSC-Exos in all stages of diabetic wound healing and the preclinical application are highlighted in this review, to pave the way for their use as an effective tool in the management of these harmful diabetic complications.

## **DIABETIC CHRONIC WOUNDS: HEALING DISORDERS CAUSED BY VARIOUS MECHANISMS**

<sup>10</sup>DM is a metabolic disease characterized by elevated blood glucose levels, which poses a serious threat to human health. The continuous progression of hyperglycemic toxicity without effective control will affect macrovascular, microvascular, and peripheral nerves throughout the body and involve various organs such as the brain, eyes, heart, kidney, and skin, resulting in various diabetic chronic complications<sup>[26]</sup>. DCWs are one of the most common and threatening chronic complications, often accompanied by infection or deep-tissue destruction<sup>[27]</sup>. Protracted wounds <sup>50</sup>are the most common cause of non-traumatic amputations. Diabetic foot ulcers (DFUs) are characterized by wounds on the feet, which are the most typical, and <sup>32</sup>patients with DFUs have a 2.5 times higher risk of 5-year mortality than those with none<sup>[28]</sup>. The overall mortality of DFUs within 5 years is nearly 50%<sup>[29]</sup>, <sup>11</sup>and approximately 20% of moderate-to-severe DFUs will lead to amputation; the 5-year mortality rate after amputation exceeds 70%<sup>[30]</sup>.

Impaired wound healing processes caused by hyperglycemia-induced disturbances in wound-linked cellular behaviors contribute to diabetic wound healing difficulties<sup>[7]</sup>.

<sup>31</sup>7. Hyperglycemia, oxidative stress, and insulin resistance affect the function of vascular smooth muscle cells, endothelial cells, and platelets, which in turn may lead to abnormal coagulation processes and affect platelets of triggering for subsequent inflammatory and proliferative phases<sup>[32]</sup>. The hyperglycemic microenvironment can lead to dysfunction of immune and inflammatory cells and dysregulation of inflammatory factors. Perpetuated inflammatory states induced by increased mast cell degranulation<sup>[33]</sup>, excessive extracellular traps produced by neutrophils<sup>[34]</sup>, dysregulated and persistent M1 (pro-inflammatory) macrophage polarization<sup>[35]</sup>, pro-inflammatory factors (IL-1 $\beta$ , TNF- $\alpha$ , and IL-6) overexpression, and anti-inflammatory factors (IL-10 and TGF- $\beta$ ) deficiency finally hinder wound healing<sup>[7]</sup>. The proliferative phase of diabetic wound healing is characterized by disturbed physiological functions of keratinocytes<sup>[36]</sup>, fibroblasts<sup>[37]</sup>, and endothelial cells<sup>[38]</sup>, then the impaired re-epithelialization, granulation tissue formation, matrix deposition, and angiogenesis affect wound healing. Various factors also affect the function and activity of these cells during this phase, including decreased chemokines with pro-angiogenesis produced by macrophages, hemoglobin glycation, vascular stenosis, increased oxygen consumption affecting oxygen-dependent cellular behaviors, and impaired nerve fiber regeneration<sup>[7,31,39,40]</sup>. Remodeling of the ECM spans the entire injury response, and fibroblasts are the major cell type responsible for this phase<sup>[31]</sup>. Sequential changes in the ECM require a balance between collagen degradation and synthesis, achieved through temporal regulation of the dynamic changes in the ratio of matrix metalloproteinases (MMPs) to tissue inhibitors of metalloproteinases (TIMPs)<sup>[41,42]</sup>. Such changes in DCWs are unbalanced and lead to difficult wound healing and excessive scarring<sup>[41,43]</sup>. However, no clear demarcation exists between the various stages of wound healing, and functionally impaired cells can interact, eventually leading to poor diabetic wound healing, progressing to local infection, gangrene, and even amputation. Therefore, the most important aspect of effectively treating DCWs is to identify an appropriate approach that can comprehensively improve abnormalities in all phases of wound healing.

## **CURRENT STRATEGIES AND PROMISING DIRECTIONS FOR DIABETIC CHRONIC WOUNDS REPAIR**

Traditional strategies for DCWs management include glycemic control, conventional dressings (*e.g.*, hydrocolloids, alginates, and silver ions, *etc.*), thorough debridement (*e.g.*, surgical, mechanical, ultrasonic waterjet, collagenase, and maggot, *etc.*), wound off-loading, autologous skin and skin substitute grafting, infection control, and revascularization, *etc.* These strategies are used to create the wound bed microenvironment suitable for repair through moisture balance maintenance, necrotic or inactivated tissues removal, systemic and local infections control, and local blood flow improvement<sup>[13,44-46]</sup>. Negative pressure wound therapy can also be used to achieve its role in improving wound exudate drainage, enhancing local perfusion, removing bacterial products, promoting granulation tissue growth, and facilitating wound healing<sup>[47]</sup>. <sup>22</sup> However, these conventional treatments are often ineffective in many patients because of impaired cell function around the wound sites caused by underlying microenvironmental alterations<sup>[48]</sup>. Several innovative wound adjuvant therapies, including exogenous supplementation of growth factors<sup>[49]</sup>, platelet-rich plasma<sup>[50]</sup>, autologous platelet-rich gels<sup>[15, 51]</sup>, and hyperbaric oxygen therapy<sup>[52]</sup> have been developed to promote the activity and function of damaged cells and offer the possibility of treating unselected refractory wounds. However, an updated systematic review has revealed that some measures had positive effects on accelerating wound healing, while others had limited impacts on diabetic ulcer healing<sup>[53]</sup>. However, the overall efficacy of various treatment modalities for DCWs remains unsatisfactory, and effective therapeutic strategies need to be continued.

## **STEM CELL-BASED THERAPIES BECOME HOT TOPICS, COMING EXOSOMES INTO BEING**

<sup>11</sup> Stem cells have the potential for self-renewal and multidirectional differentiation with great research and application value in life sciences, clinical trials and disease research.

Stem cell-based therapies are now approved by several countries, and have been widely used in various disciplines. MSCs are currently the main experimental cell sources and have shown their excellent therapeutic potential and value in clinical trials in the field of regenerative medicine<sup>[16,54]</sup>.

MSCs provide assistance in all phases of wound healing by exerting their functions of regulating skin homeostasis and wound healing through migration into the skin damage site and interaction with skin cells and can influence the function of these cells by paracrine secretion of bioactive factors and differentiation into them<sup>[55,56]</sup>. As MSCs have exhibited wound healing in many preclinical studies as powerful tools for regulating inflammation, promoting cell proliferation and migration, angiogenesis, and collagen synthesis<sup>[57-60]</sup>, the application of MSCs for DCWs contributes to progress toward clinical trials. Twenty-five clinical trials of MSCs for diabetic ulcers have been conducted or are recruiting subjects, which are recorded in the ClinicalTrials.gov database (clinicaltrials.gov).

Previous clinical studies have demonstrated that MSC transplantation in patients with DFUs is safe and feasible with the properties of improving microcirculation, wound healing, ulcer recurrence, and amputation<sup>[61-63]</sup>. However, <sup>5</sup> stem cell therapies are still in their early clinical stage, further attention and supervision are required of declined performance during production and application as cellular senescence and loss of multipotency during *ex vivo* expansion and from variable donors<sup>[64,65]</sup>, decreased survival rate caused by advanced glycosylation end products<sup>[66]</sup>, potential safety issues as proliferative lesion formation and abnormal organ reaction<sup>[20]</sup>, and unknown long-term health effects after transplantation. Basic and clinical researches related to allogeneic/autologous stem cells are subject to <sup>21</sup> the International Society for Stem Cell Research Guidelines for Clinical Translation of Stem Cells and national ethical guidelines and related guidelines/regulations<sup>[20,67]</sup>.

MSCs exert their repair and regenerative effects mainly through paracrine signaling, and exosomes are one of the important active components<sup>[21]</sup> that provide a more stable entity that minimizes the potential safety concerns for cell transplantation. MSC-Exos



13

play an important role in intercellular communication by carrying various important functional substances of parental cells, being used of promoting wound healing<sup>[68,69]</sup>. Compared to direct cell transplantation, MSC-Exos avoid the immune rejection because of low immunogenicity; allow to cross various biological barriers and avoid the risk of embolism from intravenous injection based on their smaller sizes<sup>[70]</sup>; the dose and fraction can be adjusted artificially and genetic modifications are easier and safer<sup>[71]</sup>; avoid the problem of malignant transformation; and allow to repair diabetic complications through multiple actions<sup>[72]</sup>. They can also be used as ideal carriers for carrying and delivering therapeutic drugs, genes, enzymes, or RNAs<sup>[73]</sup>, and their efficiency and targeted transport capacity can be tuned through pretreatment or engineering transformation<sup>[74]</sup>, demonstrating their promising applications in the field of repair and regeneration.

## **STEM CELL-DERIVED EXOSOMES: NOVEL CELL-FREE STRATEGIES**

### ***Exosomes biology***

44

The concept of “exosomes” was first proposed in 1981 by Trams *et al*<sup>[75]</sup>, using to collectively refer to extracellular vesicles (EVs) that originated from the exudation of various cell line cultures. The currently defined exosomes were first discovered in sheep reticulocytes and considered cellular waste<sup>[76-78]</sup>. Of note, “EVs” is the preferred term by the International Society for Extracellular Vesicles (ISEV) to describe all nanoparticles with lipid bilayer structures released by cells<sup>[79]</sup>.

33

Exosomes, the biological nanoscale spherical lipid bilayer vesicles<sup>[80]</sup>, can be secreted by almost all cell types and are widely present in cell culture supernatants and many body fluids<sup>[81]</sup>. Their diameters range from 10 to 200 nm. In addition to exosomes, EVs also include microvesicles that are also called ectosomes with a diameter of 100-1000 nm, and apoptotic bodies larger than 1000 nm according to different sizes and biogenesis<sup>[82,83]</sup>. The types and functions of the bioactive substances carried by exosomes differ according to their cellular origins and adjacent cellular components<sup>[84]</sup>. The major substances include genetic information, RNA species (mRNA, tRNA, rRNA, miRNA,

43

lncRNA, circRNA, etc.), proteins, lipids, cytokines, and growth factors<sup>[85,86]</sup>. Exosomal proteins include intrinsic components involved in exosome biogenesis, such as fusion-related proteins (GTPases, annexins, flotillin, and Rab proteins), heat shock proteins (HSP70 and HSP90), tetraspanins (CD63, CD81, CD82, and CD9), ESCRT complex, and specific functional proteins originating from parental cells<sup>[87]</sup>. Apart from serving as a medium for cellular communication, some proteins are also involved in the membrane composition and biosynthesis as identified biomarker proteins and can provide stability and permeability in concert with phospholipid bilayers.

Exosomes originate from endosomes during generation, circulation, degradation, and liberation<sup>[88]</sup>. Extracellular substances fuse with early sorting endosomes through plasma membrane invagination and endocytosis, and begin to accumulate bioactive substances. Eventually, they mature into late sorting endosomes, which invaginate to form intraluminal vesicles that can then generate multivesicular bodies (MVBs)<sup>[68,88]</sup>. MVBs can be absorbed by lysosomes comprising a degradative pathway, or they can undergo a specific exocytotic process whereby they fuse with the plasma membrane to release exosomes into the extracellular space<sup>[89]</sup>. After release, they act as mediators of intercellular and intra-organ communication to transfer the contained bioactive substances to recipient cells through direct fusion, endocytosis, and receptor-ligand binding to affect their functions<sup>[90,91]</sup>, participating in the body's physiological and pathological state adjustment<sup>[92]</sup>.

### *Isolation and characterization of exosomes*

The extraction of exosomes is primarily based on their physicochemical properties. This process is difficult because of the heterogeneity of exosomes derived from different cell origins, the possible existence of subpopulations of exosomes with different functions and phenotypes even when extracted from a single cell line, and multiple EV subtypes with similar biophysical properties<sup>[93]</sup>. Therefore, different isolation methods should be targeted for different purposes<sup>[87]</sup>. Differential ultracentrifugation is the most widely used separation technique and is also known as the gold standard for isolation, while



the main principle is to harvest the desired components based on size and density differences<sup>[94]</sup>. Polymer precipitation uses polyethylene glycol to harvest exosomes under centrifugal conditions by reducing their solubility<sup>[95]</sup>. Size-exclusion chromatography<sup>[96]</sup> and ultrafiltration<sup>[97]</sup> are both based on size differences between exosomes and other components, although they may adulterate other particles of similar size. Immunoaffinity capture is based on the specific binding of antibodies and ligands to isolate exosomes from a heterogeneous mixture<sup>[98]</sup>. Current isolation and purification techniques have varying effects and many problems such as low purity and recovery, structural damage, and time and cost consumption, making achieving efficient enrichment difficult, which has become a bottleneck of the translational applications of exosomes<sup>[87]</sup>. Hence, continuously exploring new isolation and purification techniques or combining multiple techniques is necessary to improve the isolation efficiency and thus obtain ideal exosomes.

Exosomes are mainly characterized by external characteristics (morphology and size detection) and the identification of surface markers<sup>[87]</sup>. As mentioned above, some protein components of exosomes serve as surface protein markers for identification. The ISEV has proposed the need to identify two types of proteins as follows: one is the biomarker proteins shared by exosomes to determine whether the extracted components are exosomes, and the other is cell-type-specific exosomal proteins that need to be identified to determine cellular origin<sup>[79]</sup>. Therefore, exosomes can be characterized by detecting their morphology using transmission electron microscopy, their size and concentration by dynamic light scattering, and nanoparticle tracking analysis technology, and their marker proteins by western blot, enzyme-linked immunoassay, and flow cytometry<sup>[87]</sup>.

### *MSCs derived exosomes and biological functions*

Stem cells have self-renewal abilities and multi-directional differentiation potential, while MSCs are one of the most frequently used and promising adult stem cells that can be derived from most adult tissues such as the bone marrow, adipose tissue, and

umbilical cord<sup>[99,100]</sup>. Bone marrow-derived MSC-Exos (BMSC-Exos) are biologically stable, have low immunogenicity, and exhibit good proliferation and viability after transplantation. They are most commonly used in clinical trials and can play a prominent role in various disorders, especially bone-related diseases<sup>[101]</sup>. Umbilical cord-derived MSC-Exos (UCMSC-Exos) can be isolated non-invasively, with low immunogenicity and strong self-renewal and proliferation ability, although it has limitations in maintaining bioactive and clinical therapeutic transport<sup>[102]</sup>. Adipose-derived MSC-Exos (AMSC-Exos) have relatively abundant sources that can be easily obtained by painless minimally invasive surgery; they are also pluripotent, plastic, easy to store, and stable in blood or body fluids<sup>[103]</sup>. Exosomes of different origins share most of their bioactive factors and are generally similar in their biological functions; however, their specific biological properties depend on the molecules that are specifically expressed<sup>[104]</sup>.

MSC-Exos are involved in intercellular communication through the transfer of proteins, RNA, DNA, and bioactive lipids that can be delivered to target cells to regulate their activities and functions<sup>[68]</sup>. They are generally involved in the regulation of cell survival and differentiation, the immune system, and inflammation modulation, and are also capable of promoting angiogenesis and tissue remodeling<sup>[73]</sup>. Considering these multiple biological functions, several studies have also reported that the MSC-Exos play a therapeutic role in autoimmune diseases<sup>[105]</sup>, ischemic injuries<sup>[106]</sup>, and metabolic diseases<sup>[107]</sup>, and are also related to dynamically modulating tumor biological functions<sup>[108]</sup>, promoting repair and regeneration of damaged osteochondral, neural, and tendon tissues, and facilitating wound healing<sup>[109-112]</sup>. Current studies also discovered that they can improve COVID-19-related cytokine storms and the deterioration of lung function due to severe pneumonia<sup>[113]</sup>.

## MESENCHYMAL STEM CELL-DERIVED EXOSOMES FOR REPAIRING DIABETIC WOUNDS

MSC-Exos<sup>30</sup> play an important role in each phase of wound healing<sup>[81]</sup>. They can regulate diverse cell types related to wound repair by enhancing or suppressing certain bioactivities, achieving hemostasis, inflammatory regulation,<sup>3</sup> cell migration to the wound site, cell proliferation, and differentiation to form granulation tissue, angiogenesis, and ECM reorganization<sup>[69]</sup>. They can also be expected to be therapeutic agents for different types of diabetes by alleviating autoimmune damages<sup>[114]</sup>, attenuating insulin resistance, and improving  $\beta$ -cell exhaustion<sup>[115]</sup>. Additionally, they can be used to prevent and treat DM-related complications. Based on these potentials, MSC-Exos may be of considerable importance in DCW treatment.

### *Hemostasis*

Tissue factor (TF) is an initiator of coagulation activation and was identified in the plasma membrane of exosomes<sup>[116]</sup>. TF can transfer to the platelets and<sup>15</sup> initiate the extrinsic coagulation cascade, leading to the conversion of prothrombin to thrombin and fibrin clot formation<sup>[117]</sup>. Induced coagulation and stimulated thrombogenicity were observed using<sup>3</sup> EVs carrying TF from the pericardial blood of patients who received cardiac surgery<sup>[116]</sup>. Rat BMSC-Exos were applied to the bleeding site in the hemorrhage liver model, which exhibited an inhibited amount of bleeding and shortened bleeding time, demonstrating their excellent hemostatic properties. However, no studies related to exosomes' promotion of coagulation in cutaneous wound healing have been conducted.<sup>29</sup> Further studies are needed to demonstrate the potential role of exosomes in the hemostasis phase of wound healing.

### *Inflammation*

Excessive inflammation is a major cause of persistent diabetic wounds. Abnormal macrophage polarization and cytokine overexpression lead to an uncontrolled and persistent inflammatory state and can cause secondary tissue damage<sup>[7]</sup>. MSCs-Exos can inhibit the differentiation, activation, and proliferation of T cells as well as reduce<sup>8</sup> IFN- $\gamma$  release<sup>[118]</sup>. They can reduce the concentration of the inflammatory cytokines, TNF- $\alpha$ ,<sup>18</sup>

iNOS, IL-1 $\beta$ , and IL-6<sup>[119]</sup> and upregulate the expression of the anti-inflammatory cytokine IL-10<sup>[120,121]</sup>. MSCs-Exos can also induce M2 polarization of macrophages to promote wound healing by delivering exosome-derived miR-223 to target regulating the expression of pknox1 protein<sup>[122]</sup>.

Such abilities can also be observed in diabetic wounds. Topical application of native AMSC-Exos to diabetic mice dorsal full-thickness skin wounds also downregulated inflammatory cytokines (IL-6, TNF- $\alpha$ , CD14, CD19, and CD68) expression and promoted wound healing<sup>[123]</sup>. Similar alleviated inflammatory effects achieved by regulating inflammatory factors could also be observed in the combination of intraperitoneal Nrf2 pharmaceutical activator and BMSC-Exos subcutaneous injection, demonstrating decreased inflammatory cytokines TNF- $\alpha$  and IL-1 $\beta$  and increased anti-inflammatory cytokines IL-4 and IL-10<sup>[124]</sup>. Intradermal injection of MSC-Exos derived from human menstrual blood could induce macrophage polarization from the M1 to M2 phenotype, while this capacity is better than that of menstrual blood-derived MSCs<sup>[125]</sup>. Significantly lower M1 polarized macrophages and higher M2 polarized macrophages were also observed in the diabetic mouse air pouch model and diabetic rat full-thickness skin wound model using BMSC-Exos, while melatonin-stimulated BMSC-Exos (MT-Exos) had stronger effects<sup>[121]</sup>. Immunomodulatory capacity was enhanced after preconditioning. Moreover, MT-Exos could improve wound healing by activating the PTEN/PI3K/AKT signaling pathway to promote macrophage M2 polarization, angiogenesis, and collagen synthesis; promote the resolution of persistent inflammation; and drive the transition from inflammation to proliferation<sup>[121]</sup>. HUCMSC-Exos pretreated with lipopolysaccharides have better regulatory properties for macrophage polarization and resolution of chronic inflammation by transferring miR-let7b, while the TLR4/NF- $\kappa$ B/STAT3/AKT pathway is important in regulating this mechanism to promote wound healing<sup>[126]</sup>. The use of engineered TNF- $\alpha$ /hypoxia-pretreated HUVMSC-Exos in infected DCWs also decreased proinflammatory cytokines (TNF- $\alpha$ , IL-1 $\beta$ , and IL-6), induced M2 macrophage polarization, reduced bacterial burden, and bacterial colonization at the wound sites. Reduced levels of oxidative



biomarkers and increased levels of antioxidant mediators also demonstrated the ability of oxidative stress suppression<sup>[127]</sup>. The combination of BMSC-Exos and carboxyethyl chitosan-dialdehyde carboxymethyl cellulose hydrogel revealed <sup>27</sup>skewed macrophage functional polarity from M1 toward an anti-inflammatory M2 phenotype, as well as enhanced antibacterial effects by significantly inhibiting bacterial growth<sup>[128]</sup>.

### ***Proliferation***

Fibroblasts, keratinocytes, and endothelial cells participate in the proliferative phase. Unlike the dual regulatory effects on the tumor, MSC-Exos directly affect <sup>45</sup>the proliferative phase of wound healing by stimulating the proliferation and differentiation of these cells, as well as promoting angiogenesis at injury sites<sup>[104]</sup>. Enhanced migratory and proliferative capacity and inhibited apoptosis of keratinocytes by activating the AKT/HIF-1 $\alpha$  and Wnt/ $\beta$ -catenin pathways were observed with AMSC-Exos<sup>[129,130]</sup>. BMSC-Exos demonstrated the ability to promote fibroblast proliferation, migration, and secretion of growth factors and can induce tube formation in human umbilical vein cells (HUVECs)<sup>[131]</sup>. AMSC-Exos induced angiogenesis <sup>34</sup>in both *in vivo* and *in vitro* experiments, and the promotion of angiogenesis in ECs was achieved by transferring <sup>4</sup>miR-125a to inhibit DLL4 expression, accompanied by the downregulation of pro-angiogenic genes (Ang1 and Flk1), and upregulation of anti-angiogenic genes (Vash1 and TSP1)<sup>[132]</sup>. In addition to its pro-proliferative ability *in vitro*, the pro-healing effect of MSC-Exos has also been observed in acute non-diabetic wounds. MSC-Exos from human umbilical cord Wharton's jelly could regulate HaCaT cell function by suppressing AIF nucleus translocation and PARP-1 hyperactivation, thus attenuating <sup>49</sup>full-thickness skin wounds by enhancing re-epithelialization and angiogenesis<sup>[133]</sup>. Fetal dermal-derived MSC-Exos accelerated <sup>2</sup>wound closure in a mouse full-thickness skin wound model by activating the Notch signaling pathway to promote the motility and secretory capacity of fibroblasts<sup>[134]</sup>.

Similarly, exosomes from MSCs improve proliferation and angiogenesis in diabetic wounds. AMSC-Exos <sup>4</sup>accelerated cutaneous wound healing in diabetic mice with full-

thickness skin wounds model by enhancing cell proliferation, inhibiting apoptosis, and promoting angiogenesis. They also repaired skin barrier functions, and produced large amounts, regular arrangement, and dense distribution of new collagen<sup>[123]</sup>. Shabbir *et al*<sup>[131]</sup> have also reported that these cells significantly increased their proliferation when treated with MSC-derived exosomes. Enhanced angiogenesis and fibroblasts proliferation, migration, and differentiation abilities were observed in diabetic wounds treated with human decidua derived MSC-Exos, as well as an improved fibroblast senescent state, reduced scar width, and larger and better-organized collagen deposition<sup>[135]</sup>.

Various methods have been used to modify MSC-Exos to enhance fibroblast proliferation and angiogenesis. Co-culture of lncRNA H19-transfected BMSC-Exos with fibroblasts extracted from foot tissue of patients with DFUs revealed that overexpressed exosomes regulated the PTEN-mediated PI3K/AKT signaling pathway by competitively binding miR-152-3p to enhance proliferation and migration of fibroblasts and inhibit apoptosis and inflammation<sup>[136]</sup>. Injecting such exosomes into the peri-wound tissue of diabetic mice revealed the same changes in expression and accelerated wound healing<sup>[136]</sup>. Atorvastatin-pretreated BMSC-Exos promoted proliferation, migration of HUVECs, and vascular endothelial growth factor (VEGF) expression and accelerated wound healing in diabetic full-thickness skin injury rat models<sup>[137]</sup>. Pioglitazone-pretreated BMSC-Exos-treated full-thickness wounds in diabetic rats achieved faster-wound closure, with more adequate re-epithelialization and extensive collagen deposition, significantly enhanced wound perfusion, and had significantly upregulated levels of VEGF and CD31<sup>[138]</sup>. Subcutaneous injection of mmu\_circ\_0000250-modified AMSC-Exos via miR-128-3p/SIRT1-mediated autophagy promoted wound healing in diabetic mice, and increased capillary and granulation tissue production was detected owing to promoted proliferation and migration and reduced apoptosis of endothelial cells<sup>[139]</sup>.

Biological scaffolds can improve the survival of exosomes in the inflammatory environment of diabetic wounds and maintain their sustained release. UCMSC-Exos



combined with the Pluronic F127 hydrogel revealed promoted chronic wound healing in diabetic mice. The elevated number of blood vessels and microvascular density, enhanced regeneration of granulation tissue, and cell proliferation were also observed, with the significant formation of new hair follicles in the center of the wounds, sufficient subepidermal collagen deposition, and orderly arrangement of collagen fibers<sup>[140]</sup>. Similar changes were observed in the wounds of diabetic mice using engineered bioactive self-healing antimicrobial exosome hydrogels (FHE@exo), and the elevated number of dermal appendages and differentiation and re-epithelialization of the epidermis were also observed<sup>[141]</sup>. The combination of human gingival tissue-derived MSC-Exos (GMSC-Exos) and a chitosan/silk hydrogel sponge promoted re-epithelialization, angiogenesis, and collagen deposition, while the increased nerve fiber density also reflected enhanced neuronal ingrowth in the proliferative stage<sup>[142]</sup>.

### **Matrix remodeling**

<sup>41</sup> In the final stage of wound healing, the production and remodeling of the ECM are key factors in determining the time of wound healing and degree of scarring. Recently, some studies have reported on the effects of exosomes on matrix remodeling. BMSC-Exos have been demonstrated to restore normal skin morphology in rats with full-thickness skin injury<sup>[143]</sup>, while these capacities relied on the downregulation of TGF- $\beta$ 1 and upregulation of TGF- $\beta$ 3 by inhibiting the TGF- $\beta$ /Smad signaling pathway. UCMSC-Exos had large amounts of miR-21, miR-23a, miR-125b, and miR-145, while it <sup>36</sup>inhibited the differentiation and excessive aggregation of myofibroblasts and exerted an anti-scarring effect *via* the TGF- $\beta$ 2/Smad2 pathway *in vivo*<sup>[144]</sup>. UCMSC-Exos can also promote the phosphorylation of YAP, a key site of the Hippo pathway, to negatively regulate the Wnt4/ $\beta$ -catenin pathway to balance <sup>56</sup>tissue regeneration and repair, with excessive cell proliferation and collagen deposition in the remodeling stage<sup>[145]</sup>. It was noted that intravenous injection of ADSC-Exos could increase the ratio of <sup>12</sup>type III collagen to type I and TGF- $\beta$ 3 to TGF- $\beta$ 1, prevent fibroblast-to-myofibroblast differentiation, and reduce scarring at incisions in the full-thickness skin injury

models<sup>[146]</sup>. They could also induce the ERK/MAPK pathway in fibroblasts to increase the expression of MMP3, thereby increasing MMP3/TIMP1 to regulate ECM remodeling<sup>[146]</sup>.

In contrast to the promoted cell proliferation and abundant granulation tissue in the early stage of healing, proliferative activities were reduced during the late repair stage to prohibit tissue hyperplasia when using FHE@exo, suggesting entry into the remodeling phase that prevents excessive tissue proliferation to promote wound healing<sup>[141]</sup>. The application of GMSC-Exos with chitosan/silk hydrogel sponge on the wounds of diabetic rats revealed more collagen deposition and thick wavy collagen fibers that were arranged in an orderly fashion, which is similar to that in normal skin, implying enhanced ECM remodeling<sup>[142]</sup>. These were also observed in the local transplantation of HUCMSC-Exos with polyvinyl alcohol/alginate nano hydrogel and of miR-126-3p overexpressed synovial-derived MSC-Exos with hydroxyapatite/chitosan composite hydrogel<sup>[147,148]</sup>. Altogether, these studies indicate that MSC-Exos play a pivotal role in the ECM remodeling phase of wound healing.

The various stages of wound healing are closely interwoven. MSC-Exos inherit the genetic information of their parental cells and can transfer the therapeutic bioactive substances to target cells to participate in intercellular communication, resulting in the regulation of target cell function and promotion of wound healing<sup>[81,149]</sup>. We analyzed the current preclinical application of MSC-Exos in diabetic wound models, and the cell source, administration method, dose, frequency, animal type, wound diameter, efficacy, and possible molecular mechanisms are summarized in Table 1. Additionally, MSC-Exos were not only responsible for a specific stage but also promote microenvironment changes in the wounds at each stage to exert a pro-healing effect. Although the biological functions of promoting diabetic wound healing are generally similar, certain differences exist in the regulated signaling pathways of different cell-derived exosomes or receiving different preconditioning, according to previous studies. The regulatory mechanisms most frequently studied in diabetic wound models and may potentially

confirmed in DCWs, as well as the microenvironmental changes in inflammatory and proliferative stages of wound healing after using MSC-Exos, are depicted in Figure 1.

## CURRENT STATUS AND PROSPECTS OF CLINICAL APPLICATIONS OF EXOSOMES IN DIABETIC CHRONIC WOUNDS

Preclinical studies have demonstrated the ability of MSC-Exos to promote diabetic wound healing. No evident pathological abnormalities in the heart, liver, spleen, lung, and kidneys sampled after exosome treatment were observed, and biomarkers reflecting liver and kidney function blood biochemistry were also within normal limits<sup>[127]</sup>. Meanwhile, no erythema, edema, or irritation was observed in the wound area after exosome treatment<sup>[137]</sup>, confirming the superior biosafety of exosome therapy.

We also searched for applications of exosomes secreted by stem cells from other sources in diabetic wounds and summarized them in Suppl Table S1. Noteworthy, the types of animals used for modeling were limited to mice and rats. Most of the studies involved acute diabetic wounds, that is, exosomes were administered immediately after successful modeling of full-thickness skin wounds. Only one study introduced *Staphylococcus aureus* to establish infected chronic wounds after the establishment of full-thickness cutaneous wounds and confirmed that exosomes were effective in treating infectious DCWs<sup>[127]</sup>. The efficacy and safety of MSC-Exos need to be further confirmed in larger animal models and DCW models. Because the islet morphology, structure and function, blood biochemical indices, and skin structure of minipigs are more similar to those of the human body, they are ideal animal models for studying diabetic wounds<sup>[159]</sup>. Our team has established a chronic skin ulcer model in diabetic miniature pigs in the early stage<sup>[160]</sup> and is researching on exosome products to explore the optimal administration methods and dosages and to verify their therapeutic effects.

According to the search results in ClinicalTrials.gov, no clinical trials of MSC-Exos and exosomes from other sources for diabetic cutaneous wound healing have been registered. Therefore, we expanded the scope of clinical trials to search for exosomes derived from any sources and exosome-enriched stem cell-conditioned medium in

various wound types (Table 2). None of the included four registered clinical trials had related results published, while they were all non-randomized one-arm pilot studies. Thus, more <sup>59</sup> high-quality randomized controlled trials are required to further confirm these research results. Of note, the application of cell-free therapies in clinical patients requires special attention to security, although no adverse reactions of exosomes have been reported in preclinical studies. Moreover, ADSC-Exos has been confirmed to not induce any irritation or toxicity in skin sensitization, irritation, or oral toxicity tests<sup>[161]</sup>; therefore, they can be considered in clinical practice to promote wound healing in combination with basic wound care measures. Nevertheless, toxicological analysis of different tissue-derived MSCs-Exos and more evidence of short and long-term health safety assessments are required to confirm their safety.

Exosome research is still in its infancy, and the realization of the transformation from preclinical research to clinical application still has great exploration value. The problems of optimal preparation, extraction, isolation, and storage of exosomes on a large scale and their production efficiency have not yet been determined; preparation and identification of components due to different source cells and the high heterogeneity of exosome components have not yet been solved; specific regulatory mechanisms in DCWs have not yet been fully elucidated; efficacy and safety of different cell sources and/or administrations have not been proven, and reasonable and effective methods of fusing exosomes with other biomaterials have not yet been implemented, all these issues are barriers that limit the clinical application of exosomes.

Thus, efficient, stable, safe, and mass-producible stem cells and related products for the treatment of diabetic wounds are yet to be explored and developed. More research is required in future clinical trials and routine practice to determine the most effective cell sources for diabetic wounds; to establish optimal large-scale culture conditions of MSCs; to solve the preparation problem of huge heterogeneity of exosome components; to explore standardized isolation, quality control, purification, and characterization techniques of MSC-Exos; and to determine the best approach for long-term storage<sup>[162]</sup>. Researchers also need to fully understand the abilities, loss, distribution, diffusion



efficiency, and clearance efficiency of exosomes after transporting them to target areas. Physical, chemical, or biological methods for preconditioning, genetic engineering, and transfection are used to specifically enhance a certain therapeutic potential to achieve relatively better wound healing than native exosomes, thus becoming new treatment directions<sup>[163]</sup>. Additionally, combining exosomes with biomaterials is possible to create bioactive dressings to enhance or combine repair ability, provide local microenvironment stability, and achieve sustained release of exosomes<sup>[74]</sup>. Additionally, starting clinical trials as soon as possible is necessary to verify the optimal dosages, administration methods, and efficacy evaluation of MSC-Exos in clinical patients, looking forward to its broad application prospects in promoting DCW healing in clinical practices<sup>[162]</sup>.

## **CONCLUSION**

DCWs, which are one of the most common chronic refractory wounds, pose a heavy burden to patients, families, and society. Current studies have suggested that MSC-Exos can play an important role in various aspects of wound healing and hold sufficient promise for promoting diabetic wound healing. However, recent clinical applications of MSC-Exos in DCW repair are still limited. Moreover, clinical translational issues, such as exosome production, isolation, purification, and storage processes, the most effective route of administration and dose, and efficacy evaluation remain. Accurate and efficient exosome products need to be established, and experiments in animals that have a greater resemblance to human skin tissues and clinical trials need to be initiated as soon as possible to validate the optimal dosage and administration, and efficacy evaluation for using MSC-Exos to provide safety assurance for further clinical applications. Modification of MSC-Exos and integration with biomaterials to improve their efficacy and reduce their elimination rate may be a promising direction. We look forward to the clinical application of MSC-Exos for diabetic wound healing.

## **REFERENCES**

- 1 **Broughton G 2nd**, Janis JE, Attinger CE. The basic science of wound healing. *Plast Reconstr Surg* 2006; **117**: 12S-34S [PMID: 16799372 DOI: 10.1097/01.prs.0000225430.42531.c2]
- 2 **Lazarus GS**, Cooper DM, Knighton DR, Percoraro RE, Rodeheaver G, Robson MC. Definitions and guidelines for assessment of wounds and evaluation of healing. *Wound Repair Regen* 1994; **2**: 165-170 [PMID: 17156107 DOI: 10.1046/j.1524-475X.1994.20305.x]
- 3 **Mustoe TA**, O'Shaughnessy K, Kloeters O. Chronic wound pathogenesis and current treatment strategies: a unifying hypothesis. *Plast Reconstr Surg* 2006; **117**: 35S-41S [PMID: 16799373 DOI: 10.1097/01.prs.0000225431.63010.1b]
- 4 **Frykberg RG**, Banks J. Challenges in the Treatment of Chronic Wounds. *Adv Wound Care (New Rochelle)* 2015; **4**: 560-582 [PMID: 26339534 DOI: 10.1089/wound.2015.0635]
- 5 **Martinengo L**, Olsson M, Bajpai R, Soljak M, Upton Z, Schmidtchen A, Car J, Järbrink K. Prevalence of chronic wounds in the general population: systematic review and meta-analysis of observational studies. *Ann Epidemiol* 2019; **29**: 8-15 [PMID: 30497932 DOI: 10.1016/j.annepidem.2018.10.005]
- 6 **Cheng B**, Jiang Y, Fu X, Hao D, Liu H, Liu Y, Huang Z, Tan Q, Wang L, Hu D, Yang Y, Han C, Cheng Z, Ran X, Li Y. Epidemiological characteristics and clinical analyses of chronic cutaneous wounds of inpatients in China: Prevention and control. *Wound Repair Regen* 2020; **28**: 623-630 [PMID: 32585756 DOI: 10.1111/wrr.12825]
- 7 **Patel S**, Srivastava S, Singh MR, Singh D. Mechanistic insight into diabetic wounds: Pathogenesis, molecular targets and treatment strategies to pace wound healing. *Biomed Pharmacother* 2019; **112**: 108615 [PMID: 30784919 DOI: 10.1016/j.biopha.2019.108615]
- 8 **Lim JZ**, Ng NS, Thomas C. Prevention and treatment of diabetic foot ulcers. *J R Soc Med* 2017; **110**: 104-109 [PMID: 28116957 DOI: 10.1177/0141076816688346]
- 9 **Chen L**, Gao Y, Li Y, Wang C, Chen D, Gao Y, Ran X. Severe Intermittent Hypoxia Modulates the Macrophage Phenotype and Impairs Wound Healing Through Downregulation of HIF-2 $\alpha$ . *Nat Sci Sleep* 2022; **14**: 1511-1520 [PMID: 36068885 DOI: 10.2147/NSS.S382275]



- 10 **Bowling FL**, Rashid ST, Boulton AJ. Preventing and treating foot complications associated with diabetes mellitus. *Nat Rev Endocrinol* 2015; **11**: 606-616 [PMID: 26284447 DOI: 10.1038/nrendo.2015.130]
- 11 **Kerr M**, Rayman G, Jeffcoate WJ. Cost of diabetic foot disease to the National Health Service in England. *Diabet Med* 2014; **31**: 1498-1504 [PMID: 24984759 DOI: 10.1111/dme.12545]
- 12 **Bowers S**, Franco E. Chronic Wounds: Evaluation and Management. *Am Fam Physician* 2020; **101**: 159-166 [PMID: 32003952]
- 13 **Jones RE**, Foster DS, Longaker MT. Management of Chronic Wounds-2018. *JAMA* 2018; **320**: 1481-1482 [PMID: 30326512 DOI: 10.1001/jama.2018.12426]
- 14 **Zarei F**, Negahdari B, Eatemadi A. Diabetic ulcer regeneration: stem cells, biomaterials, growth factors. *Artif Cells Nanomed Biotechnol* 2018; **46**: 26-32 [PMID: 28355923 DOI: 10.1080/21691401.2017.1304407]
- 15 **Li Y**, Gao Y, Gao Y, Chen D, Wang C, Liu G, Yang X, Ran X. Autologous platelet-rich gel treatment for diabetic chronic cutaneous ulcers: A meta-analysis of randomized controlled trials. *J Diabetes* 2019; **11**: 359-369 [PMID: 30182534 DOI: 10.1111/1753-0407.12850]
- 16 **Kanji S**, Das H. Advances of Stem Cell Therapeutics in Cutaneous Wound Healing and Regeneration. *Mediators Inflamm* 2017; **2017**: 5217967 [PMID: 29213192 DOI: 10.1155/2017/5217967]
- 17 **Lopes L**, Setia O, Aurshina A, Liu S, Hu H, Isaji T, Liu H, Wang T, Ono S, Guo X, Yatsula B, Guo J, Gu Y, Navarro T, Dardik A. Stem cell therapy for diabetic foot ulcers: a review of preclinical and clinical research. *Stem Cell Res Ther* 2018; **9**: 188 [PMID: 29996912 DOI: 10.1186/s13287-018-0938-6]
- 18 **Hyun I**, Lindvall O, Ahrlund-Richter L, Cattaneo E, Cavazzana-Calvo M, Cossu G, De Luca M, Fox IJ, Gerstle C, Goldstein RA, Hermerén G, High KA, Kim HO, Lee HP, Levy-Lahad E, Li L, Lo B, Marshak DR, McNab A, Munsie M, Nakauchi H, Rao M, Rooke HM, Valles CS, Srivastava A, Sugarman J, Taylor PL, Veiga A, Wong AL, Zoloth L, Daley GQ. New ISSCR guidelines underscore major principles for responsible

translational stem cell research. *Cell Stem Cell* 2008; **3**: 607-609 [PMID: 19041777 DOI: 10.1016/j.stem.2008.11.009]

19 **Lovell-Badge R**, Anthony E, Barker RA, Bubela T, Brivanlou AH, Carpenter M, Charo RA, Clark A, Clayton E, Cong Y, Daley GQ, Fu J, Fujita M, Greenfield A, Goldman SA, Hill L, Hyun I, Isasi R, Kahn J, Kato K, Kim JS, Kimmelman J, Knoblich JA, Mathews D, Montserrat N, Mosher J, Munsie M, Nakauchi H, Naldini L, Naughton G, Niakan K, Ogbogu U, Pedersen R, Rivron N, Rooke H, Rossant J, Round J, Saitou M, Sipp D, Steffann J, Sugarman J, Surani A, Takahashi J, Tang F, Turner L, Zettler PJ, Zhai X. ISSCR Guidelines for Stem Cell Research and Clinical Translation: The 2021 update. *Stem Cell Reports* 2021; **16**: 1398-1408 [PMID: 34048692 DOI: 10.1016/j.stemcr.2021.05.012]

20 **Marks PW**, Witten CM, Califf RM. Clarifying Stem-Cell Therapy's Benefits and Risks. *N Engl J Med* 2017; **376**: 1007-1009 [PMID: 27959704 DOI: 10.1056/NEJMp1613723]

21 **Nikfarjam S**, Rezaie J, Zolbanin NM, Jafari R. Mesenchymal stem cell derived-exosomes: a modern approach in translational medicine. *J Transl Med* 2020; **18**: 449 [PMID: 33246476 DOI: 10.1186/s12967-020-02622-3]

22 **Heldring N**, Mäger I, Wood MJ, Le Blanc K, Andaloussi SE. Therapeutic Potential of Multipotent Mesenchymal Stromal Cells and Their Extracellular Vesicles. *Hum Gene Ther* 2015; **26**: 506-517 [PMID: 26153722 DOI: 10.1089/hum.2015.072]

23 **Deng H**, Sun C, Sun Y, Li H, Yang L, Wu D, Gao Q, Jiang X. Lipid, Protein, and MicroRNA Composition Within Mesenchymal Stem Cell-Derived Exosomes. *Cell Reprogram* 2018; **20**: 178-186 [PMID: 29782191 DOI: 10.1089/cell.2017.0047]

24 **Joo HS**, Suh JH, Lee HJ, Bang ES, Lee JM. Current Knowledge and Future Perspectives on Mesenchymal Stem Cell-Derived Exosomes as a New Therapeutic Agent. *Int J Mol Sci* 2020; **21** [PMID: 31979113 DOI: 10.3390/ijms21030727]

25 **An T**, Chen Y, Tu Y, Lin P. Mesenchymal Stromal Cell-Derived Extracellular Vesicles in the Treatment of Diabetic Foot Ulcers: Application and Challenges. *Stem Cell Rev Rep* 2021; **17**: 369-378 [PMID: 32772239 DOI: 10.1007/s12015-020-10014-9]

- 26 **Demir S**, Nawroth PP, Herzig S, Ekim Üstünel B. Emerging Targets in Type 2 Diabetes and Diabetic Complications. *Adv Sci (Weinh)* 2021; **8**: e2100275 [PMID: 34319011 DOI: 10.1002/advs.202100275]
- 27 **Grennan D**. Diabetic Foot Ulcers. *JAMA* 2019; **321**: 114 [PMID: 30620372 DOI: 10.1001/jama.2018.18323]
- 28 **Saeedi P**, Petersohn I, Salpea P, Malanda B, Karuranga S, Unwin N, Colagiuri S, Guariguata L, Motala AA, Ogurtsova K, Shaw JE, Bright D, Williams R; IDF Diabetes Atlas Committee. Global and regional diabetes prevalence estimates for 2019 and projections for 2030 and 2045: Results from the International Diabetes Federation Diabetes Atlas, 9<sup>th</sup> edition. *Diabetes Res Clin Pract* 2019; **157**: 107843 [PMID: 31518657 DOI: 10.1016/j.diabres.2019.107843]
- 29 **Chen L**, Sun S, Gao Y, Ran X. Global mortality of diabetic foot ulcer: A systematic review and meta-analysis of observational studies. *Diabetes Obes Metab* 2022 [PMID: 36054820 DOI: 10.1111/dom.14840]
- 30 **Walsh JW**, Hoffstad OJ, Sullivan MO, Margolis DJ. Association of diabetic foot ulcer and death in a population-based cohort from the United Kingdom. *Diabet Med* 2016; **33**: 1493-1498 [PMID: 26666583 DOI: 10.1111/dme.13054]
- 31 **Wilkinson HN**, Hardman MJ. Wound healing: cellular mechanisms and pathological outcomes. *Open Biol* 2020; **10**: 200223 [PMID: 32993416 DOI: 10.1098/rsob.200223]
- 32 **Kaur R**, Kaur M, Singh J. Endothelial dysfunction and platelet hyperactivity in type 2 diabetes mellitus: molecular insights and therapeutic strategies. *Cardiovasc Diabetol* 2018; **17**: 121 [PMID: 30170601 DOI: 10.1186/s12933-018-0763-3]
- 33 **Dong J**, Chen L, Zhang Y, Jayaswal N, Mezghani I, Zhang W, Veves A. Mast Cells in Diabetes and Diabetic Wound Healing. *Adv Ther* 2020; **37**: 4519-4537 [PMID: 32935286 DOI: 10.1007/s12325-020-01499-4]
- 34 **Wong SL**, Demers M, Martinod K, Gallant M, Wang Y, Goldfine AB, Kahn CR, Wagner DD. Diabetes primes neutrophils to undergo NETosis, which impairs wound healing. *Nat Med* 2015; **21**: 815-819 [PMID: 26076037 DOI: 10.1038/nm.3887]

- 35 **Louiselle AE**, Niemiec SM, Zgheib C, Liechty KW. Macrophage polarization and diabetic wound healing. *Transl Res* 2021; **236**: 109-116 [PMID: 34089902 DOI: 10.1016/j.trsl.2021.05.006]
- 36 **Hu SC**, Lan CE. High-glucose environment disturbs the physiologic functions of keratinocytes: Focusing on diabetic wound healing. *J Dermatol Sci* 2016; **84**: 121-127 [PMID: 27461757 DOI: 10.1016/j.jdermsci.2016.07.008]
- 37 **Liu Y**, Liu Y, He W, Mu X, Wu X, Deng J, Nie X. Fibroblasts: Immunomodulatory factors in refractory diabetic wound healing. *Front Immunol* 2022; **13**: 918223 [PMID: 35990622 DOI: 10.3389/fimmu.2022.918223]
- 38 **Okonkwo UA**, DiPietro LA. Diabetes and Wound Angiogenesis. *Int J Mol Sci* 2017; **18** [PMID: 28671607 DOI: 10.3390/ijms18071419]
- 39 **Brem H**, Tomic-Canic M. Cellular and molecular basis of wound healing in diabetes. *J Clin Invest* 2007; **117**: 1219-1222 [PMID: 17476353 DOI: 10.1172/JCI32169]
- 40 **Papachristou S**, Pafili K, Papanas N. Skin AGEs and diabetic neuropathy. *BMC Endocr Disord* 2021; **21**: 28 [PMID: 33622304 DOI: 10.1186/s12902-021-00697-7]
- 41 **Davis FM**, Kimball A, Boniakowski A, Gallagher K. Dysfunctional Wound Healing in Diabetic Foot Ulcers: New Crossroads. *Curr Diab Rep* 2018; **18**: 2 [PMID: 29362914 DOI: 10.1007/s11892-018-0970-z]
- 42 **Li L**, Chen D, Wang C, Liu G, Ran X. The Effect of Autologous Platelet-Rich Gel on the Dynamic Changes of the Matrix Metalloproteinase-2 and Tissue Inhibitor of Metalloproteinase-2 Expression in the Diabetic Chronic Refractory Cutaneous Ulcers. *J Diabetes Res* 2015; **2015**: 954701 [PMID: 26221614 DOI: 10.1155/2015/954701]
- 43 **Chang M**. Restructuring of the extracellular matrix in diabetic wounds and healing: A perspective. *Pharmacol Res* 2016; **107**: 243-248 [PMID: 27033051 DOI: 10.1016/j.phrs.2016.03.008]
- 44 **Everett E**, Mathioudakis N. Update on management of diabetic foot ulcers. *Ann N Y Acad Sci* 2018; **1411**: 153-165 [PMID: 29377202 DOI: 10.1111/nyas.13569]

- 45 **Powers JG**, Higham C, Broussard K, Phillips TJ. Wound healing and treating wounds: Chronic wound care and management. *J Am Acad Dermatol* 2016; **74**: 607-25; quiz 625-6 [PMID: 26979353 DOI: 10.1016/j.jaad.2015.08.070]
- 46 **Braun L**, Kim PJ, Margolis D, Peters EJ, Lavery LA; Wound Healing Society. What's new in the literature: an update of new research since the original WHS diabetic foot ulcer guidelines in 2006. *Wound Repair Regen* 2014; **22**: 594-604 [PMID: 25139424 DOI: 10.1111/wrr.12220]
- 47 **Ji S**, Liu X, Huang J, Bao J, Chen Z, Han C, Hao D, Hong J, Hu D, Jiang Y, Ju S, Li H, Li Z, Liang G, Liu Y, Luo G, Lv G, Ran X, Shi Z, Tang J, Wang A, Wang G, Wang J, Wang X, Wen B, Wu J, Xu H, Xu M, Ye X, Yuan L, Zhang Y, Xiao S, Xia Z. Consensus on the application of negative pressure wound therapy of diabetic foot wounds. *Burns Trauma* 2021; **9**: tkab018 [PMID: 34212064 DOI: 10.1093/burnst/tkab018]
- 48 **Pop MA**, Almquist BD. Biomaterials: A potential pathway to healing chronic wounds? *Exp Dermatol* 2017; **26**: 760-763 [PMID: 28094868 DOI: 10.1111/exd.13290]
- 49 **Zubair M**, Ahmad J. Role of growth factors and cytokines in diabetic foot ulcer healing: A detailed review. *Rev Endocr Metab Disord* 2019; **20**: 207-217 [PMID: 30937614 DOI: 10.1007/s11154-019-09492-1]
- 50 **Mastrogiacomo M**, Nardini M, Collina MC, Di Campli C, Filaci G, Cancedda R, Odorisio T. Innovative Cell and Platelet Rich Plasma Therapies for Diabetic Foot Ulcer Treatment: The Allogeneic Approach. *Front Bioeng Biotechnol* 2022; **10**: 869408 [PMID: 35586557 DOI: 10.3389/fbioe.2022.869408]
- 51 **Li L**, Chen D, Wang C, Yuan N, Wang Y, He L, Yang Y, Chen L, Liu G, Li X, Ran X. Autologous platelet-rich gel for treatment of diabetic chronic refractory cutaneous ulcers: A prospective, randomized clinical trial. *Wound Repair Regen* 2015; **23**: 495-505 [PMID: 25847503 DOI: 10.1111/wrr.12294]
- 52 **Zhang Z**, Zhang W, Xu Y, Liu D. Efficacy of hyperbaric oxygen therapy for diabetic foot ulcers: An updated systematic review and meta-analysis. *Asian J Surg* 2022; **45**: 68-78 [PMID: 34376365 DOI: 10.1016/j.asjsur.2021.07.047]

- 53 **Vas P**, Rayman G, Dhatariya K, Driver V, Hartemann A, Londahl M, Piaggese A, Apelqvist J, Attinger C, Game F. Effectiveness of interventions to enhance healing of chronic foot ulcers in diabetes: a systematic review. *Diabetes Metab Res Rev* 2020; **36** **Suppl 1**: e3284 [PMID: 32176446 DOI: 10.1002/dmrr.3284]
- 54 **Margiana R**, Markov A, Zekiy AO, Hamza MU, Al-Dabbagh KA, Al-Zubaidi SH, Hameed NM, Ahmad I, Sivaraman R, Kzar HH, Al-Gazally ME, Mustafa YF, Siahmansouri H. Clinical application of mesenchymal stem cell in regenerative medicine: a narrative review. *Stem Cell Res Ther* 2022; **13**: 366 [PMID: 35902958 DOI: 10.1186/s13287-022-03054-0]
- 55 **Mazini L**, Rochette L, Admou B, Amal S, Malka G. Hopes and Limits of Adipose-Derived Stem Cells (ADSCs) and Mesenchymal Stem Cells (MSCs) in Wound Healing. *Int J Mol Sci* 2020; **21** [PMID: 32075181 DOI: 10.3390/ijms21041306]
- 56 **Dekoninck S**, Blanpain C. Stem cell dynamics, migration and plasticity during wound healing. *Nat Cell Biol* 2019; **21**: 18-24 [PMID: 30602767 DOI: 10.1038/s41556-018-0237-6]
- 57 **Yu Q**, Qiao GH, Wang M, Yu L, Sun Y, Shi H, Ma TL. Stem Cell-Based Therapy for Diabetic Foot Ulcers. *Front Cell Dev Biol* 2022; **10**: 812262 [PMID: 35178389 DOI: 10.3389/fcell.2022.812262]
- 58 **Maranda EL**, Rodriguez-Menocal L, Badiavas EV. Role of Mesenchymal Stem Cells in Dermal Repair in Burns and Diabetic Wounds. *Curr Stem Cell Res Ther* 2017; **12**: 61-70 [PMID: 27412677 DOI: 10.2174/1574888x11666160714115926]
- 59 **Shi R**, Lian W, Jin Y, Cao C, Han S, Yang X, Zhao S, Li M, Zhao H. Role and effect of vein-transplanted human umbilical cord mesenchymal stem cells in the repair of diabetic foot ulcers in rats. *Acta Biochim Biophys Sin (Shanghai)* 2020; **52**: 620-630 [PMID: 32484226 DOI: 10.1093/abbs/gmaa039]
- 60 **De Gregorio C**, Contador D, Díaz D, Cárcamo C, Santapau D, Lobos-Gonzalez L, Acosta C, Campero M, Carpio D, Gabriele C, Gaspari M, Aliaga-Tobar V, Maracaja-Coutinho V, Ezquer M, Ezquer F. Human adipose-derived mesenchymal stem cell-conditioned medium ameliorates polyneuropathy and foot ulceration in diabetic BKS



db/db mice. *Stem Cell Res Ther* 2020; **11**: 168 [PMID: 32357914 DOI: 10.1186/s13287-020-01680-0]

61 **Kirana S**, Stratmann B, Prante C, Prohaska W, Koerperich H, Lammers D, Gastens MH, Quast T, Negrean M, Stirban OA, Nandreaan SG, Götting C, Minartz P, Kleesiek K, Tschoepe D. Autologous stem cell therapy in the treatment of limb ischaemia induced chronic tissue ulcers of diabetic foot patients. *Int J Clin Pract* 2012; **66**: 384-393 [PMID: 22284892 DOI: 10.1111/j.1742-1241.2011.02886.x]

62 **Lu D**, Jiang Y, Deng W, Zhang Y, Liang Z, Wu Q, Jiang X, Zhang L, Gao F, Cao Y, Chen B, Xue Y. Long-Term Outcomes of BMMSC Compared with BMMNC for Treatment of Critical Limb Ischemia and Foot Ulcer in Patients with Diabetes. *Cell Transplant* 2019; **28**: 645-652 [PMID: 30917698 DOI: 10.1177/0963689719835177]

63 **Moon KC**, Suh HS, Kim KB, Han SK, Young KW, Lee JW, Kim MH. Potential of Allogeneic Adipose-Derived Stem Cell-Hydrogel Complex for Treating Diabetic Foot Ulcers. *Diabetes* 2019; **68**: 837-846 [PMID: 30679183 DOI: 10.2337/db18-0699]

64 **Rombouts WJ**, Ploemacher RE. Primary murine MSC show highly efficient homing to the bone marrow but lose homing ability following culture. *Leukemia* 2003; **17**: 160-170 [PMID: 12529674 DOI: 10.1038/sj.leu.2402763]

65 **Siddappa R**, Licht R, van Blitterswijk C, de Boer J. Donor variation and loss of multipotency during *in vitro* expansion of human mesenchymal stem cells for bone tissue engineering. *J Orthop Res* 2007; **25**: 1029-1041 [PMID: 17469183 DOI: 10.1002/jor.20402]

66 **Wang Z**, Li H, Zhang D, Liu X, Zhao F, Pang X, Wang Q. Effect of advanced glycosylation end products on apoptosis in human adipose tissue-derived stem cells *in vitro*. *Cell Biosci* 2015; **5**: 3 [PMID: 25973170 DOI: 10.1186/2045-3701-5-3]

67 **Jin J**. Stem Cell Treatments. *JAMA* 2017; **317**: 330 [PMID: 28114555 DOI: 10.1001/jama.2016.17822]

68 **Kalluri R**, LeBleu VS. The biology, function, and biomedical applications of exosomes. *Science* 2020; **367** [PMID: 32029601 DOI: 10.1126/science.aau6977]

- 69 **Than UTT**, Guanzon D, Leavesley D, Parker T. Association of Extracellular Membrane Vesicles with Cutaneous Wound Healing. *Int J Mol Sci* 2017; **18** [PMID: 28468315 DOI: 10.3390/ijms18050956]
- 70 **Nair S**, Salomon C. Extracellular vesicles and their immunomodulatory functions in pregnancy. *Semin Immunopathol* 2018; **40**: 425-437 [PMID: 29616307 DOI: 10.1007/s00281-018-0680-2]
- 71 **Lamichhane TN**, Sokic S, Schardt JS, Raiker RS, Lin JW, Jay SM. Emerging roles for extracellular vesicles in tissue engineering and regenerative medicine. *Tissue Eng Part B Rev* 2015; **21**: 45-54 [PMID: 24957510 DOI: 10.1089/ten.TEB.2014.0300]
- 72 **Newton WC**, Kim JW, Luo JZQ, Luo L. Stem cell-derived exosomes: a novel vector for tissue repair and diabetic therapy. *J Mol Endocrinol* 2017; **59**: R155-R165 [PMID: 28835418 DOI: 10.1530/JME-17-0080]
- 73 **Xunian Z**, Kalluri R. Biology and therapeutic potential of mesenchymal stem cell-derived exosomes. *Cancer Sci* 2020; **111**: 3100-3110 [PMID: 32639675 DOI: 10.1111/cas.14563]
- 74 **Phan J**, Kumar P, Hao D, Gao K, Farmer D, Wang A. Engineering mesenchymal stem cells to improve their exosome efficacy and yield for cell-free therapy. *J Extracell Vesicles* 2018; **7**: 1522236 [PMID: 30275938 DOI: 10.1080/20013078.2018.1522236]
- 75 **Trams EG**, Lauter CJ, Salem N Jr, Heine U. Exfoliation of membrane ecto-enzymes in the form of micro-vesicles. *Biochim Biophys Acta* 1981; **645**: 63-70 [PMID: 6266476 DOI: 10.1016/0005-2736(81)90512-5]
- 76 **Johnstone RM**, Adam M, Hammond JR, Orr L, Turbide C. Vesicle formation during reticulocyte maturation. Association of plasma membrane activities with released vesicles (exosomes). *J Biol Chem* 1987; **262**: 9412-9420 [PMID: 3597417]
- 77 **Pan BT**, Johnstone RM. Fate of the transferrin receptor during maturation of sheep reticulocytes in vitro: selective externalization of the receptor. *Cell* 1983; **33**: 967-978 [PMID: 6307529 DOI: 10.1016/0092-8674(83)90040-5]

78 **Pan BT**, Teng K, Wu C, Adam M, Johnstone RM. Electron microscopic evidence for externalization of the transferrin receptor in vesicular form in sheep reticulocytes. *J Cell Biol* 1985; **101**: 942-948 [PMID: 2993317 DOI: 10.1083/jcb.101.3.942]

79 **Théry C**, Witwer KW, Aikawa E, Alcaraz MJ, Anderson JD, Andriantsitohaina R, Antoniou A, Arab T, Archer F, Atkin-Smith GK, Ayre DC, Bach JM, Bachurski D, Baharvand H, Balaj L, Baldacchino S, Bauer NN, Baxter AA, Bebawy M, Beckham C, Bedina Zavec A, Benmoussa A, Berardi AC, Bergese P, Bielska E, Blenkiron C, Bobis-Wozowicz S, Boilard E, Boireau W, Bongiovanni A, Borràs FE, Bosch S, Boulanger CM, Breakefield X, Breglio AM, Brennan MÁ, Brigstock DR, Brisson A, Broekman ML, Bromberg JF, Bryl-Górecka P, Buch S, Buck AH, Burger D, Busatto S, Buschmann D, Bussolati B, Buzás EI, Byrd JB, Camussi G, Carter DR, Caruso S, Chamley LW, Chang YT, Chen C, Chen S, Cheng L, Chin AR, Clayton A, Clerici SP, Cocks A, Cocucci E, Coffey RJ, Cordeiro-da-Silva A, Couch Y, Coumans FA, Coyle B, Crescitelli R, Criado MF, D'Souza-Schorey C, Das S, Datta Chaudhuri A, de Candia P, De Santana EF, De Wever O, Del Portillo HA, Demaret T, Deville S, Devitt A, Dhondt B, Di Vizio D, Dieterich LC, Dolo V, Dominguez Rubio AP, Dominici M, Dourado MR, Driedonks TA, Duarte FV, Duncan HM, Eichenberger RM, Ekström K, El Andaloussi S, Elie-Caille C, Erdbrügger U, Falcón-Pérez JM, Fatima F, Fish JE, Flores-Bellver M, Försonits A, Frelet-Barrand A, Fricke F, Fuhrmann G, Gabrielsson S, Gámez-Valero A, Gardiner C, Gärtner K, Gaudin R, Gho YS, Giebel B, Gilbert C, Gimona M, Giusti I, Goberdhan DC, Görgens A, Gorski SM, Greening DW, Gross JC, Gualerzi A, Gupta GN, Gustafson D, Handberg A, Haraszti RA, Harrison P, Hegyesi H, Hendrix A, Hill AF, Hochberg FH, Hoffmann KF, Holder B, Holthofer H, Hosseinkhani B, Hu G, Huang Y, Huber V, Hunt S, Ibrahim AG, Ikezu T, Inal JM, Isin M, Ivanova A, Jackson HK, Jacobsen S, Jay SM, Jayachandran M, Jenster G, Jiang L, Johnson SM, Jones JC, Jong A, Jovanovic-Talisman T, Jung S, Kalluri R, Kano SI, Kaur S, Kawamura Y, Keller ET, Khamari D, Khomyakova E, Khvorova A, Kierulf P, Kim KP, Kislinger T, Klingeborn M, Klinke DJ 2nd, Kornek M, Kosanović MM, Kovács ÁF, Krämer-Albers EM, Krasemann S, Krause M, Kurochkin IV, Kusuma GD, Kuypers S, Laitinen S, Langevin SM, Languino LR, Lannigan J, Lässer C,

Laurent LC, Lavieu G, Lázaro-Ibáñez E, Le Lay S, Lee MS, Lee YXF, Lemos DS, Lenassi M, Leszczynska A, Li IT, Liao K, Libregts SF, Ligeti E, Lim R, Lim SK, Linē A, Linnemannstöns K, Llorente A, Lombard CA, Lorenowicz MJ, Lörincz ÁM, Lötvall J, Lovett J, Lowry MC, Loyer X, Lu Q, Lukomska B, Lunavat TR, Maas SL, Malhi H, Marcilla A, Mariani J, Mariscal J, Martens-Uzunova ES, Martin-Jaular L, Martinez MC, Martins VR, Mathieu M, Mathivanan S, Maugeri M, McGinnis LK, McVey MJ, Meckes DG Jr, Meehan KL, Mertens I, Minciocchi VR, Möller A, Møller Jørgensen M, Morales-Kastresana A, Morhayim J, Mullier F, Muraca M, Musante L, Mussack V, Muth DC, Myburgh KH, Najrana T, Nawaz M, Nazarenko I, Nejsun P, Neri C, Neri T, Nieuwland R, Nimrichter L, Nolan JP, Nolte-'t Hoen EN, Noren Hooten N, O'Driscoll L, O'Grady T, O'Loghlen A, Ochiya T, Olivier M, Ortiz A, Ortiz LA, Osteikoetxea X, Østergaard O, Ostrowski M, Park J, Pegtel DM, Peinado H, Perut F, Pfaffl MW, Phinney DG, Pieters BC, Pink RC, Pisetsky DS, Pogge von Strandmann E, Polakovicova I, Poon IK, Powell BH, Prada I, Pulliam L, Quesenberry P, Radeghieri A, Raffai RL, Raimondo S, Rak J, Ramirez MI, Raposo G, Rayyan MS, Regev-Rudzki N, Ricklefs FL, Robbins PD, Roberts DD, Rodrigues SC, Rohde E, Rome S, Rouschop KM, Ruggetti A, Russell AE, Saá P, Sahoo S, Salas-Huenuleo E, Sánchez C, Saugstad JA, Saul MJ, Schiffelers RM, Schneider R, Schøyen TH, Scott A, Shahaj E, Sharma S, Shatnyeva O, Shekari F, Shelke GV, Shetty AK, Shiba K, Siljander PR, Silva AM, Skowronek A, Snyder OL 2nd, Soares RP, Sódar BW, Soekmadji C, Sotillo J, Stahl PD, Stoorvogel W, Stott SL, Strasser EF, Swift S, Tahara H, Tewari M, Timms K, Tiwari S, Tixeira R, Tkach M, Toh WS, Tomasini R, Torrecilhas AC, Tosar JP, Toxavidis V, Urbanelli L, Vader P, van Balkom BW, van der Grein SG, Van Deun J, van Herwijnen MJ, Van Keuren-Jensen K, van Niel G, van Royen ME, van Wijnen AJ, Vasconcelos MH, Vechetti IJ Jr, Veit TD, Vella LJ, Velot É, Verweij FJ, Vestad B, Viñas JL, Visnovitz T, Vukman KV, Wahlgren J, Watson DC, Wauben MH, Weaver A, Webber JP, Weber V, Wehman AM, Weiss DJ, Welsh JA, Wendt S, Wheelock AM, Wiener Z, Witte L, Wolfram J, Xagorari A, Xander P, Xu J, Yan X, Yáñez-Mó M, Yin H, Yuana Y, Zappulli V, Zarubova J, Žekas V, Zhang JY, Zhao Z, Zheng L, Zheutlin AR, Zickler AM, Zimmermann P, Zivkovic AM, Zocco D, Zuba-Surma EK. Minimal

information for studies of extracellular vesicles 2018 (MISEV2018): a position statement of the International Society for Extracellular Vesicles and update of the MISEV2014 guidelines. *J Extracell Vesicles* 2018; **7**: 1535750 [PMID: 30637094 DOI: 10.1080/20013078.2018.1535750]

80 **Farooqi AA**, Desai NN, Qureshi MZ, Librelotto DRN, Gasparri ML, Bishayee A, Nabavi SM, Curti V, Daglia M. Exosome biogenesis, bioactivities and functions as new delivery systems of natural compounds. *Biotechnol Adv* 2018; **36**: 328-334 [PMID: 29248680 DOI: 10.1016/j.biotechadv.2017.12.010]

81 **Vu NB**, Nguyen HT, Palumbo R, Pellicano R, Fagoonee S, Pham PV. Stem cell-derived exosomes for wound healing: current status and promising directions. *Minerva Med* 2021; **112**: 384-400 [PMID: 33263376 DOI: 10.23736/S0026-4806.20.07205-5]

82 **Raposo G**, Stoorvogel W. Extracellular vesicles: exosomes, microvesicles, and friends. *J Cell Biol* 2013; **200**: 373-383 [PMID: 23420871 DOI: 10.1083/jcb.201211138]

83 **Lai P**, Weng J, Guo L, Chen X, Du X. Novel insights into MSC-EVs therapy for immune diseases. *Biomark Res* 2019; **7**: 6 [PMID: 30923617 DOI: 10.1186/s40364-019-0156-0]

84 **Börger V**, Bremer M, Ferrer-Tur R, Gockeln L, Stambouli O, Becic A, Giebel B. Mesenchymal Stem/Stromal Cell-Derived Extracellular Vesicles and Their Potential as Novel Immunomodulatory Therapeutic Agents. *Int J Mol Sci* 2017; **18** [PMID: 28684664 DOI: 10.3390/ijms18071450]

85 **Barile L**, Vassalli G. Exosomes: Therapy delivery tools and biomarkers of diseases. *Pharmacol Ther* 2017; **174**: 63-78 [PMID: 28202367 DOI: 10.1016/j.pharmthera.2017.02.020]

86 **Mianehsaz E**, Mirzaei HR, Mahjoubin-Tehran M, Rezaee A, Sahebhasagh R, Pourhanifeh MH, Mirzaei H, Hamblin MR. Mesenchymal stem cell-derived exosomes: a new therapeutic approach to osteoarthritis? *Stem Cell Res Ther* 2019; **10**: 340 [PMID: 31753036 DOI: 10.1186/s13287-019-1445-0]



- 87 **Zhang Y**, Bi J, Huang J, Tang Y, Du S, Li P. Exosome: A Review of Its Classification, Isolation Techniques, Storage, Diagnostic and Targeted Therapy Applications. *Int J Nanomedicine* 2020; **15**: 6917-6934 [PMID: 33061359 DOI: 10.2147/IJN.S264498]
- 88 **Jella KK**, Nasti TH, Li Z, Malla SR, Buchwald ZS, Khan MK. Exosomes, Their Biogenesis and Role in Inter-Cellular Communication, Tumor Microenvironment and Cancer Immunotherapy. *Vaccines (Basel)* 2018; **6** [PMID: 30261592 DOI: 10.3390/vaccines6040069]
- 89 **Zhang Y**, Liu Y, Liu H, Tang WH. Exosomes: biogenesis, biologic function and clinical potential. *Cell Biosci* 2019; **9**: 19 [PMID: 30815248 DOI: 10.1186/s13578-019-0282-2]
- 90 **Rai AK**, Johnson PJ. *Trichomonas vaginalis* extracellular vesicles are internalized by host cells using proteoglycans and caveolin-dependent endocytosis. *Proc Natl Acad Sci U S A* 2019; **116**: 21354-21360 [PMID: 31601738 DOI: 10.1073/pnas.1912356116]
- 91 **Costa Verdera H**, Gitz-Francois JJ, Schiffelers RM, Vader P. Cellular uptake of extracellular vesicles is mediated by clathrin-independent endocytosis and macropinocytosis. *J Control Release* 2017; **266**: 100-108 [PMID: 28919558 DOI: 10.1016/j.jconrel.2017.09.019]
- 92 **Isaac R**, Reis FCG, Ying W, Olefsky JM. Exosomes as mediators of intercellular crosstalk in metabolism. *Cell Metab* 2021; **33**: 1744-1762 [PMID: 34496230 DOI: 10.1016/j.cmet.2021.08.006]
- 93 **Zabeo D**, Cvjetkovic A, Lässer C, Schorb M, Lötval J, Höög JL. Exosomes purified from a single cell type have diverse morphology. *J Extracell Vesicles* 2017; **6**: 1329476 [PMID: 28717422 DOI: 10.1080/20013078.2017.1329476]
- 94 **Livshits MA**, Khomyakova E, Evtushenko EG, Lazarev VN, Kulemin NA, Semina SE, Generozov EV, Govorun VM. Isolation of exosomes by differential centrifugation: Theoretical analysis of a commonly used protocol. *Sci Rep* 2015; **5**: 17319 [PMID: 26616523 DOI: 10.1038/srep17319]

- 95 **Rider MA**, Hurwitz SN, Meckes DG Jr. ExtraPEG: A Polyethylene Glycol-Based Method for Enrichment of Extracellular Vesicles. *Sci Rep* 2016; **6**: 23978 [PMID: 27068479 DOI: 10.1038/srep23978]
- 96 **Böing AN**, van der Pol E, Grootemaat AE, Coumans FA, Sturk A, Nieuwland R. Single-step isolation of extracellular vesicles by size-exclusion chromatography. *J Extracell Vesicles* 2014; **3** [PMID: 25279113 DOI: 10.3402/jev.v3.23430]
- 97 **Vergauwen G**, Dhondt B, Van Deun J, De Smedt E, Berx G, Timmerman E, Gevaert K, Miinalainen I, Cocquyt V, Braems G, Van den Broecke R, Denys H, De Wever O, Hendrix A. Confounding factors of ultrafiltration and protein analysis in extracellular vesicle research. *Sci Rep* 2017; **7**: 2704 [PMID: 28577337 DOI: 10.1038/s41598-017-02599-y]
- 98 **Zarovni N**, Corrado A, Guazzi P, Zocco D, Lari E, Radano G, Muhhina J, Fondelli C, Gavrilova J, Chiesi A. Integrated isolation and quantitative analysis of exosome shuttled proteins and nucleic acids using immunocapture approaches. *Methods* 2015; **87**: 46-58 [PMID: 26044649 DOI: 10.1016/j.ymeth.2015.05.028]
- 99 **Liao HT**, Chen CT. Osteogenic potential: Comparison between bone marrow and adipose-derived mesenchymal stem cells. *World J Stem Cells* 2014; **6**: 288-295 [PMID: 25126378 DOI: 10.4252/wjsc.v6.i3.288]
- 100 **Álvarez-Viejo M**. Mesenchymal stem cells from different sources and their derived exosomes: A pre-clinical perspective. *World J Stem Cells* 2020; **12**: 100-109 [PMID: 32184935 DOI: 10.4252/wjsc.v12.i2.100]
- 101 **Wang Z**, Wu Y, Zhao Z, Liu C, Zhang L. Study on Transorgan Regulation of Intervertebral Disc and Extra-Skeletal Organs Through Exosomes Derived From Bone Marrow Mesenchymal Stem Cells. *Front Cell Dev Biol* 2021; **9**: 741183 [PMID: 34631718 DOI: 10.3389/fcell.2021.741183]
- 102 **Yaghoubi Y**, Movassaghpour A, Zamani M, Talebi M, Mehdizadeh A, Yousefi M. Human umbilical cord mesenchymal stem cells derived-exosomes in diseases treatment. *Life Sci* 2019; **233**: 116733 [PMID: 31394127 DOI: 10.1016/j.lfs.2019.116733]

- 103 **Schreml S**, Babilas P, Fruth S, Orsó E, Schmitz G, Mueller MB, Nerlich M, Prantl L. Harvesting human adipose tissue-derived adult stem cells: resection versus liposuction. *Cytotherapy* 2009; **11**: 947-957 [PMID: 19903106 DOI: 10.3109/14653240903204322]
- 104 **Pomatto M**, Gai C, Negro F, Cedrino M, Grange C, Ceccotti E, Togliatto G, Collino F, Tapparo M, Figliolini F, Lopatina T, Brizzi MF, Camussi G. Differential Therapeutic Effect of Extracellular Vesicles Derived by Bone Marrow and Adipose Mesenchymal Stem Cells on Wound Healing of Diabetic Ulcers and Correlation to Their Cargoes. *Int J Mol Sci* 2021; **22** [PMID: 33917759 DOI: 10.3390/ijms22083851]
- 105 **Shen Z**, Huang W, Liu J, Tian J, Wang S, Rui K. Effects of Mesenchymal Stem Cell-Derived Exosomes on Autoimmune Diseases. *Front Immunol* 2021; **12**: 749192 [PMID: 34646275 DOI: 10.3389/fimmu.2021.749192]
- 106 **Babaei M**, Rezaie J. Application of stem cell-derived exosomes in ischemic diseases: opportunity and limitations. *J Transl Med* 2021; **19**: 196 [PMID: 33964940 DOI: 10.1186/s12967-021-02863-w]
- 107 **Akbar N**, Azzimato V, Choudhury RP, Aouadi M. Extracellular vesicles in metabolic disease. *Diabetologia* 2019; **62**: 2179-2187 [PMID: 31690986 DOI: 10.1007/s00125-019-05014-5]
- 108 **Kalluri R**. The biology and function of exosomes in cancer. *J Clin Invest* 2016; **126**: 1208-1215 [PMID: 27035812 DOI: 10.1172/jci81135]
- 109 **Guo S**, Perets N, Betzer O, Ben-Shaul S, Sheinin A, Michaelievski I, Popovtzer R, Offen D, Levenberg S. Intranasal Delivery of Mesenchymal Stem Cell Derived Exosomes Loaded with Phosphatase and Tensin Homolog siRNA Repairs Complete Spinal Cord Injury. *ACS Nano* 2019; **13**: 10015-10028 [PMID: 31454225 DOI: 10.1021/acsnano.9b01892]
- 110 **Behera J**, Tyagi N. Exosomes: mediators of bone diseases, protection, and therapeutics potential. *Oncoscience* 2018; **5**: 181-195 [PMID: 30035185 DOI: 10.18632/oncoscience.421]
- 111 **Yu H**, Cheng J, Shi W, Ren B, Zhao F, Shi Y, Yang P, Duan X, Zhang J, Fu X, Hu X, Ao Y. Bone marrow mesenchymal stem cell-derived exosomes promote tendon

regeneration by facilitating the proliferation and migration of endogenous tendon stem/progenitor cells. *Acta Biomater* 2020; **106**: 328-341 [PMID: 32027991 DOI: 10.1016/j.actbio.2020.01.051]

112 **Guillamat-Prats R**. The Role of MSC in Wound Healing, Scarring and Regeneration. *Cells* 2021; **10** [PMID: 34359898 DOI: 10.3390/cells10071729]

113 **Askenase PW**. COVID-19 therapy with mesenchymal stromal cells (MSC) and convalescent plasma must consider exosome involvement: Do the exosomes in convalescent plasma antagonize the weak immune antibodies? *J Extracell Vesicles* 2020; **10**: e12004 [PMID: 33304473 DOI: 10.1002/jev2.12004]

114 **Nojehdehi S**, Soudi S, Hesampour A, Rasouli S, Soleimani M, Hashemi SM. Immunomodulatory effects of mesenchymal stem cell-derived exosomes on experimental type-1 autoimmune diabetes. *J Cell Biochem* 2018; **119**: 9433-9443 [PMID: 30074271 DOI: 10.1002/jcb.27260]

115 **Sun Y**, Shi H, Yin S, Ji C, Zhang X, Zhang B, Wu P, Shi Y, Mao F, Yan Y, Xu W, Qian H. Human Mesenchymal Stem Cell Derived Exosomes Alleviate Type 2 Diabetes Mellitus by Reversing Peripheral Insulin Resistance and Relieving  $\beta$ -Cell Destruction. *ACS Nano* 2018; **12**: 7613-7628 [PMID: 30052036 DOI: 10.1021/acsnano.7b07643]

116 **Biró E**, Sturk-Maquelin KN, Vogel GM, Meuleman DG, Smit MJ, Hack CE, Sturk A, Nieuwland R. Human cell-derived microparticles promote thrombus formation *in vivo* in a tissue factor-dependent manner. *J Thromb Haemost* 2003; **1**: 2561-2568 [PMID: 14738565 DOI: 10.1046/j.1538-7836.2003.00456.x]

117 **Del Conde I**, Shrimpton CN, Thiagarajan P, López JA. Tissue-factor-bearing microvesicles arise from lipid rafts and fuse with activated platelets to initiate coagulation. *Blood* 2005; **106**: 1604-1611 [PMID: 15741221 DOI: 10.1182/blood-2004-03-1095]

118 **Blazquez R**, Sanchez-Margallo FM, de la Rosa O, Dalemans W, Alvarez V, Tarazona R, Casado JG. Immunomodulatory Potential of Human Adipose Mesenchymal Stem Cells Derived Exosomes on *in vitro* Stimulated T Cells. *Front Immunol* 2014; **5**: 556 [PMID: 25414703 DOI: 10.3389/fimmu.2014.00556]

- 119 **Li X**, Xie X, Lian W, Shi R, Han S, Zhang H, Lu L, Li M. Exosomes from adipose-derived stem cells overexpressing Nrf2 accelerate cutaneous wound healing by promoting vascularization in a diabetic foot ulcer rat model. *Exp Mol Med* 2018; **50**: 1-14 [PMID: 29651102 DOI: 10.1038/s12276-018-0058-5]
- 120 **Wang J**, Xia J, Huang R, Hu Y, Fan J, Shu Q, Xu J. Mesenchymal stem cell-derived extracellular vesicles alter disease outcomes *via* endorsement of macrophage polarization. *Stem Cell Res Ther* 2020; **11**: 424 [PMID: 32993783 DOI: 10.1186/s13287-020-01937-8]
- 121 **Liu W**, Yu M, Xie D, Wang L, Ye C, Zhu Q, Liu F, Yang L. Melatonin-stimulated MSC-derived exosomes improve diabetic wound healing through regulating macrophage M1 and M2 polarization by targeting the PTEN/AKT pathway. *Stem Cell Res Ther* 2020; **11**: 259 [PMID: 32600435 DOI: 10.1186/s13287-020-01756-x]
- 122 **He X**, Dong Z, Cao Y, Wang H, Liu S, Liao L, Jin Y, Yuan L, Li B. MSC-Derived Exosome Promotes M2 Polarization and Enhances Cutaneous Wound Healing. *Stem Cells Int* 2019; **2019**: 7132708 [PMID: 31582986 DOI: 10.1155/2019/7132708]
- 123 **Zhao B**, Zhang X, Zhang Y, Lu Y, Zhang W, Lu S, Fu Y, Zhou Y, Zhang J, Zhang J. Human Exosomes Accelerate Cutaneous Wound Healing by Promoting Collagen Synthesis in a Diabetic Mouse Model. *Stem Cells Dev* 2021; **30**: 922-933 [PMID: 34167333 DOI: 10.1089/scd.2021.0100]
- 124 **Wang L**, Cai Y, Zhang Q, Zhang Y. Pharmaceutical Activation of Nrf2 Accelerates Diabetic Wound Healing by Exosomes from Bone Marrow Mesenchymal Stem Cells. *Int J Stem Cells* 2022; **15**: 164-172 [PMID: 34711700 DOI: 10.15283/ijsc21067]
- 125 **Dalirfardouei R**, Jamialahmadi K, Jafarian AH, Mahdipour E. Promising effects of exosomes isolated from menstrual blood-derived mesenchymal stem cell on wound-healing process in diabetic mouse model. *J Tissue Eng Regen Med* 2019; **13**: 555-568 [PMID: 30656863 DOI: 10.1002/term.2799]
- 126 **Ti D**, Hao H, Tong C, Liu J, Dong L, Zheng J, Zhao Y, Liu H, Fu X, Han W. LPS-preconditioned mesenchymal stromal cells modify macrophage polarization for



resolution of chronic inflammation *via* exosome-shuttled let-7b. *J Transl Med* 2015; **13**: 308 [PMID: 26386558 DOI: 10.1186/s12967-015-0642-6]

127 **Sun B**, Wu F, Wang X, Song Q, Ye Z, Mohammadniaei M, Zhang M, Chu X, Xi S, Zhou N, Wang W, Yao C, Shen J. An Optimally Designed Engineering Exosome-Reductive COF Integrated Nanoagent for Synergistically Enhanced Diabetic Fester Wound Healing. *Small* 2022; **18**: e2200895 [PMID: 35638464 DOI: 10.1002/sml.202200895]

128 **Geng X**, Qi Y, Liu X, Shi Y, Li H, Zhao L. A multifunctional antibacterial and self-healing hydrogel laden with bone marrow mesenchymal stem cell-derived exosomes for accelerating diabetic wound healing. *Biomater Adv* 2022; **133**: 112613 [PMID: 35527135 DOI: 10.1016/j.msec.2021.112613]

129 **Zhang Y**, Han F, Gu L, Ji P, Yang X, Liu M, Tao K, Hu D. Adipose mesenchymal stem cell exosomes promote wound healing through accelerated keratinocyte migration and proliferation by activating the AKT/HIF-1 $\alpha$  axis. *J Mol Histol* 2020; **51**: 375-383 [PMID: 32556903 DOI: 10.1007/s10735-020-09887-4]

130 **Ma T**, Fu B, Yang X, Xiao Y, Pan M. Adipose mesenchymal stem cell-derived exosomes promote cell proliferation, migration, and inhibit cell apoptosis *via* Wnt/ $\beta$ -catenin signaling in cutaneous wound healing. *J Cell Biochem* 2019; **120**: 10847-10854 [PMID: 30681184 DOI: 10.1002/jcb.28376]

131 **Shabbir A**, Cox A, Rodriguez-Menocal L, Salgado M, Van Badiavas E. Mesenchymal Stem Cell Exosomes Induce Proliferation and Migration of Normal and Chronic Wound Fibroblasts, and Enhance Angiogenesis In Vitro. *Stem Cells Dev* 2015; **24**: 1635-1647 [PMID: 25867197 DOI: 10.1089/scd.2014.0316]

132 **Liang X**, Zhang L, Wang S, Han Q, Zhao RC. Exosomes secreted by mesenchymal stem cells promote endothelial cell angiogenesis by transferring miR-125a. *J Cell Sci* 2016; **129**: 2182-2189 [PMID: 27252357 DOI: 10.1242/jcs.170373]

133 **Zhao G**, Liu F, Liu Z, Zuo K, Wang B, Zhang Y, Han X, Lian A, Wang Y, Liu M, Zou F, Li P, Liu X, Jin M, Liu JY. MSC-derived exosomes attenuate cell death through

suppressing AIF nucleus translocation and enhance cutaneous wound healing. *Stem Cell Res Ther* 2020; **11**: 174 [PMID: 32393338 DOI: 10.1186/s13287-020-01616-8]

134 **Wang X**, Jiao Y, Pan Y, Zhang L, Gong H, Qi Y, Wang M, Gong H, Shao M, Wang X, Jiang D. Fetal Dermal Mesenchymal Stem Cell-Derived Exosomes Accelerate Cutaneous Wound Healing by Activating Notch Signaling. *Stem Cells Int* 2019; **2019**: 2402916 [PMID: 31281370 DOI: 10.1155/2019/2402916]

135 **Bian X**, Li B, Yang J, Ma K, Sun M, Zhang C, Fu X. Regenerative and protective effects of dMSC-sEVs on high-glucose-induced senescent fibroblasts by suppressing RAGE pathway and activating Smad pathway. *Stem Cell Res Ther* 2020; **11**: 166 [PMID: 32349787 DOI: 10.1186/s13287-020-01681-z]

136 **Li B**, Luan S, Chen J, Zhou Y, Wang T, Li Z, Fu Y, Zhai A, Bi C. The MSC-Derived Exosomal lncRNA H19 Promotes Wound Healing in Diabetic Foot Ulcers by Upregulating PTEN via MicroRNA-152-3p. *Mol Ther Nucleic Acids* 2020; **19**: 814-826 [PMID: 31958697 DOI: 10.1016/j.omtn.2019.11.034]

137 **Yu M**, Liu W, Li J, Lu J, Lu H, Jia W, Liu F. Exosomes derived from atorvastatin-pretreated MSC accelerate diabetic wound repair by enhancing angiogenesis via AKT/eNOS pathway. *Stem Cell Res Ther* 2020; **11**: 350 [PMID: 32787917 DOI: 10.1186/s13287-020-01824-2]

138 **Hu Y**, Tao R, Chen L, Xiong Y, Xue H, Hu L, Yan C, Xie X, Lin Z, Panayi AC, Mi B, Liu G. Exosomes derived from pioglitazone-pretreated MSCs accelerate diabetic wound healing through enhancing angiogenesis. *J Nanobiotechnology* 2021; **19**: 150 [PMID: 34020670 DOI: 10.1186/s12951-021-00894-5]

139 **Shi R**, Jin Y, Hu W, Lian W, Cao C, Han S, Zhao S, Yuan H, Yang X, Shi J, Zhao H. Exosomes derived from mmu\_circ\_0000250-modified adipose-derived mesenchymal stem cells promote wound healing in diabetic mice by inducing miR-128-3p/SIRT1-mediated autophagy. *Am J Physiol Cell Physiol* 2020; **318**: C848-C856 [PMID: 32159361 DOI: 10.1152/ajpcell.00041.2020]

140 **Yang J**, Chen Z, Pan D, Li H, Shen J. Umbilical Cord-Derived Mesenchymal Stem Cell-Derived Exosomes Combined Pluronic F127 Hydrogel Promote Chronic Diabetic

Wound Healing and Complete Skin Regeneration. *Int J Nanomedicine* 2020; **15**: 5911-5926 [PMID: 32848396 DOI: 10.2147/IJN.S249129]

141 **Wang C**, Wang M, Xu T, Zhang X, Lin C, Gao W, Xu H, Lei B, Mao C. Engineering Bioactive Self-Healing Antibacterial Exosomes Hydrogel for Promoting Chronic Diabetic Wound Healing and Complete Skin Regeneration. *Theranostics* 2019; **9**: 65-76 [PMID: 30662554 DOI: 10.7150/thno.29766]

142 **Shi Q**, Qian Z, Liu D, Sun J, Wang X, Liu H, Xu J, Guo X. GMSC-Derived Exosomes Combined with a Chitosan/Silk Hydrogel Sponge Accelerates Wound Healing in a Diabetic Rat Skin Defect Model. *Front Physiol* 2017; **8**: 904 [PMID: 29163228 DOI: 10.3389/fphys.2017.00904]

143 **Jiang T**, Wang Z, Sun J. Human bone marrow mesenchymal stem cell-derived exosomes stimulate cutaneous wound healing mediates through TGF- $\beta$ /Smad signaling pathway. *Stem Cell Res Ther* 2020; **11**: 198 [PMID: 32448395 DOI: 10.1186/s13287-020-01723-6]

144 **Fang S**, Xu C, Zhang Y, Xue C, Yang C, Bi H, Qian X, Wu M, Ji K, Zhao Y, Wang Y, Liu H, Xing X. Umbilical Cord-Derived Mesenchymal Stem Cell-Derived Exosomal MicroRNAs Suppress Myofibroblast Differentiation by Inhibiting the Transforming Growth Factor- $\beta$ /SMAD2 Pathway During Wound Healing. *Stem Cells Transl Med* 2016; **5**: 1425-1439 [PMID: 27388239 DOI: 10.5966/sctm.2015-0367]

145 **Zhang B**, Shi Y, Gong A, Pan Z, Shi H, Yang H, Fu H, Yan Y, Zhang X, Wang M, Zhu W, Qian H, Xu W. HucMSC Exosome-Delivered 14-3-3 $\zeta$  Orchestrates Self-Control of the Wnt Response *via* Modulation of YAP During Cutaneous Regeneration. *Stem Cells* 2016; **34**: 2485-2500 [PMID: 27334574 DOI: 10.1002/stem.2432]

146 **Wang L**, Hu L, Zhou X, Xiong Z, Zhang C, Shehada HMA, Hu B, Song J, Chen L. Exosomes secreted by human adipose mesenchymal stem cells promote scarless cutaneous repair by regulating extracellular matrix remodelling. *Sci Rep* 2017; **7**: 13321 [PMID: 29042658 DOI: 10.1038/s41598-017-12919-x]

147 **Li M**, Ke QF, Tao SC, Guo SC, Rui BY, Guo YP. Fabrication of hydroxyapatite/chitosan composite hydrogels loaded with exosomes derived from

miR-126-3p overexpressed synovial mesenchymal stem cells for diabetic chronic wound healing. *J Mater Chem B* 2016; **4**: 6830-6841 [PMID: 32263577 DOI: 10.1039/c6tb01560c]

148 **Zhang Y**, Zhang P, Gao X, Chang L, Chen Z, Mei X. Preparation of exosomes encapsulated nanohydrogel for accelerating wound healing of diabetic rats by promoting angiogenesis. *Mater Sci Eng C Mater Biol Appl* 2021; **120**: 111671 [PMID: 33545836 DOI: 10.1016/j.msec.2020.111671]

149 **Hu L**, Wang J, Zhou X, Xiong Z, Zhao J, Yu R, Huang F, Zhang H, Chen L. Exosomes derived from human adipose mesenchymal stem cells accelerates cutaneous wound healing *via* optimizing the characteristics of fibroblasts. *Sci Rep* 2016; **6**: 32993 [PMID: 27615560 DOI: 10.1038/srep32993]

150 **Tao SC**, Guo SC, Li M, Ke QF, Guo YP, Zhang CQ. Chitosan Wound Dressings Incorporating Exosomes Derived from MicroRNA-126-Overexpressing Synovium Mesenchymal Stem Cells Provide Sustained Release of Exosomes and Heal Full-Thickness Skin Defects in a Diabetic Rat Model. *Stem Cells Transl Med* 2017; **6**: 736-747 [PMID: 28297576 DOI: 10.5966/sctm.2016-0275]

151 **Xiao S**, Xiao C, Miao Y, Wang J, Chen R, Fan Z, Hu Z. Human acellular amniotic membrane incorporating exosomes from adipose-derived mesenchymal stem cells promotes diabetic wound healing. *Stem Cell Res Ther* 2021; **12**: 255 [PMID: 33926555 DOI: 10.1186/s13287-021-02333-6]

152 **Yan C**, Xv Y, Lin Z, Endo Y, Xue H, Hu Y, Hu L, Chen L, Cao F, Zhou W, Zhang P, Liu G. Human Umbilical Cord Mesenchymal Stem Cell-Derived Exosomes Accelerate Diabetic Wound Healing *via* Ameliorating Oxidative Stress and Promoting Angiogenesis. *Front Bioeng Biotechnol* 2022; **10**: 829868 [PMID: 35174145 DOI: 10.3389/fbioe.2022.829868]

153 **Gondaliya P**, Sayyed AA, Bhat P, Mali M, Arya N, Khairnar A, Kalia K. Mesenchymal Stem Cell-Derived Exosomes Loaded with miR-155 Inhibitor Ameliorate Diabetic Wound Healing. *Mol Pharm* 2022; **19**: 1294-1308 [PMID: 35294195 DOI: 10.1021/acs.molpharmaceut.1c00669]

- 154 **Han ZF**, Cao JH, Liu ZY, Yang Z, Qi RX, Xu HL. Exosomal lncRNA KLF3-AS1 derived from bone marrow mesenchymal stem cells stimulates angiogenesis to promote diabetic cutaneous wound healing. *Diabetes Res Clin Pract* 2022; **183**: 109126 [PMID: 34742784 DOI: 10.1016/j.diabres.2021.109126]
- 155 **Ding J**, Wang X, Chen B, Zhang J, Xu J. Exosomes Derived from Human Bone Marrow Mesenchymal Stem Cells Stimulated by Deferoxamine Accelerate Cutaneous Wound Healing by Promoting Angiogenesis. *Biomed Res Int* 2019; **2019**: 9742765 [PMID: 31192260 DOI: 10.1155/2019/9742765]
- 156 **Zhang Y**, Bai X, Shen K, Luo L, Zhao M, Xu C, Jia Y, Xiao D, Li Y, Gao X, Tian C, Wang Y, Hu D. Exosomes Derived from Adipose Mesenchymal Stem Cells Promote Diabetic Chronic Wound Healing through SIRT3/SOD2. *Cells* 2022; **11** [PMID: 36010644 DOI: 10.3390/cells11162568]
- 157 **Born LJ**, Chang KH, Shoureshi P, Lay F, Bengali S, Hsu ATW, Abadchi SN, Harmon JW, Jay SM. HOTAIR-Loaded Mesenchymal Stem/Stromal Cell Extracellular Vesicles Enhance Angiogenesis and Wound Healing. *Adv Healthc Mater* 2022; **11**: e2002070 [PMID: 33870645 DOI: 10.1002/adhm.202002070]
- 158 **Teng L**, Maqsood M, Zhu M, Zhou Y, Kang M, Zhou J, Chen J. Exosomes Derived from Human Umbilical Cord Mesenchymal Stem Cells Accelerate Diabetic Wound Healing *via* Promoting M2 Macrophage Polarization, Angiogenesis, and Collagen Deposition. *Int J Mol Sci* 2022; **23** [PMID: 36142334 DOI: 10.3390/ijms231810421]
- 159 **Kleinert M**, Clemmensen C, Hofmann SM, Moore MC, Renner S, Woods SC, Huypens P, Beckers J, de Angelis MH, Schürmann A, Bakhti M, Klingenspor M, Heiman M, Cherrington AD, Ristow M, Lickert H, Wolf E, Havel PJ, Müller TD, Tschöp MH. Animal models of obesity and diabetes mellitus. *Nat Rev Endocrinol* 2018; **14**: 140-162 [PMID: 29348476 DOI: 10.1038/nrendo.2017.161]
- 160 **Li Y**, Liang Y, Gao Y, Chen D, Ran X. Dynamic changes of wound-related miRNAs after application of autologous platelet-rich gel in diabetic wounds. *Chin Med J* 2022; Epub ahead of print [DOI: 10.26226/morressier.5d9b6232ea541d6ca84940ad]



161 **Ha DH**, Kim SD, Lee J, Kwon HH, Park GH, Yang SH, Jung JY, Lee JH, Park SR, Youn J, Lee SH, Kim JE, Lim J, Lee HK, Cho BS, Yi YW. Toxicological evaluation of exosomes derived from human adipose tissue-derived mesenchymal stem/stromal cells. *Regul Toxicol Pharmacol* 2020; **115**: 104686 [PMID: 32450131 DOI: 10.1016/j.yrtph.2020.104686]

162 **Varderidou-Minasian S**, Lorenowicz MJ. Mesenchymal stromal/stem cell-derived extracellular vesicles in tissue repair: challenges and opportunities. *Theranostics* 2020; **10**: 5979-5997 [PMID: 32483432 DOI: 10.7150/thno.40122]

163 **Philipp D**, Suhr L, Wahlers T, Choi YH, Paunel-Görgülü A. Preconditioning of bone marrow-derived mesenchymal stem cells highly strengthens their potential to promote IL-6-dependent M2b polarization. *Stem Cell Res Ther* 2018; **9**: 286 [PMID: 30359316 DOI: 10.1186/s13287-018-1039-2]

17%

SIMILARITY INDEX

---

PRIMARY SOURCES

---

- |   |   |                      |
|---|---|----------------------|
| <div style="background-color: red; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">1</div>    | <p>Yi Zhang, Jiayao Bi, Jiayi Huang, Yanan Tang, Shouying Du, Pengyue Li. "&lt;p&gt;Exosome: A Review of Its Classification, Isolation Techniques, Storage, Diagnostic and Targeted Therapy Applications&lt;/p&gt;", International Journal of Nanomedicine, 2020</p> <p><small>Crossref</small></p> | <p>61 words — 1%</p> |
| <hr/>   |   |                      |
| <div style="background-color: purple; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">2</div> | <p><a href="http://www.frontiersin.org">www.frontiersin.org</a></p> <p><small>Internet</small></p>  | <p>49 words — 1%</p> |
| <hr/>   |   |                      |
| <div style="background-color: purple; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">3</div> | <p>Uyen Thi Trang Than, Dominic Guanzon, David Leavesley, Tony Parker. "Association of Extracellular Membrane Vesicles with Cutaneous Wound Healing", International Journal of Molecular Sciences, 2017</p> <p><small>Crossref</small></p>  | <p>39 words — 1%</p> |
| <hr/>   |   |                      |
| <div style="background-color: teal; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">4</div>   | <p><a href="http://www.mdpi.com">www.mdpi.com</a></p> <p><small>Internet</small></p>  | <p>39 words — 1%</p> |
| <hr/>   |   |                      |
| <div style="background-color: green; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">5</div>  | <p>Nina Kosaric, Harriet Kiwanuka, Geoffrey C Gurtner. "Stem cell therapies for wound healing", Expert Opinion on Biological Therapy, 2019</p> <p><small>Crossref</small></p>   | <p>34 words — 1%</p> |
| <hr/>   |   |                      |
| <div style="background-color: brown; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin-bottom: 10px;">6</div>  | <p>Holly N. Wilkinson, Matthew J. Hardman. "Wound healing: cellular mechanisms and pathological outcomes", Open Biology, 2020</p> <p><small>Crossref</small></p>  | <p>32 words — 1%</p> |

|    |  |                 |
|----|--|-----------------|
| 7  | link.springer.com<br>Internet  | 30 words — < 1% |
| 8  | stemcellres.biomedcentral.com<br>Internet  | 28 words — < 1% |
| 9  | Roi Isaac, Felipe Castellani Gomes Reis, Wei Ying, Jerrold M. Olefsky. "Exosomes as mediators of intercellular crosstalk in metabolism", Cell Metabolism, 2021<br>Crossref   | 24 words — < 1% |
| 10 | repository.unair.ac.id<br>Internet   | 23 words — < 1% |
| 11 | dokumen.pub<br>Internet  | 22 words — < 1% |
| 12 | Lu Wang, Li Hu, Xin Zhou, Zehuan Xiong, Chenguang Zhang, Hassan M. A. Shehada, Bo Hu, Jinlin Song, Lili Chen. "Exosomes secreted by human adipose mesenchymal stem cells promote scarless cutaneous repair by regulating extracellular matrix remodelling", Scientific Reports, 2017<br>Crossref | 21 words — < 1% |
| 13 | Ziwei Shen, Wei Huang, Jun Liu, Jie Tian, Shengjun Wang, Ke Rui. "Effects of Mesenchymal Stem Cell-Derived Exosomes on Autoimmune Diseases", Frontiers in Immunology, 2021<br>Crossref   | 20 words — < 1% |
| 14 | www.wjgnet.com<br>Internet   | 19 words — < 1% |
| 15 | Ali Golchin, Simzar Hosseinzadeh, Abdolreza Ardeshirylajimi. "The exosomes released from   | 18 words — < 1% |

- 16 Zhao, Shichang, Le Li, Hui Wang, Yadong Zhang, Xiangguo Cheng, Nai Zhou, Mohamed N. Rahaman, Zhongtang Liu, Wenhai Huang, and Changqing Zhang. "Wound dressings composed of copper-doped borate bioactive glass microfibers stimulate angiogenesis and heal full-thickness skin defects in a rodent model", Biomaterials, 2015. 16 words — < 1%

- 17 Chuanjiang He, Shu Zheng, Yan Luo, Ben Wang. "Exosome Theranostics: Biology and Translational Medicine", Theranostics, 2018 15 words — < 1%

- 18 Jana Janockova, Lucia Slovinska, Denisa Harvanova, Timea Spakova, Jan Rosocha. "New therapeutic approaches of mesenchymal stem cells-derived exosomes", Journal of Biomedical Science, 2021 15 words — < 1%

- 19 Joo, Suh, Lee, Bang, Lee. "Current Knowledge and Future Perspectives on Mesenchymal Stem Cell-Derived Exosomes as a New Therapeutic Agent", International Journal of Molecular Sciences, 2020 15 words — < 1%

- 20 "Local Wound Care for Dermatologists", Springer Science and Business Media LLC, 2020 14 words — < 1%

- 21 "Translational Stem Cell Research", Springer Science and Business Media LLC, 2011 14 words — < 1%

22 Chenggui Wang, Min Wang, Tianzhen Xu, Xingxing Zhang, Cai Lin, Weiyang Gao, Huazi Xu, Bo Lei, Cong Mao. "Engineering Bioactive Self-Healing Antibacterial Exosomes Hydrogel for Promoting Chronic Diabetic Wound Healing and Complete Skin Regeneration", *Theranostics*, 2019

14 words — < 1%

Crossref

23 Zhou Xunian, Raghu Kalluri. "Biology and therapeutic potential of mesenchymal stem cell - derived exosomes", *Cancer Science*, 2020

14 words — < 1%

Crossref

24 Yan Chang, Xinwei Wang, Susu Li, Yueye Wang, Chengyan Jia, Xuezhi Yang, Siyu Li, Bingjie Zhang, Yingjie Zhao, Wei Wei. "Activation of KYN-AHR axis impairs the chondrogenic and chondroprotective effects of human umbilical cord-derived mesenchymal stem cells on osteoarthritis", *Research Square Platform LLC*, 2022

13 words — < 1%

Crossref Posted Content

25 Dongdong Ti, Haojie Hao, Chuan Tong, Jiejie Liu, Liang Dong, Jingxi Zheng, Yali Zhao, Huiling Liu, Xiaobing Fu, Weidong Han. "LPS-preconditioned mesenchymal stromal cells modify macrophage polarization for resolution of chronic inflammation via exosome-shuttled let-7b", *Journal of Translational Medicine*, 2015

12 words — < 1%

Crossref

26 Lan Li, Dawei Chen, Chun Wang, Guanjian Liu, Xingwu Ran. "The Effect of Autologous Platelet-Rich Gel on the Dynamic Changes of the Matrix Metalloproteinase-2 and Tissue Inhibitor of Metalloproteinase-2 Expression in the Diabetic Chronic Refractory Cutaneous Ulcers", *Journal of Diabetes Research*, 2015

12 words — < 1%

Crossref



---

28 "Cell Biology and Translational Medicine, Volume 11", Springer Science and Business Media LLC, 2021

Crossref

---

29 "Exosomes in Cardiovascular Diseases", Springer Science and Business Media LLC, 2017

Crossref

---

30 András Mester, Diana Opincariu, Imre Benedek, István Benedek. "Stem Cell Therapy in Wound Healing", Journal of Interdisciplinary Medicine, 2017

Crossref

---

31 Jayashree Vijaya Raghavan, Vinod Kumar Dorai, Shruthi Ksheera Sagar, Archana Sivaraman, Kalpana S R, Siddharth Jhunjhunwala. "Immunomodulatory Bandage for Accelerated Healing of Diabetic Wounds", ACS Bio & Med Chem Au, 2022

Crossref

---

32 José Luis Lázaro-Martínez, Yolanda García-Álvarez, Francisco Javier Álvaro-Afonso, Esther García-Morales et al. "Hard-to-heal diabetic foot ulcers treated using negatively charged polystyrene microspheres: a prospective case series", Journal of Wound Care, 2019

Crossref

---

33 Razieh Dalirfardoue, Khadijeh Jamialahmadi, Amir Hossein Jafarian, Elahe Mahdipour. "Promising effects of exosomes isolated from menstrual blood - derived mesenchymal stem cell on wound - healing process in diabetic mouse model", Journal of Tissue Engineering and Regenerative Medicine, 2019

Crossref

34 Shichang Zhang, Li Chen, Guoying Zhang, Bo Zhang. "Umbilical cord-matrix stem cells induce the functional restoration of vascular endothelial cells and enhance skin wound healing in diabetic mice via the polarized macrophages", Stem Cell Research & Therapy, 2020

11 words — < 1%

Crossref

35 Zhi-Xiang Tan, Rui Tao, Si-Cheng Li, Bing-Zheng Shen, Lan-Xia Meng, Zhan-Yong Zhu. "Role of defensins in diabetic wound healing", World Journal of Diabetes, 2022

11 words — < 1%

Crossref

36 Ziyin Pan, Weiyan Sun, Yi Chen, Hai Tang, Weikang Lin, Jiafei Chen, Chang Chen. "Extracellular Vesicles in Tissue Engineering: Biology and Engineered Strategy", Advanced Healthcare Materials, 2022

11 words — < 1%

Crossref

37 eurjmedres.biomedcentral.com

Internet

11 words — < 1%

38 f6publishing.blob.core.windows.net

Internet

11 words — < 1%

39 www.cjter.com

Internet

11 words — < 1%

40 "Ganoderma and Health", Springer Science and Business Media LLC, 2019

10 words — < 1%

Crossref

41 Gurvinder Kaur, Ganesh Narayanan, Deepa Garg, Abhay Sachdev, Ishita Matai. "Biomaterials-Based Regenerative Strategies for Skin Tissue Wound Healing", ACS Applied Bio Materials, 2022

10 words — < 1%

Crossref

42 He Qiu, Shuo Liu, Kelun Wu, Rui Zhao, Lideng Cao, Hang Wang. "Prospective application of exosomes derived from adipose - derived stem cells in skin wound healing: A review", Journal of Cosmetic Dermatology, 2019

10 words — < 1%

[Crossref](#)

43 Jianrui Li, Chuanyuan Wei, Yang Yang, Zixu Gao, Zheng Guo, Fazhi Qi. "Apoptotic bodies extracted from adipose mesenchymal stem cells carry microRNA-21-5p to induce M2 polarization of macrophages and augment skin wound healing by targeting KLF6", Burns, 2022

10 words — < 1%

[Crossref](#)

44 Jin Cheng, Yixin Sun, Yong Ma, Yingfang Ao, Xiaoqing Hu, Qingyang Meng. "Engineering of MSC-Derived Exosomes: A Promising Cell-Free Therapy for Osteoarthritis", Membranes, 2022

10 words — < 1%

[Crossref](#)

45 Joana Cabral, Aideen E. Ryan, Matthew D. Griffin, Thomas Ritter. "Extracellular vesicles as modulators of wound healing", Advanced Drug Delivery Reviews, 2018

10 words — < 1%

[Crossref](#)

46 Na Li, Jinlian Hua. "Interactions between mesenchymal stem cells and the immune system", Cellular and Molecular Life Sciences, 2017

10 words — < 1%

[Crossref](#)

47 Qinan Wu, Bing Chen, Ziwen Liang. "Mesenchymal Stem Cells as a Prospective Therapy for the Diabetic Foot", Stem Cells International, 2016

10 words — < 1%

[Crossref](#)

48 Shihuan Cheng, Zhiyu Xi, Guang Chen, Kai Liu, Renshi Ma, Chen Zhou. "Extracellular vesicle-carried microRNA-27b derived from mesenchymal stem cells accelerates cutaneous wound healing via E3 ubiquitin ligase ITCH", Journal of Cellular and Molecular Medicine, 2020

10 words — < 1%

Crossref

49 [rimi.imi.bg.ac.rs](http://rimi.imi.bg.ac.rs)

Internet

10 words — < 1%

50 "Chronic Wounds, Wound Dressings and Wound Healing", Springer Science and Business Media LLC, 2021

9 words — < 1%

Crossref

51 "Exosomes, Stem Cells and MicroRNA", Springer Science and Business Media LLC, 2018

9 words — < 1%

Crossref

52 Bo Li, Song Luan, Jing Chen, Yue Zhou, Tingting Wang, Zhijuan Li, Yili Fu, Aixia Zhai, Changlong Bi. "The MSC-Derived Exosomal lncRNA H19 Promotes Wound Healing in Diabetic Foot Ulcers by Upregulating PTEN via MicroRNA-152-3p", Molecular Therapy - Nucleic Acids, 2020

9 words — < 1%

Crossref

53 Bo Zhao, Xingliao Zhang, Yuanlin Zhang, Yijun Lu, Wanting Zhang, Shoutao Lu, Yu Fu, Yang Zhou, Jun Zhang, Jing Zhang. "Human Exosomes Accelerate Cutaneous Wound Healing by Promoting Collagen Synthesis in a Diabetic Mouse Model", Stem Cells and Development, 2021

9 words — < 1%

Crossref

54 Douglas D. Taylor, Cicek Gercel-Taylor. "Exosomes/microvesicles: mediators of cancer-associated immunosuppressive microenvironments", Seminars in Immunopathology, 2011

9 words — < 1%

55 Endong Zhang, Philana Phan, Zongmin Zhao. "Cellular nanovesicles for therapeutic immunomodulation: A perspective on engineering strategies and new advances", *Acta Pharmaceutica Sinica B*, 2022

9 words — < 1%

Crossref

56 Hongyan Deng, Yong Chen. "The role of adipose-derived stem cells-derived extracellular vesicles in the treatment of diabetic foot ulcer: Trends and prospects", *Frontiers in Endocrinology*, 2022

9 words — < 1%

Crossref

57 Hui Nie, Zhujun Liao, Yutong Wang, Jianhua Zhou, Xiaoyun He, Chunlin Ou. "Exosomal long non-coding RNAs: Emerging players in cancer metastasis and potential diagnostic biomarkers for personalized oncology", *Genes & Diseases*, 2020

9 words — < 1%

Crossref

58 Jianying Song, Lixin Hu, Bo Liu, Nan Jiang et al. "The Emerging Role of Immune Cells and Targeted Therapeutic Strategies in Diabetic Wounds Healing", *Journal of Inflammation Research*, 2022

9 words — < 1%

Crossref

59 Jinlong Shao, Yating Cui, Ye Liang, Hong Liu, Baojin Ma, Shaohua Ge. "Unilateral Silver-Loaded Silk Fibroin Difunctional Membranes as Antibacterial Wound Dressings", *ACS Omega*, 2021

9 words — < 1%

Crossref

60 Juan Cen, Yichen Zhang, Yindu Bai, Shenqian Ma, Chuan Zhang, Lin Jin, Shaofeng Duan, Yanan Du, Yuqi Guo. "Research progress of stem cell therapy for endometrial injury", *Materials Today Bio*, 2022

9 words — < 1%

Crossref

---

61 MeiHua Wan, Bo Ning, Sarah Spiegel, Christopher J. Lyon, Tony Y. Hu. "Tumor - derived exosomes (TDEs): How to avoid the sting in the tail", Medicinal Research Reviews, 2019

9 words — < 1%

Crossref

---

62 Meng Kou, Li Huang, Jinjuan Yang, Zhixin Chiang et al. "Mesenchymal stem cell-derived extracellular vesicles for immunomodulation and regeneration: a next generation therapeutic tool?", Cell Death & Disease, 2022

9 words — < 1%

Crossref

---

63 Ping-Ruey Chou, Yun-Nan Lin, Sheng-Hua Wu, Sin-Daw Lin, Periasamy Srinivasan, Dar-Jen Hsieh, Shu-Hung Huang. "Supercritical Carbon Dioxide-decellularized Porcine Acellular Dermal Matrix combined with Autologous Adipose-derived Stem Cells: Its Role in Accelerated Diabetic Wound Healing", International Journal of Medical Sciences, 2020

9 words — < 1%

Crossref

---

64 Raquel Guillamat-Prats. "The Role of MSC in Wound Healing, Scarring and Regeneration", Cells, 2021

9 words — < 1%

Crossref

---

65 Shuo Wang, Biyu Lei, E Zhang, Puyang Gong, Jian Gu, Lili He, Lu Han, Zhixiang Yuan. "Targeted Therapy for Inflammatory Diseases with Mesenchymal Stem Cells and Their Derived Exosomes: From Basic to Clinics", International Journal of Nanomedicine, 2022

9 words — < 1%

Crossref

---

66 Wei Dai, Yuchen Dong, Ting Han, Jing Wang, Bin Gao, Hui Guo, Feng Xu, Jing Li, Yufei Ma. "Microenvironmental cue-regulated exosomes as therapeutic

9 words — < 1%



67 Yiqiang Hu, Bin Wu, Yuan Xiong, Ranyang Tao et al. "Cryogenic 3D printed hydrogel scaffolds loading exosomes accelerate diabetic wound healing", Chemical Engineering Journal, 2021

Crossref

68 [accounts.public.ce.basespace.illumina.com](https://accounts.public.ce.basespace.illumina.com)

Internet

69 [camacs.ca](https://camacs.ca)

Internet

70 [journals.lww.com](https://journals.lww.com)

Internet

71 [journals.plos.org](https://journals.plos.org)

Internet

72 [www.ncbi.nlm.nih.gov](https://www.ncbi.nlm.nih.gov)

Internet

73 [www.researchgate.net](https://www.researchgate.net)

Internet

EXCLUDE QUOTES ON

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES < 9 WORDS

EXCLUDE MATCHES < 9 WORDS