

79357_Auto_Edited.docx

Name of Journal: *World Journal of Diabetes*

Manuscript NO: 79357

Manuscript Type: ORIGINAL ARTICLE

Basic Study

miR-124 is upregulated in diabetic mice and inhibits proliferation and promotes apoptosis of high-glucose-induced β -cells by targeting EZH2

miR-124 and EZH2 in diabetes

Duan Xiao-Kai, Sun Yong-xiang, Wang Hong-yun, Xu Yan-yan, Fan Shi-zhen, Tian Jin-ya, Yu Yong, Zhao Yan-yun, Jiang Yan-li

Abstract

BACKGROUND

Diabetes is a chronic metabolic disease, and a variety of miRNA are involved in the occurrence and development of diabetes. In clinical studies, miR-124 is highly expressed in the serum of patients with diabetes and in pancreatic islet β -cells. However, few reports exist concerning the role and mechanism of action of miR-124 in diabetes.

AIM

To investigate the expression of miR-124 in diabetic mice and the potential mechanism of action in islet β -cells.

METHODS

The expression levels of miR-124 and enhancer of zeste homolog 2 (EZH2) in pancreatic tissues of diabetic mice were detected. The targeted relationship between miR-124 and EZH2 was predicted by Targetscan software and verified by a double luciferase reporter assay. Mouse islet β -cells Min6 were grown in a high glucose (HG) medium to mimic a diabetes model. The insulin secretion, proliferation, cell cycle and apoptosis of HG-induced Min6 cells were detected after interference of miR-124a and/or EZH2.

RESULTS

The expression of miR-124 was upregulated and EZH2 was downregulated in the pancreatic tissue of diabetic mice compared with control mice, and the expression of miR-124 was negatively correlated with that of EZH2. miR-124 was highly expressed in HG-induced Min6 cells. Inhibition of miR-124 promoted insulin secretion and cell proliferation, induced the transition from the G0/G1 phase to the S phase of the cell cycle, and inhibited cell apoptosis in HG-induced Min6 cells. EZH2 was one of the targets of miR-124. Co-transfection of miR-124 inhibitor and siRNA-EZH2 could reverse the effects of the miR-124 inhibitor in HG-induced Min6 cells.

CONCLUSION

miR-124 is highly expressed in diabetic mice and HG-induced Min6 cells and regulates insulin secretion, proliferation and apoptosis of islet β -cells by targeting EZH2

Key Words: miR-124; EZH2; Diabetes; Insulin secretion; Proliferation; Apoptosis

Xiao-Kai D, Yong-xiang S, Hong-yun W, Yan-yan X, Shi-zhen F, Jin-ya T, Yong Y, Yan-yun Z, Yan-li J. miR-124 is upregulated in diabetic mice and inhibits proliferation and promotes apoptosis of high-glucose-induced β -cells by targeting EZH2. *World J Diabetes* 2023; In press

Core Tip: In clinical studies, miR-124 is highly expressed in the serum of patients with diabetes and in pancreatic islet β -cells. In this study, the role and mechanism of miR-124 in diabetes was explored. miR-124 was highly expressed in the diabetic mice, suggesting that the miR-124 was involved in the pathological process of diabetes. In high glucose (HG)-induced Min6 cells, miR-124 regulated the insulin secretion, proliferation and apoptosis of islet β -cells by targeting enhancer of zeste homolog 2 (EZH2), indicating that miR-124 was involved in the pathological process of diabetes by targeting EZH2, which might provide a new idea and target for the diagnosis and treatment of diabetes.

INTRODUCTION

Diabetes is a chronic metabolic disease characterized by hyperglycemia and clinical manifestations including polydipsia, polyuria and weight loss. In China, the incidence of diabetes is increasing, and about 90% of patients with diabetes have type 2 diabetes mellitus (T2DM)^[1]. At present, it is widely accepted that hyperglycemia plays a vital role in the apoptosis and insulin secretion of pancreatic islet β -cells, and dysfunctional islet β -cells contribute to the serious clinical symptoms of T2DM^[2, 3]. However, the exact

mechanisms of T2DM remain poorly understood because of its complex etiology and pathogenesis. It is, therefore, important to explore the pathogenesis of T2DM in depth.

Evidence exists that a variety of micro RNAs (miRNAs), such as miR-375, miR-29 and miR-146 negatively regulate the synthesis and secretion of insulin, and that this process is associated with the development of T2DM^[4, 5]. Several studies have shown that miR-124 is highly expressed in uninephrectomized diabetic rats treated with streptozotocin (STZ). Furthermore, overexpression of miR-124a in the Min6 cell line decreases glucose-stimulated insulin secretion by directly affecting the levels of Rab27A^[6, 7]. At present, evidence is lacking concerning the role and mechanism of miR-124 in diabetes.

Enhancer of zeste homolog 2 (EZH2) is an enzymatic catalytic subunit of polycomb repressive complex 2 that can mediate gene silencing by coordinating with other epigenetic modifying enzymes^[8]. Many studies have shown that miR-124 targets EZH2 to regulate the cell proliferation and apoptosis in a variety of cancer cells and is involved in the occurrence and development of cancers^[9, 10]. Previous studies have shown that with the aging of pancreatic islet cells, the expression of EZH2 gradually decreases, along with a decline in proliferative ability, an increase in apoptosis, and a decrease in insulin levels. These findings suggest that EZH2 plays an important role in the proliferation and apoptosis of pancreatic islet β -cells and the pathogenesis and progression of diabetes^[11]. However, whether miR-124 targeting EZH2 is involved in the occurrence and development of diabetes remains unclear.

This study explored the expression of miR-124 and EZH2 in the pancreatic tissue of diabetic mice and the effect of miR-124 on insulin secretion and the proliferation and apoptosis of pancreatic islet β -cells. The aim was to provide new insights and targets for the diagnosis and treatment of diabetes.

MATERIALS AND METHODS

Animals

Twenty male, specified-pathogen free BALB/c mice aged 6-8 wk, were obtained from Sanxia University(Hubei, China). All mice were fed at 25°C with relative humidity between 60% and 75%, and a 12-h light/dark cycle with free access to food and water. The mice were randomly divided into a normal control group (NC group) and diabetes group (DM group), with 10 mice in each group.

Establishment of the diabetic mouse model

After acclimation for 1 wk, mice in the DM group were fed with a high sucrose/high-fat diet, whilst mice in the NC group were fed with standard chow for 4 wk. All mice were fasted for 12 h and then the mice in the DM group were intraperitoneally injected with STZ solution (45 mg/kg, dissolved in 0.1 mol/L citrate buffer solution; pH4.4) for 3 days to induce T2DM^[12, 13]. Mice in the NC group were injected with an equal volume of the same citrate buffer solution for 3 days. Fasting blood glucose levels were measured by the glucose oxidase method at 72 h after the last injection. Mice with fasting blood glucose levels above 16.7 mmol/L were regarded as having T2DM^[14, 15]. One week after this measurement, the mice were killed under deep anesthesia with excessive ether. Pancreases were immediately dissected and washed with 0.9% cold saline solution and fixed in 10% formalin solution for subsequent experiments.

Hematoxylin and eosin staining

Part of the pancreatic tissue was fixed in 4% paraformaldehyde for 24 h. After dehydration by ethanol gradients, the pancreatic tissue was embedded in paraffin and sliced into 4 µm sections. Then after dewaxing and rehydration, the sections were stained with hematoxylin and eosin (HE). The histopathological changes of the pancreatic tissue were observed under a light microscope.

Immunohistochemistry

Paraffin sections of pancreatic tissue were dewaxed with xylene, rehydrated with ethanol gradients and treated with 3% H₂O₂ solution at 37°C for 15 min. Rabbit anti-

mouse EZH2 and insulin antibody (Abcam, 1:1000) was incubated overnight at 4°C. After being washed by PBS, HRP-labeled goat anti-rabbit IgG secondary antibody (Abcam, 1:1000) was incubated at 37°C for 30 min. DAB solution was added to provide the staining for 6-8 min. Next, the sections were counterstained with hematoxylin and observed under a light microscope. The positive cells of EZH2 or insulin were characterized by brown granules in the nucleus. The expression levels of EZH2 and insulin were scored semi-quantitatively based on staining intensity and the percentage of positive cells using Image J software (National Institutes of Health, USA). Briefly, Immunohistochemistry score = staining intensity × percentage of positive cells. Four score categories in immunostaining intensity: negative (0), weak (1), moderate (2), and strongly (3). Percentage of positive defined as 0=0%-25%; 1=26%-50%; 2=51%-75%; 3=76%-100%^[16]. Five fields were randomly selected in each section.

Cell culture

Mice pancreatic islet β -cells lines Min6 were purchased from Shanghai Meilian Biotechnology Co. Ltd., and the cells were cultured in Dulbecco's Modified Eagle's Medium (DMEM, HyClone) containing 10% fetal bovine serum and 50 mmol/L β -mercaptoethanol in a incubator at 37°C with 5% CO₂. The medium was changed once every other day. When reached 80-90% confluence, the cells were passaged. And the Min6 cells were incubated in DMEM medium containing 25 mmol/L glucose for 24 h to mimic diabetes model for subsequent experiments.

Transfection and grouping

Small interference RNA targeting EZH2 (si-EZH2) and its negative control (si-NC), miR-124 inhibitor and its negative control (inhibitor-NC), and miR-124 mimic and its negative control (mimic-NC) were purchased from Shanghai GenePharma. The cells were divided into a blank control (BC) group, high glucose culture (HG) medium group (Min6 cells were grown in medium with 25 mmol/L glucose), miR-124 inhibitor group (miR-124 inhibitor was transfected into HG-induced Min6 cells), inhibitor NC group

(inhibitor-NC was transfected into HG-induced Min6 cells), miR-124 inhibitor+si-EZH2 group (miR-124 inhibitor and si-EZH2 were co-transfected into HG-induced Min6 cells), and a miR-124 inhibitor+si-NC group (miR-124 inhibitor and si-NC were co-transfected into HG-induced Min6 cells). The cells at logarithmic phase were seeded into a 6-well plate. When the cells reached 70% confluence, the cells were transfected according to the instructions of Lipofectamine 2000. After transfection for 24 h, the following experiments were conducted.

ELISA

The cells were seeded into 24 well plates at a density of 5×10^3 /well and cultured with 1mL Krebs-Ringer bicarbonate (KRB) buffer containing 25 mmol/L glucose for 2 h at 37°C and 5% CO₂. Then the supernatants were collected and insulin content was measured using ELISA according to the instructions of the mouse insulin ELISA kit (Millipore, USA).

CCK-8 assay

Cell viability was detected by the CCK-8 method. The transfected cells were seeded into 96-well plates with five multiple wells in each group. After cell were cultured for 12 h, 24 h, 36 h, and 48 h, 10 µl of CCK-8 solution was added into each well and the cells were incubated for 4 h. Then, the absorbance (OD value) at 450 nm of each well was detected by a microplate reader. The cell proliferation curve was depicted according to time as abscissa and OD value as coordinate.

Flow cytometry

After transfection for 48 h, the cells were washed with PBS and fixed overnight with 70% pre-cooled ethanol at 4 °C For detection of cell cycle distribution, 2 µl of RNase A (0.25 mg/mL) and 500 µl of propidium iodide (PI) dye solution (50 µg/ml) were added to the cells and they were incubated for 30 min in the dark at room temperature before cell cycle distribution was assessed by flow cytometry.

For the detection of cell apoptosis, cell density was adjusted to 1×10^5 /mL. For a 100 μ l cell suspension, 5 μ l Annexin V-FITC was added, and the cells were incubated for 15min at room temperature. Then, 10 μ l of PI was added and the cells were incubated for 5 min in the dark before cell apoptosis was detected by CytoFLEX flow cytometry.

RT-qPCR

The expression levels of miR-124 and EZH2 in pancreatic tissues and cells were detected by RT-qPCR. The tissue samples were ground under liquid nitrogen and the total RNA from tissues and cells were extracted by Trizol (Invitrogen, USA). The transcriptor First Strand cDNA Synthesis kit (Takara, Japan) was used for the reverse transcription of RNA into cDNA. The SYBR Green Master Mix (Takara, Japan) was used for RT-qPCR. The primers were as follows: miR-124, forward: 5'-GTCGTATCCAGTGC GTGTCGTGGAGTCGGCAATTGCACTGGATACGACGGCATTCA-3'; U6, forward: 5'-CTCGCTTCGGCAGCACACA-3'; and reverse: 5'-AACGCTTCACGAATTTGCGT-3'; EZH2, forward: 5'-GTAGACACTC CTCCAAGAA-3', and reverse: 5'-GGTCACAGGGTTGATAGTT-3'; β -actin, forward: 5'-CACGAAACTACCTTCAACTCCAT-3', and reverse: 5'-ATCTTGATCTTCATTGTGCTGGG-3'. The reaction conditions: 95 $^{\circ}$ C 3 min, 95 $^{\circ}$ C 15 s, 58 $^{\circ}$ C 30 s, a total of 40 cycles. The relative expression of miR-124 and EZH2 were calculated by $2^{-\Delta\Delta C_t}$, with U6 and β -actin as the internal references.

Western blot

The total proteins in tissues and cells were extracted by ProteoPrep[®] total protein extraction kit (Millipore, USA), and the protein concentration was detected by the BCA method. The protein samples were separated by SDS-PAGE, and transferred onto PVDF membranes. The membranes were blocked using TBST solution with 5% skim milk for 1h at room temperature. Next, the samples were incubated with first antibodies: EZH2 (Abcam, ab186006, 1:1000), caspase-3 (Abcam, ab13847, 1:500), Bax (Abcam, ab32503,

1:2000) and Bcl-2 (Abcam, ab182858, 1:2000) were incubated overnight at 4°C. After washing with TBST solution, the samples were incubated with the second antibody HRP-IgG (Abcam, ab7090, 1:5000) for 1 h at room temperature. Images were developed by an ECL chemiluminescent reagent (Thermo-Fisher Scientific, USA) and photographed using a gel imaging system (BioRad, USA). The gray value ratios were analyzed using Image J software to represent the relative expression levels.

Double luciferase reporter assay

Targetscan (http://www.targetscan.org/vert_71/) was used to predict the target gene of miR-124. According to the sequence of pmirGLO and the EZH2 gene in PubMed, the wild-type and mutant primers (containing cleavage sites of NheI and SalI) were synthesized. The wild-type and mutant fragment of the EZH2 gene 3'-UTR were amplified to construct the recombinant plasmid pmirGLO-EZH2-WT and pmirGLO-EZH2-MUT. Min6 cells were seeded into 24-well plates. After the cells were cultured for 24 h, pmirGLO-EZH2-WT or pmirGLO-EZH2-MUT plasmids and miR-124 mimic or mimic-NC were co-transfected into the cells. After transfection for 48 h, luciferase activity was detected by the double luciferase reporter assay kit (Promega).

Statistical analysis

All data were analyzed by SPSS21.0. Data with a normal distribution are presented as mean±standard deviation. The comparison among multiple groups was analyzed using one-way analysis of variance, followed by a Tukey post hoc test. In the case of small sample sizes or non-parametric distribution, the difference among groups was analyzed by the Kruskal-Wallis test. The correlation between miR-124 and EZH2 was tested by Pearson correlation analysis. Statistical significance was considered to be $P < 0.05$.

RESULTS

miR-124 was upregulated and EZH2 was downregulated in pancreatic tissues of diabetic mice

16

After 3 days of STZ injection, the fasting blood glucose level of mice in the DM group was more than 16.7 mmol/L. Furthermore, HE staining showed that the pancreatic islets of diabetic mice were atrophic and irregular with a fuzzy boundary and fewer cells, which were disordered and swelling. By contrast, the pancreatic islets of NC mice were oval with a clear boundary and many cells, which were evenly distributed and arranged in the islets (Figure 1A). These findings demonstrated that the diabetic mice models had been successfully established.

13

RT-qPCR was used to detect the expression of miR-124 in pancreatic tissue of mice. The expression of miR-124 in the DM group was significantly higher than that in the NC group (Figure 1B). By contrast, the expression of EZH2 in the DM group was significantly lower than that in the NC group, as detected by RT-qPCR, western blot and immunohistochemistry (Figure 1C, D and E). Pearson correlation analysis showed that the expression of miR-124 was significantly negatively correlated with EZH2 in pancreatic tissue ($r=-0.911$, $P<0.05$) (Figure 1F). Furthermore, immunohistochemical staining showed that insulin expression in the DM group was lower than that of the NC group (Figure 1G).

3

miR-124 was upregulated in HG-induced Min6 cells

Min6 cells were induced by HG to mimic glucolipotoxicity and establish the diabetes cell model^[17]. Firstly, the expression of miR-124 was upregulated in HG-induced Min6 cells detected by RT-qPCR (Figure 2A). Moreover the expression of miR-124 was downregulated by miR-124 inhibitor in HG-induced Min6 cells, suggesting that miR-124 was highly expressed in the diabetes cell model and was downregulated by miR-124 inhibitor.

18

Silencing of miR-124 promoted the proliferation of HG-induced Min6 cells

2

The CCK-8 assay was used to detect the proliferation activity of Min6 cells to explore the effect of miR-124 on the proliferation of HG-induced Min6 cells (Figure 2B). The results showed that the proliferation viability of HG-induced Min6 cells was inhibited,

whereas knockdown of miR-124 significantly improved the proliferation of HG-induced Min6 cells. These findings suggested that silencing of miR-124 promoted the proliferation of HG-induced Min6 cells.

Silencing of miR-124 promoted the HG-induced Min6 cells at the transition from the G0/G1 phase to the S phase of the cell cycle

The cell cycle was detected by flow cytometry to study the effect of miR-124 on the cell cycles. As shown in Figure 2C, HG-induced Min6 cells were arrested at the transition from the G0/G1 phase to the S phase of the cells cycle. In the miR-124 inhibitor group, the proportion of G0/G1 phase cells was significantly reduced and the proportion of S phase cells was significantly increased. This findings suggested that miR-124 inhibitor promoted the cells at the transition from the G0/G1 phase to the S phase of the cell cycle in HG-induced Min6 cells.

Silencing of miR-124 suppressed the apoptosis of HG-induced Min6 cells

As shown in Figure 2D, the apoptosis rate of HG-induced Min6 cells was increased. By contrast, knockdown of miR-124 inhibited the apoptosis of HG-induced Min6 cells. In addition, the expression levels of the apoptosis-related proteins Bax, Bcl-2 and caspase-3 were detected by western blot (Figure 2E). In the HG group, the expression of Bax and caspase-3 were increased, while the expression of Bcl-2 was decreased. In the miR-124 inhibitor group, the opposite effect was shown. miR-124 inhibitor inhibited the increase in expression of Bax and caspase-3 and the decrease in expression of Bcl-2 induced by HG. All results emphasized the pro-apoptotic effect of miR-124 in HG-induced Min6 cells.

Silencing of miR-124 promoted insulin secretion in HG-induced Min6 cells

To further explore the potential function of miR-124, insulin secretion was detected by ELISA (Figure 2F). The levels of insulin secretion decreased in HG-induced Min6 cells. In addition, insulin content was increased in the miR-124 inhibitor group in response to

HG stimulus, suggesting that miR-124 inhibitor promoted the insulin secretion in HG-induced Min6 cells.

miR-124 targeted EZH2

TargetScan predicted that there was a ¹⁰ binding site for miR-124 in the 3'-UTR region of the EZH2 gene (Figure 3A). Furthermore, a double luciferase reporter gene assay showed that in Min6 cells transfected with the pmirGLO-EZH2-WT plasmid, the miR-124 mimic reduced the fluorescence intensity ($P < 0.05$). This result suggested ¹⁵ that miR-124 could bind to the 3'-UTR of the EZH2 gene and inhibit luciferase activity. The results indicated that EZH2 was a target gene of miR-124.

Silencing of miR-124 protected pancreatic β -cell function by targeting EZH2

¹ Firstly, RT-qPCR and western blot were used to detect the expression of EZH2 in Min6 cells. The expression of EZH2 was downregulated in HG-induced Min6 cells, and the miR-124 inhibitor could upregulate EZH2 expression in HG-induced Min6 cells (Figure 3B and C). To further investigate whether the miR-124 inhibitor protected pancreatic β -cell function by targeting EZH2, the si-EZH2 and miR-124 inhibitor were co-transfected into HG-induced Min6 cells. In this experiment, the elevation of EZH2 expression induced by the miR-124 inhibitor was inhibited by si-EZH2 (Figure 4A and B). In addition, effect on the proliferation, insulin secretion, cell cycle and apoptosis were investigated (Figure 4 C-F). si-EZH2 reduced proliferation activity and insulin secretion in HG-induced Min6 cells transfected with miR-124 inhibitor. The result of flow cytometry showed that si-EZH2 inhibited the promotioal effect of miR-124 inhibitor on the cell cycle. Moreover, si-EZH2 reversed the inhibitory effect of the miR-124 inhibitor on apoptosis of HG-induced Min6 cells. Together, the results indicated that silencing of miR-124 promoted the proliferation, insulin secretion, induced ¹ the transition from the G0/G1 phase to the S phase of the cell cycle, and suppressed the apoptosis of HG-induced Min6 cells by targeting EZH2.

DISCUSSION

Diabetes is a metabolic disorder disease characterized by chronic hyperglycemia. The pathological features of diabetes include insulin resistance and a secretory deficiency in pancreatic islets β -cell. Hyperglycemia is the direct clinical manifestation of diabetes, which can induce pancreatic β -cell apoptosis and decrease β -cell proliferation^[18]. Dysfunctional β -cells result in low insulin secretion and subsequently accelerated development of diabetes^[19]. In this study, mice were fed with a high sucrose/high-fat diet combined with a low-dose STZ injection to establish the mouse model of T2DM^[12]. To model diabetic mouse islets β -cells, Min6 cells were grown in HG medium to mimic glucolipotoxicity. These cells exhibited more suppressive effects on β -cell viability and proliferation compared with cells grown in normal medium, which is consistent with the findings of previous studies^[20]. Insulin secretion also decreased in HG-induced Min6 cells. This decrease might result from hypofunction of HG-induced Min6 cells, leading to decreased glucose uptake and, therefore, decreased insulin secretion^[21].

A variety of miRNAs were abnormally expressed in patients with diabetes and affected insulin secretion by regulating the differentiation of islet β -cells, or affecting insulin synthesis and secretion^[22]. Kong *et al*^[23] found that miR-9, miR-29a, miR-30d, miR-34a, miR-124a, miR-146a, and miR-375 Levels were significantly elevated in the serum of individuals with T2DM. Guo *et al*^[24] also found that miR-124 was highly expressed in T2DM patients and involved in glucose and lipid metabolism in diabetes. In this study, miR-124 was upregulated in the pancreatic tissue of diabetic mice and in HG-induced Min6 cells. Inhibition of miR-124 expression in HG-induced Min6 cells promoted proliferation activity and insulin secretion and also inhibited cell apoptosis. The results suggested that miR-124 was involved in the occurrence and development of diabetes by regulating insulin secretion, proliferation and apoptosis of pancreatic islets β -cells.

Previous studies of miR-124 have found that miR-124 regulates the expression of SNAP25, Rab3A, Noc2 or other target genes to mediate insulin secretion of islets β -

cells^[25]. Weng *et al*^[26] found that insulin secretion levels were decreased in ISD-INS-1 cells and speculated that upregulation of miR-124 was one mechanisms by which insulin secretion was reduced in INS-1 cells. Another study also reported that miR-124 could negatively target FOXA2 to regulate the function of islet β -cells^[27]. Parsa E *et al*^[28] pointed out that the levels of miR-124a in plasma were significantly reduced in patients with diabetes compared with individuals without diabetes and that miR-124a may affect the pathogenesis and progression of diabetes by regulating TRIB3. In the current study, EZH2 was downregulated in the pancreatic tissue of diabetic mice, which was negatively correlated with the level of miR-124, consistent with the results of Lu *et al*^[11]. Furthermore, bioinformatics and a double luciferase reporter gene assay confirmed that EZH2 was one of the target genes of miR-124 and was negatively regulated by miR-124. Neo *et al*^[29] showed that EZH2 was downregulated in neural P19 cells and negatively correlated with the expression of miR-124, and miR-124 promoted the differentiation of P19 by targeting EZH2. Ma *et al*^[30] found that miR-124 played an anti-tumor role in cholangiocarcinoma by regulating the EZH2-STAT3 signal axis and inducing autophagy. Tian *et al*^[31] showed that miR-124 negatively regulated the expression of EZH2 to inhibit the proliferation, migration and invasion of esophageal cancer cells. In order to explore whether miR-124 regulated the proliferation and apoptosis of Min6 cells by targeting EZH2, the miR-124 inhibitor and si-EZH2 were co-transfected into Min6 cells and the biological characteristics were detected. The results showed that miR-124 could target EZH2, to inhibit insulin secretion and proliferation activity and induce apoptosis in HG-induced Min6 cells.

CONCLUSION

In conclusion, this study found that miR-124 was highly expressed in diabetic mice and HG-induced Min6 cells. miR-124 inhibited insulin secretion and cell proliferation, promoted apoptosis of islet β -cells *via* targeting EZH2 and was involved in the pathological process of diabetes. This study indicates a new target for the treatment or research into the pathogenesis of diabetes. However, the study had some limitations:

First, how EZH2 promotes insulin secretion was not studied. Furthermore, one miRNA can potentially target several genes. Therefore, miR-124 might target other genes to regulate islet β cells function and the development of diabetes. Further experiments are needed to elaborate the role and mechanism of miR-124 in diabetes. ⁹ Currently, long non-coding RNAs (lncRNAs) are another important type of RNA involved in the pathogenesis of several diseases in cooperation with miRNAs. Several lncRNAs cooperate with miR-124 to regulate diseases, including cancer or stroke. Therefore, the lncRNA that regulates miR-124 in T2DM should be investigated.

ARTICLE HIGHLIGHTS

Research background

Diabetes is a metabolic disorder disease characterized by chronic hyperglycemia. The pathological features of diabetes include insulin resistance and a secretory deficiency in pancreatic islets β -cell. In clinical studies, a variety of miRNA are involved in the occurrence and development of diabetes. And miR-124 is highly expressed in the serum of patients with diabetes and in pancreatic islet β -cells. So the role and mechanism of action of miR-124 in diabetes was explored in the T2DM mice and HG-induced Min6 cells.

Research motivation

A variety of miRNAs were abnormally expressed in patients with diabetes and affected insulin secretion by regulating the differentiation of islet β -cells, or affecting insulin synthesis and secretion. So what is the role and mechanism of miR-124 in diabetes.

Research objectives

² The present study aimed to investigate the expression of miR-124 in diabetic mice and the potential mechanism of action in islet β -cells.

Research methods

4
mice were fed with a high sucrose/high-fat diet combined with a low-dose STZ injection to establish the mouse model of T2DM. The expression levels of miR-124 and enhancer of zeste homolog 2 (EZH2) in pancreatic tissues of diabetic mice were detected. The targeted relationship between miR-124 and EZH2 was predicted by Targetscan software and verified by a double luciferase reporter assay.

To model diabetic mouse islets β -cells, Min6 cells were grown in HG medium to mimic glucolipotoxicity. The insulin secretion, proliferation, cell cycle and apoptosis of HG-induced Min6 cells were detected after interference of miR-124a and/or EZH2.

Research results

The expression of miR-124 was upregulated and EZH2 was downregulated in the pancreatic tissue of diabetic mice compared with control mice, and the expression of miR-124 was negatively correlated with that of EZH2. miR-124 was highly expressed in HG-induced Min6 cells. Inhibition of miR-124 promoted insulin secretion and cell proliferation, induced the transition from the G0/G1 phase to the S phase of the cell cycle, and inhibited cell apoptosis in HG-induced Min6 cells. EZH2 was one of the targets of miR-124. Co-transfection of miR-124 inhibitor and siRNA-EZH2 could reverse the effects of the miR-124 inhibitor in HG-induced Min6 cells.

Research conclusions

miR-124 is highly expressed in diabetic mice and HG-induced Min6 cells and regulates insulin secretion, proliferation and apoptosis of islet β -cells by targeting EZH2.

Research perspectives

miR-124-EZH2 axis might be one of the pathogenesis mechanisms of diabetes, providing a new target for the treatment of diabetes.

10%

SIMILARITY INDEX

PRIMARY SOURCES

1	www.ncbi.nlm.nih.gov Internet	150 words — 3%
2	www.spandidos-publications.com Internet	79 words — 2%
3	Jianhui Li, Ye Nie, Wenlong Wu, Xinjun Lei, Tianchen Zhang, Zhenzhen Mao, Yanfang Wang, Zhuolin Cao, Kaishan Tao, Wenjie Song. "The mechanism by which tertiary lymphoid structures in tumours affect prognosis and immunotherapy in patients with hepatocellular carcinoma", Research Square Platform LLC, 2022 Crossref Posted Content	27 words — 1%
4	lipidworld.biomedcentral.com Internet	27 words — 1%
5	rnajournal.cshlp.org Internet	21 words — < 1%
6	www.mdpi.com Internet	21 words — < 1%
7	research-repository.griffith.edu.au Internet	20 words — < 1%
8	www.termedia.pl Internet	19 words — < 1%

9 Huizhen Zheng, Xinying Li, Xinyu Yang, Fei Yan, Chuan Wang, Jinbo Liu. "miR-217/Mafb Axis Involve in High Glucose-Induced β -TC-tet Cell Damage Via Regulating NF- κ B Signaling Pathway", Biochemical Genetics, 2020

18 words — < 1%

Crossref

10 www.hindawi.com

Internet

17 words — < 1%

11 Yanzhi Lu, Min Long, Zhaowei Gao, Chong Liu, Ke Dong, Huizhong Zhang. "Long non-coding RNA ENST00000469812 promotes Enterovirus type 71 replication via targeting the miR-4443/NUPR1 axis in rhabdomyosarcoma cells", Archives of Virology, 2022

15 words — < 1%

Crossref

12 hepatitiscnewdrugs.blogspot.fr

Internet

15 words — < 1%

13 Yongqi Wang, Xiaoqing Wang, Hong Zhang, Biao Han, Yuanmei Ye, Mengjie Zhang, Yingbin Wang, Jianjun Xue, Chun'ai Wang. "Transforming Growth Factor- β 1 Promotes M1 Alveolar Macrophage Polarization in Acute Lung Injury by Up-Regulating DNMT1 to Mediate the microRNA-124/PELI1/IRF5 Axis", Frontiers in Cellular and Infection Microbiology, 2021

14 words — < 1%

Crossref

14 Jin Guo, Hong-zhu Li, Lu-chuan Wang, Wei-hua Zhang, Guang-wei Li, Wen-jing Xing, Rui Wang, Chang-qing Xu. "Increased expression of calcium-sensing receptors in atherosclerosis confers hypersensitivity to acute myocardial infarction in rats", Molecular and Cellular Biochemistry, 2012

13 words — < 1%

Crossref

15 Shiyong Lin, Qianwen Liu, Jing Wen, Kunhao Bai, Yandong Guo, Jing Wang. "STAT3-mediated TH17 cell Differentiation was Related to the Carcinogenesis of Colitis Related Cancer", Research Square, 2020

13 words — < 1%

Crossref Posted Content

16 www.nature.com
Internet

13 words — < 1%

17 Shujun Liu, Guigang Yan, Junfu Zhang, Lianzhi Yu. "Knockdown of Long Noncoding RNA (lncRNA) Metastasis-Associated Lung Adenocarcinoma Transcript 1 (MALAT1) Inhibits Proliferation, Migration, and Invasion and Promotes Apoptosis by Targeting miR-124 in Retinoblastoma", Oncology Research Featuring Preclinical and Clinical Cancer Therapeutics, 2018

12 words — < 1%

Crossref

18 Xin Lv, Jun Qiu, Tao Hao, Haoran Zhang, Haiping Jiang, Yang Tan. "The Mechanism of Catalpol in Alzheimer's Disease by Modulating Mitochondrial Function Through miR-124-Mediated STIM2", Research Square Platform LLC, 2021

12 words — < 1%

Crossref Posted Content

EXCLUDE QUOTES ON

EXCLUDE SOURCES < 12 WORDS

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES < 12 WORDS