

81513_Auto_Edited.docx

Ca²⁺/cAMP ratio: An inflammatory index for diabetes, hypertension and COVID-19

Bergantin L. Ca²⁺/cAMP ratio as an inflammatory index

Leandro Bergantin

Abstract

Ca²⁺/cAMP ratio could serve as an inflammatory index for diseases like hypertension, diabetes, and coronavirus disease 2019.

Key Words: Ca²⁺/cAMP ratio; COVID-19; Inflammatory index; Diabetes

Bergantin L. Ca²⁺/cAMP ratio: An inflammatory index for diabetes, hypertension and COVID-19. *World J Diabetes* 2023; In press

Core Tip: Ca²⁺/cAMP ratio could serve as an inflammatory index for diseases like hypertension, diabetes, and coronavirus disease 2019.

TO THE EDITOR

There is a clear relationship between Ca²⁺ signaling, *e.g.*, increased Ca²⁺ signals, and inflammatory responses^[1,2]. Considering the cumulative data from the scientific literature, including data of high evidence such as meta-analysis and systematic reviews, we can now link a Ca²⁺ dyshomeostasis as an upstream factor for hypertension, diabetes, and other inflammatory processes^[1,2]. In fact, severe inflammatory outcomes are described to be linked to a critical coronavirus disease 2019 (COVID-19) result^[1,3]. Intriguingly, some reports have also observed an increased severity of COVID-19 in patients with diabetes^[1,3]. To assess this issue, a meta-analysis^[3] was performed by conducting a literature review of Scopus, PubMed, Science Direct, and Web of Science. Observational studies, case-reports, and case-series reports that

analyzed diabetes in COVID-19 patients were included in this meta-analysis^[3]. The authors concluded that diabetes is a risk factor and plays a role in the disease severity and in the mortality of individuals with COVID-19. In fact, a bidirectional relationship between COVID-19 and diabetes has been established^[4]. Recent data have shown that severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) can produce a direct damage to the pancreas that could worsen hyperglycemia, and even cause the onset of diabetes in previously non-diabetic subjects^[4]. Like diabetes, hypertension has also been recognized as a prevalent cardiovascular comorbidity in patients infected with COVID-19^[5]. It is well established that hypertension increases the severity of SARS-CoV-2-infected patients^[5]. In addition, obesity is a well-known risk factor for metabolic syndrome (MetS), hypertension, and diabetes. Recent reports revealed that it is also a vital risk factor for COVID-19, as demonstrated by a recent phenome-wide analysis of COVID-19^[6].

Moreover, a link between infection with viruses and a Ca^{2+} dyshomeostasis is well-discussed, *e.g.*, altering host cellular processes in the benefit of the viruses^[7-10]. Thus, a Ca^{2+} dyshomeostasis induced by viruses may trigger an alteration of the host cellular system that benefits virus survival and could serve as a link for an increased severity of COVID-19 in patients with diabetes^[1,3].

An interesting longitudinal study also evaluated the relationship between serum Ca^{2+} levels and the incidence of MetS, diabetes, and hypertension^[11]. This study^[11] was performed through cross-sectional and longitudinal analyses (period 2010–2016). Logistic regression was used for cross-sectional analysis of the association between serum Ca^{2+} levels or albumin-corrected calcium (ACCA) and the prevalence of MetS, diabetes, or hypertension. Receiver operating characteristic curve analysis was applied for calculating an optimal cut-off value of serum Ca^{2+} levels and ACCA^[11]. Cox proportional regression for the development of MetS, diabetes, and hypertension according to different cut-off values of serum Ca^{2+} levels and ACCA were performed. At baseline, there were 27364 participants in this study^[11]. The authors^[11] concluded that higher serum Ca^{2+} levels were associated with an increased risk of MetS, diabetes,

and hypertension. A hypothesis which could link hypertension and higher serum Ca^{2+} levels postulates that ¹ the influx of Ca^{2+} into the smooth muscle of the arteries could lead to muscle contracture, then increasing vascular resistance and, therefore, could lead to the development of hypertension^[11]. Our previous reports^[1,2,12-15] also discussed this issue. In fact, we postulated that a dysregulation of Ca^{2+} signaling is linked to a sympathetic hyperactivity, then leading to hypertension^[1,12,16]. In addition, a hypothesis which could link diabetes and higher serum Ca^{2+} levels postulates that serum Ca^{2+} ¹ is associated with an insulin resistance in adipocytes and skeletal muscle^[17,18]. Our previous reports are in accordance with this hypothesis^[1,12,15]. In fact, whereas a physiological increase in the cytoplasmic concentration of Ca^{2+} is a significant trigger for releasing insulin, an abnormal elevation of Ca^{2+} could stimulate β -cell apoptosis, then decreasing insulin levels, contributing to diabetes^[1,15].

Furthermore, reports of our group undoubtedly established ⁷ that a rise of the concentration of cAMP can stimulate a Ca^{2+} release from endoplasmic reticulum, entitled as Ca^{2+} /cAMP signalling interaction^[1,2,12-15]. ² In fact, a rise of the concentration of Ca^{2+} can markedly cause a decrease of the concentration of cAMP because of the negative feedback (Ca^{2+} /cAMP signalling interaction). Thus, a disruption of this interaction can be linked with disorders, *e.g.*, hypertension, diabetes, COVID-19^[1,2,15]. ² Bearing in mind that the interaction between these disorders may be linked with continued increases of the concentration of Ca^{2+} , if these increases could disturb Ca^{2+} /cAMP signalling interaction needs more studies, *e.g.*, in animal models and clinical trials. Indeed, previous studies corroborate this concept. For instance, in patients with diabetes, plasma concentrations of cAMP were significantly lower than those of normal subjects^[19]. In addition, cAMP contents of platelets were measured and observed to be lower in hypertensive than in normotensive subjects^[20]. Figure 1 summarizes previous discussion.

In conclusion, Ca^{2+} /cAMP ratio could serve as an inflammatory index for diseases like hypertension, diabetes, and COVID-19^[21,22].

2

ACKNOWLEDGEMENTS

Dr. Bergantin would like to thank the reviewers and editors for their suggestions to improve this manuscript.

2

Figure 1 Altered Ca^{2+} /cAMP ratio resulted from an inflammatory stimulus and infection with SARS-CoV-2. Altered Ca^{2+} /cAMP ratio stimulates uncontrolled inflammation, leading to virus replication, increase of β -cell apoptosis and decrease of insulin levels, and sympathetic hyperactivity. Up/down arrows: Increase/decrease. COVID-19: Coronavirus disease 2019.

46%

SIMILARITY INDEX

PRIMARY SOURCES

- 1

www.ncbi.nlm.nih.gov
Internet

126 words — 12%
- 2

Leandro Bueno Bergantin. " Mental Disorders and Poor COVID-19 Prognosis: Reevaluating the Relationship through Ca /cAMP Signalling ", Current Topics in Medicinal Chemistry, 2022
Crossref

98 words — 10%
- 3

www.esp.org
Internet

77 words — 8%
- 4

Leandro B. Bergantin. "The Interplay Among Epilepsy, Parkinson's Disease and Inflammation: Revisiting the Link through Ca²⁺/cAMP Signalling", Current Neurovascular Research, 2021
Crossref

37 words — 4%
- 5

Leandro B. Bergantin. "A Timeline of Ca²⁺/cAMP Signalling: From Basic Research to Potential Therapeutics for Dementia", Current Alzheimer Research, 2022
Crossref

25 words — 2%
- 6

Nikolai Khaltsev, Svetlana Akselrod. "Role of Global Alliance against Chronic Respiratory Diseases (GARD) in achievement of the UN sustainable development goals (SDG) and targets", Journal of Thoracic Disease, 2021
Crossref

21 words — 2%

-
- 7 Leandro Bueno Bergantin. "The clinical link between depression and obesity: Role of Ca²⁺/cAMP signalling", Psychiatry Research, 2020 20 words — 2%
Crossref
-
- 8 Mei Peng, Jia He, Ying Xue, Xue Yang, Shao Liu, Zhicheng Gong. "Role of Hypertension on the Severity of COVID-19: A Review", Journal of Cardiovascular Pharmacology, 2021 15 words — 1%
Crossref
-
- 9 hdl.handle.net 11 words — 1%
Internet
-
- 10 novapublishers.com 11 words — 1%
Internet
-
- 11 digilib.uinsby.ac.id 8 words — 1%
Internet
-
- 12 A. Murphy, A. Gheorghe, U. Griffiths, P. Perel, H. Legido-Quigley, P. Lamptey. "PM033 The Economic Burden of CVD and Diabetes in Low- and Middle-Income Countries: A Systematic Literature Review", Global Heart, 2016 7 words — 1%
Crossref
-
- 13 Marcos M. Lima-Martínez, Carlos Carrera Boada, Marialaura D. Madera-Silva, Waleskha Marín, Miguel Contreras. "COVID-19 and diabetes: A bidirectional relationship", Clínica e Investigación en Arteriosclerosis (English Edition), 2021 6 words — 1%
Crossref
-
- 14 Matteo Di Maso, Serena Delbue, Maurizio Sampietro, Monica Ferraroni et al. "Biomarkers for predicting COVID-19 mortality using the joint modelling approach", Research Square Platform LLC, 2023 6 words — 1%

EXCLUDE QUOTES	OFF	EXCLUDE SOURCES	OFF
EXCLUDE BIBLIOGRAPHY	OFF	EXCLUDE MATCHES	OFF