

\xB2\xE9\xD6\xD8/90171_Auto_Edited
-.docx

Name of Journal: *World Journal of Diabetes*

Manuscript NO: 90171

Manuscript Type: ORIGINAL ARTICLE

Clinical Trials Study

Effect of special types of bread with select herbal components on postprandial glucose levels in diabetic patients

Special diet in diabetic patients

Drasko . M Gostiljac, Srdjan S Popovic, Vesna Dimitrijevic-Sreckovic, Sasa Ilic, Jelena Jevtovic, Dragan M Nikolić, Ivan Soldatovic

Abstract

BACKGROUND

The abstract of the revised manuscript does not contain this paragraph

AIM

The objective was to compare organoleptic characteristics and effects of two types of bread on postprandial glucose and insulin levels in patients with type 2 diabetes mellitus (T2DM).

METHODS

This trial included 97 patients with T2DM. A parallel group of 16 healthy subjects was also investigated.

All participants were given 50 g of rye bread and the same amount of a special type of bread (STB) with an herbal mixture on two consecutive days. Postprandial blood glucose and insulin levels were compared at the 30th, 60th, 90th and 120th minutes. A

questionnaire was used for subjective estimation of the organoleptic and satiety features of the two types of bread.

RESULTS

Compared with those in patients who consumed rye bread, significantly lower postprandial blood glucose and insulin concentrations were found in patients with T2DM who consumed special types of bread. No relevant differences were found among the healthy subjects. Subjectively estimated organoleptic and satiety characteristics are better for special types of bread than for rye bread.

CONCLUSION

Special types of bread (STB) have better effects than rye bread on postprandial glucoregulation in patients with T2DM. Subjectively estimated organoleptic and satiety characteristics are better for STB than for rye bread. STB can be recommended for nutrition in patients with T2DM.

Key Words: special types of bread; postprandial glucoregulation; insulin; nutrition; patients with type 2 diabetes mellitus (T2DM)

Gostiljac D.M, Popovic SS, Dimitrijevic-Sreckovic V, Ilic S, Jevtovic J, Nikolić DM, Soldatovic I. Effect of special types of bread with select herbal components on postprandial glucose levels in diabetic patients. *World J Diabetes* 2024; In press

Core Tip: Bread that we are testing is novel, tasty and very effective in population with compromised glucoregulation. In our study we have compared novel bread with standard hospital rye bread and obtained significant differences regarding glucose metabolism of TopiGluk, compared to rye bread. This research might be interesting for readers because this mixture TopiGluk could become standard supplement in bread for patients with compromised glucoregulation and those in high risk for diabetes.

INTRODUCTION

Diabetes mellitus is one of the main health issues worldwide, and its incidence has been increasing for several decades ^[1,2]. Additionally, a large number of people have reduced glucose tolerance ^[3]. A Global Report on Diabetes showed that the number of adult diabetes mellitus patients has increased almost fourfold, ³ from 108 million patients in 1980 to 422 million patients in 2014 ^[4]. This dramatic increase is mostly due to type 2 diabetes mellitus (T2DM) and its risk factors, especially obesity ^[5].

Dietary interventions are the main economic and effective strategies aimed at reducing blood glucose and insulin levels in the population ^[3,6]. Physical activity also improves glucose tolerance as a supplemental treatment for obesity but not as a replacement for dietary measures ^[7]. Controlled nutrition improves glucose tolerance by reducing endogenous glucose production and improving sensitivity to insulin ^[8]. The consumption of foods with a low glycemic index (GI) decreases postprandial blood glucose and insulin levels and their fluctuations ^[9,10].

One of the basic foodstuffs worldwide is bread. Recommendations for this population include the consumption of rye or whole grain bread with higher dietary fiber content instead of white bread ^[11]. Several trials have shown that a positive effect on glucoregulation may also be ¹ achieved by enriching food with various biologically active substances of herbal origin ^[12].

TopiGluk bread was patented by the National Intellectual Property Office (registration No. 73932) as a supplementary treatment for people with disease. TopiGluk bread is made by adding a mixture of nutrients to whole wheat, oat and buckwheat flour.

We hypothesized that TopiGluk bread would significantly improve glucose tolerance ⁵ in patients with T2DM.

The aim of this study was to evaluate the effects of TopiGluk bread on postprandial glucose levels in patients with T2DM. Additionally, the insulin levels and organoleptic properties of the examined bread were evaluated as secondary objectives.

MATERIALS AND METHODS

Subjects and Methods

Time and place of study

A prospective study was conducted at the Clinic of Endocrinology, Diabetes and Metabolic Disorders, Clinical Center of Serbia, Belgrade. The study was carried out from 20 May 2016 to 25 March 2017.

Subjects

The trial included 97 patients with T2DM who were treated as outpatients at the Clinic of Endocrinology, Diabetes and Metabolic Disorders, Faculty of Medicine, University of Belgrade. The inclusion criteria were an age older than 18 years and confirmed T2DM. The exclusion criteria were acute T2DM complications; acute inflammatory conditions; chronic diseases of the liver or gastrointestinal tract; malignant diseases; immunodeficiency; narcotic or alcoholic addictions; and pregnancy (Figure 1).

A parallel group of 16 healthy controls were analyzed; they had no history of metabolic disorders (including diabetes mellitus) and had preserved glucose homeostasis based on their medical history.

All the subjects provided informed consent for inclusion in the study. The investigation was carried out in accordance with the Declaration of Helsinki and was revised in 2013. The study was carried out in accordance with the EU Directive 20/2001/EC, the Commission Directive 2005/28/EC, and the European Parliament Declaration on Good Clinical Practice, as well as the International Conference on Harmonization ICH-GCP (E6); approval for this research was given by the Ethics Committee, Clinical Center of Serbia, No. 105/39 on 19 May 2016. The study is registered in the German Clinical Trials Register DRKS00023611. The clinical trial was registered after completion because the law of the Republic of Serbia does not require registration in the international registry. The authors confirm that all ongoing and related trials for this food supplement have been registered.

Study protocol

Prior to the trial, a detailed history was taken, which included data on physical activity and lifestyle, the duration of T2DM and therapy, chronic complications of T2DM and nutritional habits. Data on anthropometric characteristics, weight and height were obtained from medical records if no significant changes were observed in the last six months. Body mass index was calculated by dividing the body weight in kilograms by the squared body height in meters (kg/m^2).

All participants were instructed to fast for at least 12 h prior to blood sampling.

A slice of bread weighing 50 g was provided to each participant, which is the usual portion for patients with T2DM. The nutritional content of the bread is presented in Table 1. One day, all participants were given a piece of rye bread, and on another day, they were given a piece of special bread. Patients were advised to perform similar activities on both days. Furthermore, 12 h of fasting prior to each blood sampling was obligatory.

Participants were instructed to complete a questionnaire about the look, smell, taste, quantity, and satiety of the bread. Answers were given on a numerical scale from 1 (worst) to 5 (best). Upon arrival at the clinic at approximately 08:00 in the morning, the responsible researcher accommodated each participant in the laboratory. A venous cannula was inserted in the arm of the patient. The first blood sample was taken before the meal. Next, four blood samples were taken every 30 minutes after the meal. Healthy subjects were with evaluated with a glucose meter, and blood samples were collected using finger sticks (capillary blood sampling) in the infirmary.

The laboratory analysis of the T2DM patients included HbA1c and C-peptide levels (at baseline only) and glucose and insulin levels (at baseline and 30 minutes, 60 minutes, 90 minutes and 120 minutes after the meal) on both examination days. Blood glucose measurements were performed using the spectrophotometric (hexokinase) method with a COBAS 6000 (ROCHE diagnostics) with a reference range of 3.9-6.1 mmol/L. Insulin

and C-peptide levels were measured using an immunoradiometric assay method with a gamma counter for *in vitro* diagnosis (LKB-WALLAC ChinGamma 1272).

The laboratory analysis of healthy controls included only glucose levels at baseline (before the meal) and at 30 minutes, 60 minutes, 90 minutes and 120 minutes after the meal. Blood measurements from capillary blood were performed with an electrochemical method using a BIOSEN C LINE machine (EKG diagnostics) with a reference range of 3.9-6.1 mmol/L.

Due to differences in sampling methods, diabetic and healthy subjects were not compared, except for blood sugar changes from baseline to 120 minutes (the sampling method did not affect the change).

Differences between 30 minutes and baseline, 60 minutes and baseline, 90 minutes and baseline and 120 minutes and baseline levels were calculated (delta glucose and delta insulin). The area under the curve (AUC) was calculated using a trapezoidal model [13].

Ingredients of special types of bread

The bread was a brand of the Delhaize Serbia distributor in cooperation with partners for product development—the international company Puratos and the national bakery products manufacturer Alimpije—and with a partner for product improvement—ZZ Zdravlje (Čačak, Serbia), which is the name of the author of TopiGluk®.

The ingredients of TopiGluk® bread are as follows: Basil (*Ocimum basilicum*), garlic (*Allium sativum*), Greek seed (*Trigonella foenum graecum*), ginger (*Zingiber officinale*), oat (*Avena sativa*), Jerusalem artichoke (*Helianthus tuberosus*), and cinnamon (*Cinnamomum verum*). Sunflower and linen seeds are among the supplements. The nutritional content of 100 g of the special type of bread (STB) with the TopiGluk mixture is presented in Table 1. Approximately one-third of the mentioned mixture contains active principles (Topi Gluk), while the remaining two-thirds are equivalent parts buckwheat, oat and whole wheat flour. The representation of the components in Topi Gluk was 0.5% garlic, 0.5% ginger, 2% basil, 3% Greek seed, 5% cinnamon, 8% oat, and 13% Jerusalem artichoke.

Statistical analysis

A sample size of 97 patients with T2DM achieved 90% power to detect a mean difference of 1.0 ± 3.0 mmol/L with a significance level for alpha error of 0.05 using a two-sided paired samples t test.

The results are presented as counts (%), means \pm standard deviations depending on the data type and distribution. Measurements were compared using parametric (paired sample t test) and nonparametric (Wilcoxon signed rank test) tests. Glucose changes (delta) between healthy subjects and patients were compared using an independent samples t test. Deltas were calculated as the difference between the 30 min, 60 min, 90 min and 120 min time points and baseline. The area under the curve was calculated using the trapezoidal rule. All p values less than 0.05 were considered significant. All the data were analyzed using R 3.4.2. (R Core Team (2017). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>).

RESULTS

The majority of participants were males in the sixth to seventh decade of life, preobese to obese individuals, and physically active individuals. The majority of patients were receiving oral antidiabetic therapy, while one-third were receiving insulin. Coronary disease and neuropathy were present in one-third of patients. The average duration of diabetes was shorter than 10 years. Most of the patients consumed rye bread, for which the median daily intake was 1 slice, while other bread types (whole wheat bread and white bread) were less represented in terms of nutritional status (both medians were 0) (Table 2).

All the evaluated characteristics were greater for the special type of bread (STB) than for rye bread, as presented in Table 3.

In patients with T2DM, we found better glucoregulation after the STN portion than after the rye bread portion. Significantly lower delta values of blood glucose were

observed after the STB portion at the 90th and 120th minutes than after the rye bread portion. In healthy subjects, no significant differences in blood glucose delta values were observed, except at the 90th minute. Concerning the area under the curve of blood glucose (AUC), a significant difference was found only at the 90th minute. In T2DM patients, the area under the curve (AUC) was 21.2±5.1% after the rye bread portion and 19.6±4.5% after the STB with TopiGluk portion ($p < 0.001$). In healthy subjects, no significant differences in the AUC were found after the two types of bread were consumed ($P = 0.924$) (Figure 2).

When comparing patients and healthy subjects regarding delta values, all differences were significant at the 0.001 Level for both the rye bread and the STB with the TopiGluk[®] mixture, except for the change from baseline to 30 minutes after the rye ($P = 0.919$) and STB with TopiGluk[®] ($P = 0.313$) portions (Figure 2).

The insulin levels of the patients were analyzed at each time point (0, 30, 60, 90 and 120 minutes) and for each type of bread (Figure 3). Initially, the median insulin concentrations were identical. However, significant differences in insulin levels were observed at the 90th minute and the 120th minute after the two types of bread were consumed. After the STB with TopiGluk[®] was consumed, we found significantly lower median insulin concentrations compared to after rye bread was consumed. A comparison of the deltas revealed significant differences at the same time points. The median value and interquartile range of the AUC of insulin in patients with T2DM were 46.7 (29.7-61.1) after consuming rye bread and 39.9 (30.1-57.0) after consuming the STB with TopiGluk[®] ($P = 0.035$).

Diabetic patients were divided into two subgroups: patients receiving oral antidiabetic therapy and patients receiving insulin therapy (Figure 4). In patients on oral antidiabetic therapy, significantly higher blood glucose levels were observed at the 60th, 90th and 120th minutes after the rye bread was consumed than after the STB with TopiGluk[®] was consumed. Differences in blood glucose changes from baseline were significant at the 90th minute and at the 120th minute. In patients receiving insulin therapy, all mean glucose levels were significantly greater after rye bread was

consumed than after the STB with TopiGluk® was consumed, but the changes (deltas) were significantly different only at the 90th minute and at the 120th minute (Figure 4).

Table 1. Comparative nutritional composition of the rye bread and TopiGluk bread (100 g)

DISCUSSION

This study compared the effects of rye bread an (STB) with TopiGluk® on glucoregulation. Patients subjectively assessed the STB as better than the rye bread regarding appearance, aroma, taste, amount, satiety level and duration of satiety. We observed lower in glycemic and insulin levels from baseline to the 120th minute in diabetic patients after the STB was consumed than after the rye bread was consumed. In healthy subjects, this difference was not observed, as expected.

The special type of bread with TopiGluk® can be described as a “functional food” because it may have the following positive effects on glucose metabolism: a reduction in ¹glucose absorption from the intestine to blood, an increase in glucose utilization, an increase in cell sensitivity to insulin, a reduction in insulin resistance, an increase in endogenous insulin production and an increase in glycogen in the liver ^[14].

Glucose absorption has been evaluated in several studies. Glucose absorption can be reduced using foods rich in dietary fibers (DFs). DFs include ³lignin and a range of polysaccharides derived from cell walls that are poorly digested in the upper intestine ^[15]. Patients with diabetes mellitus are usually instructed to eat rye bread, which is rich in DF. ²Viscous and gel-forming soluble DFs inhibit macronutrient absorption and reduce the postprandial glucose response. ²However, in prospective cohort studies, insoluble cereal DF and whole grains, but not soluble DF, are consistently associated with reduced diabetes mellitus risk, suggesting that further unknown mechanisms are likely involved ^[16,17]. TopiGluk® contains whole wheat flour, buckwheat and oatmeal, which are rich in DF, especially beta glucans. TopiGluk is also rich in the soluble DF inulin from Jerusalem artichoke, which binds water and forms a viscous solution that delays gastric emptying and intestinal transit, thus reducing glucose absorption. This

leads to a decreased blood glucose response [16,17]. DF also decreases insulin secretion and reduces the chance of reactive hypoglycemia during the postprandial period. On the other hand, this promotes satiety and satiation, increases fat oxidation and decreases fat storage [18].

Several authors have examined the effect of whole-meal and whole-kernel rye breads on glucose metabolism compared to that of white wheat bread. Leinonen *et al* concluded that whole kernel rye bread has no effect on the glucose response but has an effect on the postprandial insulin response [19]. In healthy subjects, we found no differences in blood glucose levels after the consumption of the STB with TopiGluk® or after consumption of the rye bread. This is probably due to preserved regulatory mechanisms of glucose metabolism in healthy subjects.

However, in patients with T2DM, we found lower blood glucose and insulin levels after the consumption of the STB with TopiGluk® than after the consumption of the rye bread. This approach is very important for individuals with impaired glucose tolerance or an increased risk of diabetes. Starchin bread decomposes during digestion to simple sugars, which affect glucoregulation. TopiGluk is a mixture of natural metabolically active ingredients that play a proven role in metabolic regulation. Studies in diabetic rats revealed the significant effect of basil on glucose metabolism and its potential usefulness in treating humans with T2DM (20). This effect is likely mediated by decreased glucose absorption and glucose mobilization in the liver [20]. A recent meta-analysis revealed that garlic, in addition to therapy, contributes to improved glucose and lipid control [21,22]. Fenugreek is mostly produced in Eastern countries, but in the West, it is often used as a medicinal herb or spice. A recent meta-analysis revealed that fenugreek acutely reduces postprandial glucose levels, but the long-term effect on glucose metabolism has not been sufficiently tested [23]. A meta-analysis by Huang FY *et al* revealed that ginger has no significant effect on fasting glucose levels but significantly improves HbA1c levels [24]. Another meta-analysis by Hou Q *et al* revealed the significant effect of oat on fasting blood glucose, HbA1c and lipid levels [25]. Oat is a common choice for people with obesity, digestive problems or diabetes. It is a grain

with a low glycemic index and is rich in vitamins and minerals. Jerusalem artichoke is a sunflower plant that ¹ affects blood sugar levels favorably. Its most valuable component is inulin. Dry-ground Jerusalem artichoke contains 70% inulin. Fructose is a result of inulin hydrolysis and is an excellent replacement for glucose. A study of diabetic rats revealed that Jerusalem artichoke improved metabolism, microcirculation, and blood vessel conditions and thereby reduced the severity of diabetes complications [26]. A recent study of newly diagnosed diabetes mellitus patients revealed a positive effect of Jerusalem artichoke on postprandial glucose levels [27]. Cinnamon, a spice with high global consumption, also affects fasting blood glucose levels in patients with T2DM as well as in prediabetic patients [12,28,29]. While choosing biologically active substances for TopiGluk[®], numerous factors should be considered; the most ¹ important are the availability of the substance, its impact on glucoregulation, other favorable metabolic effects, its applicability in the technological process of bread preparation and manufacturing, and the ¹ impact of certain phases of the technological procedure; for example, the ¹ impact of the bread baking temperature on the active principles of the ingredients (duration), specific aroma and taste at planned concentrations, mutual compatibility and cumulative effect on glucoregulation.

A comparison of the organoleptic properties of two kinds of bread, an STB with TopiGluk[®] and rye bread, revealed that the former was better than the latter. A number of respondents emphasized the sweet taste of the STB with TopiGluk[®], which may be particularly important for patients with diabetes mellitus because they are usually not allowed to consume sweet food.

The results of our research agree with the latest data from the literature indicating that the plants we added to the TopiGluk[®] bread (*Ocimum basilicum*, *Allium sativum*, *Trigonella foenum graecum*, *Zingiber officinale*, *Avena sativa*, *Helianthus tuberosus*, and *Cinnamomum verum*) have antidiabetic properties [30–35].

CONCLUSION

Based on the present results, we can conclude that postprandial blood glucose levels in patients with T2DM are lower after consuming Topi Gluk bread than after consuming the same amount of rye bread. Improved glucoregulation was noted in patients with T2DM at 90 and 120 minutes, both in patients who were taking oral antidiabetic drugs and in patients receiving insulin therapy. A special type of bread made with TopiGluk® has better subjectively assessed organoleptic and fine characteristics than does rye bread. The STB with TopiGluk® can be recommended as part of the diet in patients with T2DM.

ARTICLE HIGHLIGHTS

Research background

Bread that we are testing is novel, tasty and very effective in population with compromised glycoregulation. In our study we have compared novel bread with standard hospital rye bread and obtained significant differences regarding glucose metabolism of special types of bread (STB) with TopiGluk, compared to rye bread.

Research motivation

All participants were given 50 g of rye bread, and the same amount of special types of bread with herbal mixture (STB) on two consecutive days. In the continuation of these studies, it would be interesting to increase the amount of tested bread and see how it would affect postprandial glycemia.

Research objectives

To compare organoleptic characteristics of two sorts of bread on postprandial glucose and insulin levels in patients with type 2 diabetes melitus.

Research methods

This trial included 97 patients with T2DM. A parallel group of 16 healthy subjects was also investigated.

All participants were given 50 g of rye bread and the same amount of a special type of bread (STB) with an herbal mixture on two consecutive days. Postprandial blood glucose and insulin levels were compared at the 30th, 60th, 90th and 120th minutes. A questionnaire was used for subjective estimation of the organoleptic and satiety features of the two types of bread.

Research results

Compared with those in patients who consumed rye bread, significantly lower postprandial blood glucose and insulin concentrations were found in patients with T2DM who consumed special types of bread. No relevant differences were found among the healthy subjects. Subjectively estimated organoleptic and satiety characteristics are better for special types of bread than for rye bread.

Research conclusions

Special types of bread (STB) have better effects than rye bread on postprandial glucoregulation in patients with T2DM. Subjectively estimated organoleptic and satiety characteristics are better for STB than for rye bread. STB can be recommended for nutrition in patients with T2DM.

Research perspectives

The results of this research can be the basis and incentive for future research that would determine which biochemical substances from plant components added to special types of bread (STB) are responsible for the effect of postprandial glycemia.

ACKNOWLEDGEMENTS

The authors wish to thank the Puratos and Delhaize companies for the administrative and technical support.

ORIGINALITY REPORT

6%

SIMILARITY INDEX

PRIMARY SOURCES

1	juniperpublishers.com Internet	105 words — 3%
2	www.science.gov Internet	40 words — 1%
3	www.ncbi.nlm.nih.gov Internet	33 words — 1%
4	barcelona2019.euromedlab.org Internet	14 words — < 1%
5	zenodo.org Internet	14 words — < 1%
6	joe.endocrinology-journals.org Internet	12 words — < 1%

EXCLUDE QUOTES ON
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES < 12 WORDS
EXCLUDE MATCHES < 12 WORDS