

Name of Journal: *World Journal of Diabetes*

ESPS Manuscript NO: 23692

Manuscript Type: Retrospective Cohort Study

Does parity worsen diabetes-related chronic complications in women with type 1 diabetes?

Running title: Parity in women with type 1 diabetes

Marilia Brito Gomes, M.D, PhD†, Carlos Antonio Negrato, M.D, PhD††, Ana Almeida, M.D, PhD†, Antonio Ponce de Leon, Ph.D †††

On behalf of the Brazilian Type 1 Diabetes Study Group (BrazDiab1SG)

## Match Overview

1	<b>CrossCheck</b> 322 words "Abstracts of the 50th EASD Annual Meeting", Diabetologia, 2014.	5%
2	<b>Internet</b> 98 words crawled on 03-Sep-2015 <a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a>	2%
3	<b>CrossCheck</b> 58 words Rodacki, M., L. Zajdenverg, J. R. Dantas, J. E. P. de Oliveira, R. R. Luiz, R. A. Cobas, C. C. S. Palma, C. A. Negrato	1%
4	<b>CrossCheck</b> 44 words Qin, Jian-Ping. "Contrast enhanced computed tomography and reconstruction of hepatic vascular system for tra ...	1%
5	<b>Internet</b> 36 words crawled on 21-Dec-2015 <a href="http://journals.plos.org">journals.plos.org</a>	1%
6	<b>CrossCheck</b> 27 words Tannus, Lucianne Righeti Monteiro, Karla Rezende Guerra Drummond, Eliete Leão da Silva Clemente, Maria de	<1%
7	<b>CrossCheck</b> 18 words Garn, Satish K. and Irl B. Hirsch. "Self-Monitoring of Blood	<1%

学术搜索

找到约 188 条结果 (用时0.08秒)

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☒ 包含引用

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

## Does pregnancy increase the risk for development and progression of diabetic nephropathy?

M Miodovnik, BM Rosenn, JC Khoury... - American journal of ..., 1996 - Elsevier

... diabetes mellitus does not increase the risk of subsequent nephropathy and does not accelerate ... did not develop (n = 123), Significance, ESRD developed (n = 12), ESRD did not develop (n ... prospective clinical trials have shown that modification of these factors can ameliorate the ...

被引用次数: 99 相关文章 所有 8 个版本 引用

## Factors Associated With Preterm Delivery in Women With Type 1 Diabetes A cohort study

J Lepercq, J Coste, A Theau, D Dubois-Laforgue... - Diabetes ..., 2004 - Am Diabetes Assoc

... reference groups regarding ethnicity, social class, marital status, smoking, prepregnancy, BMI, parity, the occurrence ... A higher rate of LGA infants would have been expected as glycemic control was ... Our study did not allow the identification of the mechanisms of this association....

被引用次数: 62 相关文章 所有 5 个版本 引用

## Periconceptional care of women with diabetes mellitus

G Leguizamón, ML Igarzabal, EA Reece - Obstetrics and gynecology ..., 2007 - Elsevier

... With these limitations in mind, three general observations can be made: (1) ... a CABG before pregnancy while 38% maternal or fetal death occurred in those who did not receive ... The authors concluded that parity does not influence the long-term prevalence or severity of diabetic ...

被引用次数: 30 相关文章 所有 5 个版本 引用

## [HTML] Metabolic syndrome components worsen lower urinary tract symptoms in women with type 2 diabetes

HC Tai, SD Chung, CH Ho, TY Tai... - The Journal of ..., 2013 - press.endocrine.org

... It seemed that MS did not impair the detrusor contractility nor increase the bladder outlet ... Clinically, OAB symptom (urgency, frequency and nocturia) can be measured with storage symptom part ... This result suggests that neuropathy alone does not explain the full amount of future ...

被引用次数: 40 相关文章 所有 6 个版本 引用

## Effect of medical therapy on progressive nephropathy: influence of pregnancy, diabetes and hypertension

G Leguizamon, EA Reece - Journal of Maternal-Fetal and ..., 2000 - informahealthcare.com



学术搜索

找到约 188 条结果 (用时0.10秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利

☒ 包含引用

☐ 创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

## Does pregnancy increase the risk for development and progression of diabetic nephropathy?

M Miodovnik, BM Rosenn, JC Khoury... - American journal of ..., 1996 - Elsevier

... diabetes mellitus does not increase the risk of subsequent nephropathy and does not accelerate ... did not develop (n = 123). Significance, ESRD developed (n = 12), ESRD did not develop (n ... prospective clinical trials have shown that modification of these factors can ameliorate the ...

被引用次数: 97 相关文章 所有 9 个版本 引用 保存

## Metabolic syndrome components worsen lower urinary tract symptoms in women with type 2 diabetes

HC Tai, SD Chung, CH Ho, TY Tai... - The Journal of ..., 2010 - press.endocrine.org

... It seemed that MS did not impair the detrusor contractility nor increase the bladder outlet ... Clinically, OAB symptom (urgency, frequency and nocturia) can be measured with storage symptom part ... This result suggests that neuropathy alone does not explain the full amount of future ...

被引用次数: 45 相关文章 所有 5 个版本 引用 保存

## Periconceptional care of women with diabetes mellitus

G Leguizamón, ML Igarzabal, EA Reece - Obstetrics and gynecology ..., 2007 - Elsevier

... With these limitations in mind, three general observations can be made: (1). ... a CABG before pregnancy while 38% maternal or fetal death occurred in those who did not receive ... The authors concluded that parity does not influence the long-term prevalence or severity of diabetic ...

被引用次数: 31 相关文章 所有 5 个版本 引用 保存

## [HTML] Factors Associated With Preterm Delivery in Women With Type 1 Diabetes A cohort study

J Lepercq, J Coste, A Theau, D Dubois-Laforgue... - Diabetes ..., 2004 - Am Diabetes Assoc

... reference groups regarding ethnicity, social class, marital status, smoking, prepregnancy, BMI, parity, the occurrence ... A higher rate of LGA infants would have been expected as glycemic control was ... Our study did not allow the identification of the mechanisms of this association. ...

被引用次数: 61 相关文章 所有 7 个版本 引用 保存

## Effect of medical therapy on progressive nephropathy: influence of pregnancy, diabetes and hypertension

G Leguizamón, EA Reece - Journal of Maternal-Fetal Medicine, 2000 - Taylor & Francis

... Furthermore, pregnancy per se does not appear to worsen the natural progression to end-stage renal disease for most ... 2 years, 26% of patients in the control group developed overt nephropathy, while in the study group only 9% did. ... Numerous drugs can achieve this goal safely ...

被引用次数: 40 相关文章 所有 5 个版本 引用 保存

## [HTML] Event rates, hospital utilization, and costs associated with major complications of



全部

新闻

图片

视频

更多 ▾

搜索工具

找到约 8,730,000 条结果 (用时 0.74 秒)

## Google 学术: Does parity worsen diabetes-related chronic complications in women with type 1 diabetes?

... the risk for development and progression of diabetic ... - Miodovnik - 被引用次数: 97

... urinary tract symptoms in women with type 2 diabetes - Tai - 被引用次数: 45

Periconceptional care of women with diabetes mellitus - Leguizamón - 被引用次数: 31

## Breastfeeding in Women With Type 1 Diabetes

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > ... > PubMed Central (PMC) ▾ 翻译此页

作者: C Sparud-Lundin - 2011 - 被引用次数: 23 - 相关文章

2011年1月20日 - On multivariable analysis, type 1 diabetes did not remain an independent predictive factor. ... postpartum period, affects the likelihood of long-term breastfeeding. ... childbearing woman in the same gestational week and parity (primi- or ... of birth-related complications that affects the breastfeeding duration.

## Complications of diabetes mellitus - Wikipedia, the free encyc...

[https://en.wikipedia.org/.../Complications\\_of\\_diabetes\\_mellitus](https://en.wikipedia.org/.../Complications_of_diabetes_mellitus) ▾ 翻译此页

The complications of diabetes mellitus are far less common and less severe in people who ... Ketoacidosis is much more common in type 1 diabetes than type 2. ... Several studies also show diabetes associated with a worse disease course and ... Diabetic nephropathy, damage to the kidney which can lead to chronic renal ...

缺少字词: parity

## Care during pregnancy for women with type 1 or 2 diabetes m...

[www.uptodate.com/.../care-during-pregnancy-for-women-with...](http://www.uptodate.com/.../care-during-pregnancy-for-women-with...) ▾ 翻译此页

2015年9月28日 - Most women with type 1 diabetes will require two to four insulin injections per day ... treat any medical complications related to diabetes (such as