

Name of Journal: *World Journal of Diabetes*

ESPS Manuscript NO: 27249

Manuscript Type: Letter to the Editor

Response to Comment on: **2** Statin use and risk of diabetes mellitus

Bharti Chogtu, Rahul Magazine, KL Baiyy

Abstract

In letter to the editor "Comment on: Statin use and risk of diabetes mellitus" authors found the statement "pravastatin 40 mg/day reduced the risk of diabetes by 30% in West of Scotland Coronary Prevention study" erroneous. As per our opinion the statement is right but had been referenced incorrectly.

Key words: Pravastatin; Diabetes mellitus

Core tip: The statement "Pravastatin has shown to decrease the risk of developing diabetes by 30%" is correct and had been wrongly referenced in "Statin use and risk of diabetes mellitus" by Chogtu et al

Letter to the Editor

We thank Eren MA et al for showing interest in the review article.^[1] Eren MA and co-authors ^[2] have pointed to an apparent factual error in the following statement:

"pravastatin 40 mg/day reduced the risk of diabetes by 30% in West of Scotland Coronary Prevention study". However, in our opinion, this statement is correct if we

Match Overview

- | Match Number | Source | Words | Similarity |
|--------------|------------------------------------------------------------------------------------------------------------------------------------|----------|------------|
| 1 | CrossCheck | 19 words | 6% |
| | Sattar, Naveed, and Marja-Riitta Taskinen. "Statins are di...
etogenic â€" Myth or reality?", <i>Atherosclerosis Supplement</i> | | |
| 2 | Internet | 13 words | 4% |
| | crawled on 17-Jul-2016
www.wjnet.com | | |

[全部](#)[新闻](#)[图片](#)[视频](#)[地图](#)[更多 ▾](#)[搜索工具](#)

找到约 2,240,000 条结果 (用时 0.93 秒)

Statin use and risk of diabetes mellitus in postmenopausal women in ...

www.ncbi.nlm.nih.gov/pubmed/22231607 ▾ [翻译此页](#)

作者: AL Culver - 2012 - 被引用次数: 335 - [相关文章](#)

2012年1月9日 - [Statin use](#) and [risk of diabetes mellitus](#) in postmenopausal women in the Women's Health Initiative. Culver AL(1), Ockene IS, ... [Comment](#) in Arch Intern Med. 2012 Jun 11;172(11):896; author [reply](#) 896-7. Arch Intern Med.

Statin Use and Risk of Diabetes Mellitus in Postmenopausal Women ...

archinte.jamanetwork.com/article.aspx?articleid=1108676 ▾ [翻译此页](#)

作者: AL Culver - 2012 - 被引用次数: 335 - [相关文章](#)

2012年1月23日 - [Comments](#) (4) ... [Statin use](#) at baseline was associated with an increased [risk](#) of DM ([hazard ratio](#) [Risk of Diabetes Mellitus \(DM\)](#) by [Statin Use](#) Among Women With and The filled radio buttons indicate your [responses](#).

Response to Comment on Stegman et al. High-Intensity Statin ...

care.diabetesjournals.org/cgi/content/full/38/2/e28 - [翻译此页](#)

High-Intensity [Statin Therapy](#) Alters the Natural History of [Diabetic](#) Coronary ... We agree that the [diabetic](#) population is heterogeneous in terms of cardiovascular [risk](#), [Diabetes mellitus](#) is a major negative determinant of coronary plaque ...

Largest Risk for Diabetes With Statins Yet Seen, in New Study

www.medscape.com/viewarticle/840884 ▾ [翻译此页](#)

2015年3月4日 - [Statin therapy](#) appears to increase the [risk](#) for type 2 [diabetes](#) by 46%, ... Asked to [comment](#), Alvin C Powers, MD, from Vanderbilt University ...

Statins Associated With Significant Increase in Diabetes Risk

www.medscape.com/viewarticle/756617 ▾ [翻译此页](#)

2012年1月9日 - January 9, 2012 (Boston, Massachusetts) — [Statin use](#) in postmenopausal women is associated with a significantly increased [risk of diabetes mellitus](#), research ... [Statin use](#) and [diabetes risk](#) was also observed in women across a range of ... any conflicts of interest related to your [comments](#)

[全部](#)[新闻](#)[图片](#)[视频](#)[地图](#)[更多 ▾](#)[搜索工具](#)

找到约 2,620,000 条结果 (用时 1.02 秒)

Statin Use and Risk of Diabetes Mellitus in Postmenopausal Women ...

archinte.jamanetwork.com/article.aspx?articleid=1108676 ▾ [翻译此页](#)

作者: AL Culver - 2012 - 被引用次数: 351 - [相关文章](#)

2012年1月23日 - [Comments](#) (4) ... [Statin use](#) at baseline was associated with an increased [risk](#) of DM ([hazard ratio](#) [Risk of Diabetes Mellitus \(DM\) by Statin Use](#) Among Women With and The filled radio buttons indicate your [responses](#).

Statin use and risk of diabetes mellitus in postmenopausal women in ...

www.ncbi.nlm.nih.gov/pubmed/22231607 ▾ [翻译此页](#)

作者: AL Culver - 2012 - 被引用次数: 350 - [相关文章](#)

2012年1月9日 - [Statin use](#) and [risk of diabetes mellitus](#) in postmenopausal women in the Women's Health Initiative. Culver AL(1), Ockene IS, ... [Comment](#) in Arch Intern Med. 2012 Jun 11;172(11):896; author [reply](#) 896-7. Arch Intern Med.

Response to Comment on Stegman et al. High-Intensity Statin ...

care.diabetesjournals.org/content/38/2/e28 - [翻译此页](#)

作者: B Stegman - 2015 - 被引用次数: 23 - [相关文章](#)

It is important to note that this [high-risk](#) patient subset still demonstrated significant atheroma ... High-intensity [statin therapy](#) alters the natural history of [diabetic](#) coronary ... [Diabetes mellitus](#) is a major negative determinant of coronary plaque ...

Comment on: Statin use and risk of diabetes mellitus

[www.ncbi.nlm.nih.gov/NCBI/Literature/PubMed Central \(PMC\)](http://www.ncbi.nlm.nih.gov/NCBI/Literature/PubMedCentral(PMC)) - [翻译此页](#)

作者: MA Eren - 2016 - [相关文章](#)

2016年4月25日 - In manuscript named "[Statin use](#) and [risk of diabetes mellitus](#)" by Chogtu et al, authors defined that pravastatin 40 mg/dL reduced the [risk](#) of ...

Largest Risk for Diabetes With Statins Yet Seen, in New Study

www.medscape.com/viewarticle/840884 ▾ [翻译此页](#)

2015年3月4日 - [Statin therapy](#) appears to increase the [risk](#) for type 2 [diabetes](#) by 46% ... Asked to

[全部](#)[新闻](#)[图片](#)[视频](#)[地图](#)[更多 ▾](#)[搜索工具](#)

找到约 2,760,000 条结果 (用时 0.77 秒)

Statin use and risk of diabetes mellitus in postmenopausal women in ...

www.ncbi.nlm.nih.gov/pubmed/22231607 ▾ 翻译此页

作者: AL Culver - 2012 - 被引用次数: 355 - 相关文章

2012年1月9日 - Statin use and risk of diabetes mellitus in postmenopausal women in the Women's Health Initiative. Culver AL(1), Ockene IS, ... Comment in Arch Intern Med. 2012 Jun 11;172(11):896; author reply 896-7. Arch Intern Med.

Statin Use and Risk of Diabetes Mellitus in Postmenopausal Women ...

archinte.jamanetwork.com/article.aspx?articleid=1108676 ▾ 翻译此页

作者: AL Culver - 2012 - 被引用次数: 355 - 相关文章

2012年1月23日 - Comments (4) ... Statin use at baseline was associated with an increased risk of DM (hazard ratio Risk of Diabetes Mellitus (DM) by Statin Use Among Women With and The filled radio buttons indicate your responses.

Increased Diabetes Mellitus Risk With Statin Use: Comment on "Statin ...

archinte.jamanetwork.com/article.aspx?articleid=1108650 ▾ 翻译此页

作者: KL Johansen - 2012 - 被引用次数: 7 - 相关文章

2012年1月23日 - Increased Diabetes Mellitus Risk With Statin Use: Comment on "Statin Use and Risk of Diabetes Mellitus in Postmenopausal Women in the Women's ... Note: You must get at least of the answers correct to pass this quiz.

Response to Comment on Stegman et al. High-Intensity Statin ...

care.diabetesjournals.org/content/38/2/e28 - 翻译此页

作者: B Stegman - 2015 - 被引用次数: 26 - 相关文章

It is important to note that this high-risk patient subset still demonstrated significant atheroma ... High-intensity statin therapy alters the natural history of diabetic coronary ... Diabetes mellitus is a major negative determinant of coronary plaque ...