

Match Overview

1	Crossref 176 words Nunley, Karen A., Caterina Rosano, Christopher M. Ryan, J. Richard Jennings, Howard J. Aizenstein, Janice C. Z	4%
2	Crossref 15 words "Behavioral Medicine, Clinical Nutrition, Education, and Exercise", Diabetes, 2014.	<1%
3	Crossref 13 words Abhimanyu Beri. "Association Between Statin Therapy ... and Tendon Rupture: A Case-Control Study", Journal of	<1%
4	Crossref 13 words Joosten, Hanneke, Sipke T. Visser, Marlies E. van Eersee I, Ron T. Gansevoort, Henk J. G. Bilo, Joris P. Slaets, an	<1%
5	Internet 12 words crawled on 15-Feb-2017 eprints.whiterose.ac.uk	<1%
6	Crossref 10 words "Epidemiology/Genetics", Diabetes, 2014.	<1%

Name of Journal: *World Journal of Diabetes*

ESPS Manuscript NO: 31210

Manuscript Type: Original Article

4
Observational Study

2
Statin use and cognitive function in middle-aged adults with type 1 diabetes

Karen A Nunley, Trevor J Orchard, Christopher M Ryan, Rachel Miller, Tina Costacou, Caterina Rosano

Abstract

AIM: To test associations between statin use and cognitive impairment in adults with childhood-onset type 1 diabetes (T1D).

METHODS: In 2010-13, N=108 middle-aged participants from ongoing observational

[全部](#)[新闻](#)[图片](#)[购物](#)[地图](#)[更多 ▾](#)[搜索工具](#)

找到约 58,300 条结果 (用时 0.92 秒)

Google 学术: Statin use and cognitive function in middle-aged adults with childhood-onset type 1 diabetes

Type 1 diabetes through the life span: a position ... - Chiang - 被引用次数: 251

Intensive diabetes therapy and carotid intima-media ... - Diabetes Control and ... - 被引用次数: 784

Childhood adversity heightens the impact of later-life ... - Kiecolt-Glaser - 被引用次数: 216

Clinically Relevant Cognitive Impairment in Middle-Aged Adults With ...

www.ncbi.nlm.nih.gov/pubmed/26153270 ▾ 翻译此页

作者: KA Nunley - 2015 - 被引用次数: 8 - 相关文章

2015年7月7日 - 1. Diabetes Care. 2015 Sep;38(9):1768-76. doi: 10.2337/dc15-0041. Impairment in Middle-Aged Adults With Childhood-Onset Type 1 Diabetes. ... Readers may use this article as long as the work is properly cited, the use is ...

Management of Type 1 Diabetes in Older Adults

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) ▾ 翻译此页

作者: R Dhaliwal - 2014 - 被引用次数: 7 - 相关文章

2014年2月12日 - The clinical characteristics of youth-onset type 1 diabetes and type 1 diabetes ... early in the course of the disease and with onset of type 1 diabetes in adulthood, ... Older adults may also have other medical comorbidities, functional, Statin therapy is indicated in healthy older adults and frail older adults ...

Management of Type 1 Diabetes in Older Adults | Diabetes Spectrum

spectrum.diabetesjournals.org/content/27/1/9 - 翻译此页

作者: R Dhaliwal - 2014 - 被引用次数: 7 - 相关文章

Older adults with type 1 diabetes are at high risk for severe hypoglycemia and ... Type 1 diabetes is



Statin use and cognitive function in middle-aged adults with type 1 diabetes



全部

新闻

图片

购物

地图

更多

设置

工具

找到约 152,000 条结果 (用时 0.86 秒)

Google 学术: Statin use and cognitive function in middle-aged adults with type 1 diabetes

Diabetes in older adults - Kirkman - 被引用次数: 546

Diabetes and cognitive dysfunction - McCrimmon - 被引用次数: 305

Cognitive function, dementia and type 2 diabetes ... - Strachan - 被引用次数: 212

Clinically Relevant Cognitive Impairment in Middle-Aged Adults With ...

<https://www.ncbi.nlm.nih.gov/pubmed/26153270> - 翻译此页

作者: KA Nunley - 2015 - 被引用次数: 16 - 相关文章

2015年7月7日 - 1. Diabetes Care. 2015 Sep;38(9):1768-76. doi: 10.2337/dc15-0041. ... Impairment in Middle-Aged Adults With Childhood-Onset Type 1 Diabetes. ... Readers may use this article as long as the work is properly cited, the use is ...

缺少字词: statin

Management of Type 1 Diabetes in Older Adults - NCBI - NIH

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: R Dhaliwal - 2014 - 被引用次数: 10 - 相关文章

2014年2月12日 - Older adults with type 1 diabetes are at high risk for severe ... early in the course of the disease and with onset of type 1 diabetes in adulthood, have better ... With aging and long duration of diabetes, there is impairment of and use of cognitive aids can all be helpful for older patients with type 1 diabetes.

[全部](#)[新闻](#)[图片](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 227,000 条结果 (用时 0.66 秒)

Google 学术: Statin use and cognitive function in middle-aged adults with type 1 diabetes

Diabetes in older adults - Kirkman - 被引用次数: 599

Diabetes and cognitive dysfunction - McCrimmon - 被引用次数: 328

Cognitive function, dementia and type 2 diabetes ... - Strachan - 被引用次数: 229

Clinically Relevant Cognitive Impairment in Middle-Aged Adults With ...

<https://www.ncbi.nlm.nih.gov/pubmed/26153270> - 翻译此页

作者: KA Nunley - 2015 - 被引用次数: 21 - 相关文章

2015年7月7日 - Clinically Relevant Cognitive Impairment in Middle-Aged Adults With Childhood-Onset Type 1 Diabetes. ... Readers may use this article as long as the work is properly cited, the use is educational and not for profit, and the ...

缺少字词: statin

Statin Use and Cognitive Impairment in Patients With Type 1 Diabetes ...

<https://www.ncbi.nlm.nih.gov/pubmed/27046662> - 翻译此页

作者: JM Moryś - 2016 - 被引用次数: 1 - 相关文章

Statin Use and Cognitive Impairment in Patients With Type 1 Diabetes: An ... and ¶Neurology of Adults, Medical University of Gdańsk, Gdańsk, Poland. ... 20% with statin treatment) and 36 age-matched control subjects (77.8% with ...

缺少字词: middle

Statin Use and Cognitive Impairment in Patients With Type 1 Diabetes ...

https://www.researchgate.net/.../289211300_Statin_Use_and_Cognitive_Imp... - 翻译此页

Statin Use and Cognitive Impairment in Patients With Type 1 Diabetes: An Observational ... 20% with statin treatment) and 36 age-matched control subjects (77.8% with ... With Distal Symmetrical