

Name of Journal: *World Journal of Gastrointestinal Surgery*

Manuscript NO: 57659

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Partial pancreatic tail preserving subtotal pancreatectomy for pancreatic cancer: improving glycemic control and quality of life without compromising oncological outcomes

Subtotal pancreatectomy for pancreatic cancer

Abstract

BACKGROUND

Total pancreatectomy (TP) is usually considered as a therapeutic option for pancreatic cancer in which Whipple surgery and distal pancreatectomy are undesirable, but brittle diabetes and poor quality of life (QoL) remain major concerns. A subset of patients who underwent TP even died due to the severe hypoglycemia. For pancreatic cancer

Match Overview

1	Internet 16 words crawled on 28-Sep-2015 pancreas.imedpub.com	<1%
2	Internet 12 words crawled on 04-Sep-2019 www.karger.com	<1%
3	Internet 12 words crawled on 01-Feb-2020 www.frontiersin.org	<1%

Partial pancreatic tail preserving subtotal pancreatectomy for panci



ALL

IMAGES

VIDEOS

14,200 Results

Any time ▾

Pylorus-preserving Total Pancreatectomy for Pancreatic Cancer

https://www.researchgate.net/publication/12707565_Pylorus-preserving_Total...

Pylorus-preserving total **pancreatectomy** (PPTP) and standard TP for **pancreatic cancer** were compared in terms of operative **outcomes**, nutritional recovery, and long-term survival.

Mo1498 Quality of Life in Patients After Total ...

<https://www.researchgate.net/publication/259697084...>

Quality of life after total **pancreatectomy** (TP) is perceived to be poor secondary to insulin-dependent diabetes and **pancreatic** insufficiency. As a result, surgeons may be reluctant to offer TP for ...

Abstracts - PubMed Central (PMC)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3152263>

Distal **pancreatectomy** may be inadequate treatment for post gastric bypass nesidioblastosis. A Child **subtotal pancreatectomy** or duodenal **preserving pancreatectomy** may be indicated in some cases. Optimal **pancreatic** resection volume in this disorder will require further experience and analysis. 57

Postoperative Glycemic Control after Central ...

<https://www.researchgate.net/publication/6627570...>

The incidence of new onset or worsening diabetes is surprisingly low in patients after **partial pancreatectomy** for **cancer**, leading us to question what factors predict diminished **glycemic control** in ...

(PDF) Middle-Preserving Pancreatectomy for Multifocal ...

<https://www.researchgate.net/publication/47742747...>

PDF | Postoperative endocrine and exocrine insufficiencies following traditional pancreatectomies might cause a deterioration of the quality of life and ... Find, read and cite all the research

ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

13,100 Results

Any time ▾

[Middle-Segment-Preserving Pancreatectomy | Request PDF](#)

<https://www.researchgate.net/publication/6425623...>

Middle-preserving pancreatectomy (MPP) is a surgical procedure recently advocated as treatment for multicentric periampullary-pancreatic lesions in which the middle portion of the pancreas is ...

[Pylorus-preserving Total Pancreatectomy for Pancreatic Cancer](#)

https://www.researchgate.net/publication/12707565_Pylorus-preserving_Total...

Pylorus-preserving total pancreatectomy (PPTP) and standard TP for pancreatic cancer were compared in terms of operative outcomes, nutritional recovery, and long-term survival.

[Middle-preserving pancreatectomy for multicentric body ...](#)

<https://www.researchgate.net/publication/26776695...>

Even for multifocal diseases, if the pancreatic body is spared, a middle-preserving pancreatectomy (MPP) can be performed to assure maximal pancreatic function and uncompromised quality of life.

[Abstracts - PubMed Central \(PMC\)](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3152263>

Distal pancreatectomy may be inadequate treatment for post gastric bypass nesidioblastosis. A Child subtotal pancreatectomy or duodenal preserving pancreatectomy may be indicated in some cases.

Optimal pancreatic resection volume in this disorder will require further experience and analysis. 57

[Mo1498 Quality of Life in Patients After Total ...](#)

<https://www.researchgate.net/publication/259697084...>

Quality of life after total pancreatectomy (TP) is perceived to be poor secondary to insulin-dependent diabetes and pancreatic insufficiency. As a result, surgeons may be reluctant to offer TP for ...

[Pancreatic Resection: Effects on Glucose Metabolism ...](#)

<https://www.researchgate.net/publication/11993828...>

[1] [2][3] The correlation of partial pancreatectomy to pancreatic endocrine insufficiency is less completely understood and in particular the impact of pancreatic region of surgery on the outcome

[27th Annual Meeting of the American Pancreatic Association](#)