

Match Overview

- 1** **Crossref** 28 words
Nobuhiro Tsuchiya. "Biomarkers for the early diagnosis of hepatocellular carcinoma", World Journal of Gastroenterology, 20... 1%
- 2** **Crossref** 15 words
Dongling Teng, Keyan Wu, Yunyun Sun, Min Zhang et al. "Significant increased CA199 levels in acute pancreatitis patient..." 1%
- 3** **Crossref** 13 words
Miranda Uzoma, Gaurav Singh, Laurie Kohen. "Green palm plantar vesicular eruption in a patient with hyperbilirubinemia" 1%

Name of Journal: *World Journal of Gastrointestinal Surgery*

Manuscript NO: 58424

Manuscript Type: CASE REPORT

Giant simple hepatic cyst with multiple elevated serum tumor markers: a case report

Simple hepatic cyst with elevated tumor markers

Abstract

BACKGROUND

Simple hepatic cysts are relatively common in adults, and mostly appear as asymptomatic incidental radiologic findings. Occasionally, a large cyst will cause symptoms. Elevations in the serum biomarkers PIVKA-II, CA12-5 and CA19-9 are associated with malignant tumors in the liver or bile ducts. This is the first report to describe a case of hepatic cyst with elevated levels of PIVKA-II and CA12-5.

CASE SUMMARY

Giant simple hepatic cyst with multiple elevated serum tumor mark



ALL

IMAGES

VIDEOS

90,200 Results

Any time ▼

(PDF) A simple hepatic cyst with elevated serum and cyst ...

<https://www.researchgate.net/publication/23316225...>

A **simple hepatic cyst** with **elevated serum** and **cyst** fluid CA19-9 levels: **A case report**.pdf Available via license: CC BY 2.0 Content may be subject to copyright.

[PDF] Nonparasitic Solitary Giant Hepatic Cyst Causing ...

https://www.jstage.jst.go.jp/article/internalmedicine/45/9/45_9_621/_pdf/-char/en

Markedly elevated tumor marker levels have been re-ported even in **histologically** diagnosed simple cysts (11, 12). Iwase et al (13) reported CA19-9 levels in hepatic cyst fluid at least 100 times higher than in normal serum concen-trations. The present patient also had markedly elevated cyst fluid tumor marker levels, and imaging studies indicated a

Adenocarcinoma and infection in a solitary hepatic cyst: A ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4305896>

Mar 28, 2005 · Given the progressive nature of the lesion plus the **high** levels of **tumor markers** in the fluid, malignancy was increasingly likely. The patient therefore underwent an exploratory laparotomy. A 10 cm×10 cm×10 cm cystic lesion containing turbid fluid was found in segments 4, 5, and 8 of the **liver**. The **cyst** was aspirated and then opened.

Cited by: 8

Author: Ching-Chung Lin, Shee-Chan Lin, Wen-Chin...

Publish Year: 2005

[PDF] A Giant Simple Liver Cyst That Caused Increases in Serum ...

<https://jocmr.org/index.php/JOCMR/article/download/2002/885>

A **giant simple liver cyst** that caused an increase in se-rum CA 19-9 and CA 15-3 levels is presented here. Case Report **A 76-year-old woman** who had complains of painless **intra-abdominal mass** was

ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

64,900 Results

Any time ▾

[PDF] [Nonparasitic Solitary Giant Hepatic Cyst Causing ...](#)

https://www.jstage.jst.go.jp/article/internalmedicine/45/9/45_9_621/_pdf/-char/en

Markedly elevated tumor marker levels have been re-ported even in **histologically** diagnosed simple cysts (11, 12). Iwase et al (13) reported CA19-9 levels in hepatic cyst fluid at least 100 times higher than in normal serum concen-trations. The present patient also had markedly elevated cyst fluid tumor marker levels, and imaging studies indicated a

[PDF] [A Giant Simple Liver Cyst That Caused Increases in Serum ...](#)

<https://jocmr.org/index.php/JOCMR/article/download/2002/885>

A giant simple liver cyst that caused an increase in se-rum CA 19-9 and CA 15-3 levels is presented here. Case Report **A 76-year-old woman** who had complains of painless **intra-abdominal mass** was admitted to the clinic. During physical examination, an elastic mass extending from the right upper quadrant to the pelvis was detected. Laboratory tests showed normal liver functions. Serum CA 19-9 ...

Cited by: 1

Author: Bulent Dinc, Ayhan Mesci, Selcan Enver Din...

Publish Year: 2014

(PDF) [A simple hepatic cyst with elevated serum and cyst ...](#)

<https://www.researchgate.net/publication/23316225...>

A simple hepatic cyst with elevated serum and cyst fluid CA19-9 levels: A case report.pdf Available via license: CC BY 2.0 Content may be subject to copyright.

[A Giant Simple Liver Cyst That Caused Increases in Serum ...](#)

europepmc.org/articles/PMC4169093

Dec 01, 2014 · Simple cysts (SCs) of the liver are not associated with the biliary malformations in intrahepatic bile duct biliary. Seen in 0.1% to 7% of adult population, biliary malformations are more common in women. The levels of **glycoprotein-like** tumor markers (**carbohydrate antigen** (CA) 19-9) in the cysts and serum could be high.

[Giant mucinous cystadenoma: a case report | Journal of ...](#)

<https://jmedicalcasereports.biomedcentral.com/...> ▾

Jun 14, 2019 · **Giant** ovarian **cysts** are rarely described in the literature, owing to the availability of