

Is a Preoperative Assessment of the Early Recurrence of ...

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3916679

Nov 05, 2013 · A preoperative CA 19-9 level >228 U/mL, tumor size >3.1 cm, and the presence of pathological preoperative lymph nodes statistically correlated with early recurrence. Together, these...

Cited by: 19 Author: Marco La Torre, Giuseppe Nigri, Annalisa...

Publish Year: 2014

A preoperative risk model for early recurrence after ...

https://pubmed.ncbi.nlm.nih.gov/32321665

The early recurrence, middle recurrence, and late/non-recurrence groups differed in preoperative carbohydrate antigen 19-9 and carcinoembryonic antigen levels, "resectability" on cross-sectional...

Cited by: 2 Author: Shi-wei Guo, Jing Shen, Jun-hui Gao, Xia..

Publish Year: 2020





Preoperative serum carbohydrate antigen 19-9 levels ...

https://europepmc.org/article/MED/34226331 -

Jun 01, 2021 \cdot The optimal cutoff values for **early recurrence** were 70 U/mL and 2.85 cm for carbohydrate antigen (CA) 19-9 levels and tumor size, respectively, on ROC curve analysis. Of 181 patients with CA 19-9 levels of > 70 U/mL, 59 (32.6%) had early recurrence, compared to 39 (17.4%) among 226 patients with CA 19-9 levels of \leq 70 U/mL (p < 0.001).


Author: Sarang Hong, Ki Byung Song, Dae Wook... Publish Year: 2021

Is a Preoperative Assessment of the Early Recurrence of ...

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3916679

Nov 05, 2013 · Carbohydrate antigen 19-9 (CA 19-9) is the most accurate serum biomarker for pancreatic cancer available and has shown some utility as diagnostic and prognostic marker. 9 Several studies have shown that high levels of preoperative CA 19-9 (>300 U/mL) correlate with advanced disease. 10 Additional evidence indicates that serum CA 19-9 is an ...

Cited by: 21 Author: Marco La Torre, Giuseppe Nigri, Annalisa L...
Publish Year: 2014




Feedback

A combination of platelet-to-lymphocyte ratio and ...

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6803228

Δ combination of platelet-to-lymphocyte ratio and carbohydrate antigen 19-9 predict early recurrence



Preoperative serum carbohydrate antigen 19-9 levels predict early







ALL

IMAGES

VIDEOS

11 700 Results

Any time ▼



ancbi.nlm.nih.gov - Nov 05, 2013

Is a Preoperative Assessment of the Early Recurrence of ...

Cited by: 21 Author: Marco La Torre, Giuseppe Nigri, Annalisa ...

Publish Year: 2014

The serum carbohydrate antigen 19-9 (CA 19-9) level has been identified as a prognostic indicator of recurrence and reduced overall survival. The aim of this study was to identify preoperative prognostic factors and to create a prognostic model able to assess the early recurrence risk for patients with resectable PAC.

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3916679

A combination of platelet-to-lymphocyte ratio and ...

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6803228

Jul 08, 2019 - High preoperative carbohydrate antigen (CA) 19-9 is a representative risk factor of ER (13-18), but its predictive ability is not always sufficient, and no other reliable risk factors have been identified. Here, we examined predictive risk factors for ER in PDA patients submitted for macroscopic curative pancreatectomy.

Cited by: 6 Author: Shinichi Ikuta, Takashi Sonoda, Tsukasa ...

Publish Year: 2019

Preoperative Serum CA 19-9 Level as a Predictive Factor ...

https://link.springer.com/article/10.1245/s10434-010-1529-7 •

Jan 06, 2011 - There are few studies about preoperative predictive factors for the recurrence of BTC, but in pancreatic cancer, several studies have shown the correlation between preoperative serum levels of the carbohydrate antigen (CA) 19-9 and the rate of recurrence after curative resection 8-10 CA 19-9 is the anti-blood group antibody, which is produced by bile duct epithelium and detects circulating ...

Cited by: 21 Author: Moon Jae Chung, Kyong Joo Lee, Seung...

Publish Year: 2011

Preoperative Serum CA 19-9 Level as a Predictive Factor