

80552_Auto_Edited -check.docx

Meta-analysis of transanal⁶ *vs* laparoscopic total mesorectal excision of low rectal cancer: Importance of appropriate patient selection

Bhattacharya P²³ *et al.* Transanal *vs* laparoscopic total mesorectal excision of low rectal cancer

Abstract

BACKGROUND

Achieving a clear resection margins for low rectal cancer is technically challenging.¹⁰ Transanal approach to total mesorectal excision (TME)¹ was introduced in order to address the challenges associated with the laparoscopic approach in treating low rectal cancers. However, previous meta-analyses have included mixed population with mid and low rectal tumours when comparing both approaches which has made the interpretation of the real differences between two approaches in treating low rectal cancer difficult.

AIM

To investigate the outcomes of transanal TME (TaTME) and laparoscopic TME (LaTME)²⁶ in patients with low rectal cancer.

METHODS

A comprehensive systematic review of comparative studies was performed in line with PRISMA standards. Intraoperative and postoperative complications, anastomotic leak, R0 resection, completeness of mesorectal excision,¹⁰ circumferential resection margin (CRM), distal resection margin (DRM), harvested lymph nodes, and operation time were the investigated outcome measures.

RESULTS

We included twelve comparative studies enrolling 969 patients comparing TaTME ($n = 969$) and LaTME ($n = 476$) in patients with low rectal tumours. TaTME was associated with significantly lower risk of postoperative complications (OR: 0.74, $P = 0.04$), anastomotic leak (OR: 0.59, $P = 0.02$), and conversion to an open procedure (OR: 0.29, $P = 0.002$) in comparison with LaTME. Moreover, the rate of R0 resection was significantly higher in the TaTME group (OR: 1.96, $P = 0.03$). Nevertheless, TaTME and LaTME were comparable in terms of rate of intraoperative complications (OR: 1.87; $P = 0.23$), completeness of mesorectal excision (OR: 1.57, $P = 0.15$), harvested lymph nodes (MD: -0.05, $P = 0.96$), DRM (MD: -0.94; $P = 0.17$), CRM (MD: 1.08, $P = 0.17$), positive CRM (OR: 0.64, $P = 0.11$) and procedure time (MD: -6.99 min, $P = 0.45$).

CONCLUSION

Our findings indicated that for low rectal tumours, TaTME is associated with better clinical and short term oncological outcomes compared to LaTME. More randomised controlled trials are required to confirm these findings and to evaluate long term oncological and functional outcomes.

Key Words: Total mesorectal excision; Laparoscopic; Transanal; Rectal cancer

INTRODUCTION

The incidence of rectal cancer is increasing making it one of the most common cancers worldwide^[1]. Rapidly evolving use of total mesorectal excision (TME) and neoadjuvant chemotherapy have led to considerable improvements in the outcomes of rectal cancer surgery^[2]. A clear resection margin associated with a high quality TME is important for an ideal oncological resection, reducing the incidence of local or regional recurrence, and increasing survival from cancer^[3,4].

Achieving a negative resection margins during resection of low rectal tumours can be challenging due to existence of diminishing gap between the wall of the rectum and mesorectal fascia towards the anal canal^[5]. This has resulted in worse oncological outcomes associated with resection of lower rectal tumours, in comparison with

resection of middle or high rectal tumours, because of greater incidence of local recurrence and positive resection margin^[6]. Transanal approach to TME was introduced in order to address the challenges associated with the laparoscopic and even open TME in surgical management of low rectal cancers^[7].

In 2020, in a comprehensive meta-analysis of comparative studies, we reported that Transanal TME (TaTME) led to higher R0 resection rate and number of harvested lymph nodes while decreasing rates of positive circumferential resection margin (CRM) and conversion to open procedure when compared to laparoscopic TME (LaTME)^[8]. Moreover, our findings indicated that TaTME and LaTME may have similar risk of perioperative morbidity^[8]. Nevertheless, most of the evaluated studies in the aforementioned meta-analysis compared TaTME and LaTME in patients with middle and low rectal tumours subjecting the findings to bias. Considering the existence of new studies focusing on the clinical outcomes of TaTME and LaTME in management of low rectal cancer, conduction of another meta-analysis is worthwhile in order to help defining more appropriate patient selection.

This study aimed to systematically evaluate the best available comparative evidence surrounding TaTME and LaTME in surgical management of low rectal cancer only and compare the outcome so both approaches using meta-analytical model.

MATERIALS AND METHODS

Study design and selection of eligible studies

In our review protocol, we highlighted the inclusion and exclusion criteria, our methodology, and evaluated outcome measures. This study was carried out in line with standards of Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) statement^[9].

All comparative studies investigating the outcomes of transanal and laparoscopic TME in patients with low cancer were considered for inclusion. A rectal tumour within 6 cm of anal verge was considered as a low rectal tumour. We considered all adult

(aged > 18 years) ⁶ patients undergoing TaTME or LaTME for low rectal cancer. TaTME was the intervention of interest and LaTME was the comparison of interest.

The primary outcome measures were intraoperative and postoperative complications, and anastomotic leak. The investigated primary oncological outcome measures were R0 resection, CRM, positive CRM, distal resection margin (DRM), completeness of mesorectal excision, and number of harvested lymph nodes. Moreover, conversion to open and operative time ⁵ were defined as secondary outcome measures.

Literature search strategy

Following sources: MEDLINE, Web of Science, and CENTRAL were searched by two independent authors. Appendix 1 outlines the used search strategy. ¹⁴ The most recent literature search was carried out on 08 July, 2022. Moreover, we screened ⁷ the reference lists of the included studies and previous review articles in order to identify more relevant articles.

Study selection

Two independent review authors screened the title and abstract of the identified studies. This was followed by retrieval of the full-texts of the related studies and their assessment in line with our inclusion and exclusion criteria. Discrepancies in this stage were addressed by discussion among the reviewers.

Extraction and management of data

We created a data extraction tool and extracted details of study-related data, data regarding demographic characteristics of the included patients in each study and outcome data. Two independent reviewers were involved in this process. ⁹ Disagreements between the authors were resolved following discussion. In case of no resolution, an additional reviewer was consulted.

¹⁴ *Assessment of risk of bias*

The methodological quality of the included studies was assessed by 2 review authors who determined their associated risk of bias using the Newcastle-Ottawa scale^[10] for observational studies and Cochrane's tool^[11] for randomized controlled trials (RCTs). We resolved disagreements in methodological quality assessment by discussion between the reviewers. However, if disagreement remained unresolved, a third reviewer was consulted as an adjudicator.

Summary measures and synthesis

For dichotomous outcome measures the odds ratio (OR) was calculated as the summary measures. For continuous outcome parameters, the mean difference (MD) between the two groups was calculated. If mean values were not reported, we extracted data on median and interquartile range and converted those to mean and standard deviation using Hozo *et al*^[12]'s equation.

The unit of analysis for all of the analyzed outcome measures in this study was an individual participant. We did not require contacting the authors of the included studies to ask for any potential missing information.

Data analysis was carried out via Review Manager 5.4 software^[11]. One author extracted and entered the data into the software and another author cross-checked the data. Random-effects modelling were used for analysis of all outcomes. We reported outcome of analyses in Forest plots with 95% confidence intervals (CIs).

The Cochran Q test (χ^2) was used to assess between-study heterogeneity. We calculated I^2 and used the following guide for interpreting the degree of heterogeneity: 0% to 50% might not be important; 50% to 75%: May represent moderate heterogeneity; 75% to 100% may represent substantial heterogeneity. Moreover, we constructed funnel plots for any outcome synthesis involving more than 10 studies.

We performed sensitivity analyses to assess for potential sources of heterogeneity and evaluate the robustness of our findings. Finally, we conducted leave-one-out sensitivity analysis to assess the effect of each study on the overall effect size and heterogeneity.

RESULTS

The literature search resulted in 2120 articles. Following further assessment of the aforementioned articles, 12 comparative studies (2 randomised and 10 observational studies)^[13-24] met the inclusion criteria (Figure 1). The included studies enrolled 969 patients of whom 493 underwent TaTME and the remaining 476 patients had LaTME for rectal cancer.

Table 1 presents the included studies related data. Table 2 presents baseline demographic and clinical characteristics of the included patients. The patients in the transanal and laparoscopic groups were of similar age ($P = 0.53$), gender ($P = 0.19$), and BMI ($P = 0.68$). No significant difference was found between the TaTME and LaTME groups in rectal cancer stages I ($P = 0.29$), II ($P = 0.30$) and III ($P = 0.95$). Furthermore, the mean distance of the tumour to the anal verge in the TaTME and LaTME groups were $3.4 \text{ cm} \pm 1.4 \text{ cm}$ and $3.6 \text{ cm} \pm 1.5 \text{ cm}$, respectively, which was not significantly different ($P = 0.07$). Neoadjuvant chemotherapy was carried out similarly between two groups ($P = 0.22$).

Methodological appraisal

The methodological assessment of 10 observational studies is presented in Table 3. In 7 studies, the risk of bias was low and in 3 studies it was moderate. Moreover, the outcome of methodological assessment of the included randomized controlled trials is demonstrated by Figure 2.

Data synthesis

Outcomes are summarised in Figures 3 and 4.

Intraoperative complications: Six studies (382 patients) reported intraoperative complications as an outcome. The rate of intraoperative complications in the TaTME and LaTME were 7.3% and 4.2%, respectively. There was no significant difference in

intraoperative complications **between** TaTME and LaTME (OR: 1.87; 95%CI, 0.68-5.18, $P = 0.23$). There was **low between-study heterogeneity** ($I^2: 6\%$, $P = 0.36$).

Postoperative complications: Eleven studies (923 patients) reported postoperative complications as an outcome. The rate of overall postoperative complications in the TaTME and LaTME were 30.0% and 35.9%, respectively. TaTME significantly reduced postoperative complications when compared to LaTME (OR: 0.74; 95%CI, 0.56-0.99, $P = 0.04$). There was moderate heterogeneity among the included studies ($I^2: 2\%$, $P = 0.42$).

Anastomotic leak: This outcome was reported by eleven studies (896 patients). Anastomotic leak occurred in 10.1% and 15.5% of patients in the TaTME and LaTME groups, respectively. TaTME was associated with a significantly lower rate of anastomotic leak compared with LaTME (OR: 0.59; 95%CI, 0.38-0.91, $P = 0.02$). Heterogeneity among the included studies was low ($I^2: 0\%$, $P = 0.49$).

R0 resection: Nine studies (609 patients) reported R0 resection as an outcome. An R0 resection was achieved in 93.5% and 87.8% of patients in the TaTME and LaTME groups, respectively. The rate of R0 resection was significantly higher in the TaTME group (OR: 1.96; 95%CI, 1.07-3.58, $P = 0.03$). Low between-study heterogeneity was detected ($I^2: 0\%$, $P = 0.51$).

Completeness of mesorectal excision: This outcome was reported by nine studies (766 patients). The rate of completeness of mesorectal excision in the TaTME and LaTME groups were 81.4% and 74.0%, respectively. The pooled analysis did not demonstrated similar rate of completeness of mesorectal excision between two groups (OR: 1.57; 95%CI, 0.85-2.90, $P = 0.15$). There was moderate between-study heterogeneity ($I^2: 60\%$, $P = 0.01$).

Number of harvested lymph nodes: Eight studies (747 patients) reported the number of harvested lymph nodes in the TaTME and LaTME groups. The mean number of harvested lymph nodes in the TaTME was 16.1 ± 2.1 , while it was 16.3 ± 3.2 in the LaTME group. The pooled analysis demonstrated no significant difference in the number of harvested lymph nodes between two groups (MD: -0.05; 95%CI, -1.98-1.89, $P = 0.96$). The between-study heterogeneity was moderate ($I^2 = 71\%$, $P = 0.001$).

DRM: Eight studies (745 patients) reported DRM in their study groups. The mean DRM in the TaTME group was $15.8 \text{ mm} \pm 3.9 \text{ mm}$ whereas it was $17.6 \text{ mm} \pm 3.8 \text{ mm}$ in the LaTME group. The pooled analysis found no significant difference in DRM between two groups (MD: -0.94; 95%CI, -2.26-0.39, $P = 0.17$). There was low heterogeneity among the included studies ($I^2 = 0\%$, $P = 0.53$).

CRM: Six studies (465 patients) reported CRM in their study groups. The mean CRM in the TaTME group was $8.5 \text{ mm} \pm 1.2 \text{ mm}$ and it was $8.1 \text{ mm} \pm 2.9 \text{ mm}$ in the LaTME group. The pooled analysis did not identify any significant difference in CRM between two groups (MD: 1.08; 95%CI, -0.46-2.61, $P = 0.17$). There was moderate between-study heterogeneity ($I^2 = 71\%$, $P = 0.004$).

Positive CRM: Eight studies (717 patients) reported the rate of positive CRM in their study groups. The rate of positive CRM in the TaTME group was 9.0% and it was 13.3% in the LaTME group. There was no significant difference in the rate of positive CRM between two groups (OR: 0.64; 95%CI, 0.37-1.10, $P = 0.11$). Between-study heterogeneity was low ($I^2 = 0\%$, $P = 0.59$).

Procedure time: Ten studies (889 patients) reported the procedure time as an outcome. The mean procedure time in the TaTME and LaTME groups were $274.1 \text{ min} \pm 91.8 \text{ min}$ and $282.4 \text{ min} \pm 103.0 \text{ min}$, respectively. There was no significant difference in

procedure time between two groups (MD: -6.99 min; 95%CI, -25.28-11.30, $P = 0.45$). Heterogeneity among the studies was significant ($I^2 = 86\%$, $P < 0.00001$).

Conversion to open: This outcome was reported by eleven studies (923 patients). The rate of conversion to an open procedure in the TaTME group was 1.5% and it was 7.5% in the LaTME group. The conversion rate was significantly lower in the TaTME group compared to the LaTME group (OR: 0.29; 95%CI, 0.13-0.64, $P = 0.002$). There was low between-study heterogeneity ($I^2: 0\%$, $P = 0.54$).

Considering that the included study inadequately reported length of hospital stay as an outcome, we were unable to conduct an analysis on this outcome.

Sensitivity analysis

There was no change in the direction of pooled effect size when the risk ratio, or risk difference was calculated or during leave-one-out sensitivity analysis.

DISCUSSION

In view of ongoing debates regarding the best surgical approach for resection of low rectal cancer, we conducted a comprehensive systematic review and meta-analysis to evaluate comparative outcomes of transanal *vs* laparoscopic TME in management of low rectal cancer. We identified two RCTs and 10 observational studies^[13-24] enrolling 969 patients of whom 493 had TaTME and 476 patients had LaTME for low rectal tumour. The subsequent outcome synthesis showed that TaTME significantly reduced rate of postoperative complications, anastomotic leak, and conversion to open in comparison to LaTME. Moreover, TaTME resulted in significantly higher rate of R0 resection. However, no significant difference was found in intraoperative complications, completeness of mesorectal excision, harvested lymph nodes, DRM, CRM, positive CRM and procedure time between TaTME and LaTME.

The between-study heterogeneity in the analyses of intraoperative and postoperative complications, anastomotic leak, R0 resection, DRM, positive CRM, and conversion to

open were low suggesting that the reported findings with respect to these outcomes can be considered robust. Moderate heterogeneity among the included studies in the analyses of completeness of mesorectal excision, and number of harvested lymph nodes may suggest variation of reporting in the included studies on these outcomes. There was high between-study heterogeneity regarding procedure time suggesting that our findings about procedure time may be less robust.

The findings of our meta-analysis are not consistent with some of the findings of our previous meta-analysis on this topic published in 2020^[8]. The simple explanation for such disagreement is the difference in the inclusion criteria of the two studies with regards to the location of the rectal cancer. We only included low rectal cancer patients in this meta-analysis while previously we included both middle and low rectal cancer patients. In fact, as a direction for future research, in our previous meta-analysis we encouraged future studies to consider patients with low rectal cancer only when comparing TaTME and LaTME to evaluate a more realistic comparison between these two management approaches^[8]. This is indeed reassuring to observe growing evidence in the context of comparative outcomes of TaTME and LaTME in management of low rectal cancer. The appropriate patient selection in this context is of great importance as inappropriate patient selection for TaTME has been demonstrated to result in unfavourable outcomes of TaTME leading to suspension of TaTME in some countries. Wasmuth *et al*^[25] reported high rate of anastomotic leak and local recurrence associated with TaTME, the findings that led to suspicion of TaTME in Norway. However, only 5% of their included patients had low rectal tumours with the remaining patients having middle or high rectal cancers. Moreover, the study lacked a control group, hence low level of evidence.

In the current meta-analysis, we independently evaluated the baseline characteristics of the study population to assess if the patients in the TaTME and LaTME groups were comparable. We found no significant difference in age, gender, BMI, rate of neoadjuvant chemotherapy, and stage of cancer between two groups. Moreover, we demonstrated similar distance between the distal tumour and anal verge between the

TaTME and LaTME patients. This is of a cardinal importance as TaTME has been introduced to address the challenges associated with open and laparoscopic approaches in resecting very low rectal cancers, particularly in male patients with narrow pelvis^[8]. Therefore, comparability of our included populations in both groups makes our findings more robust.

We were not able to conduct any analyses on comparative functional outcomes of TaTME and LaTME considering that only two of the included studies reported such outcomes. Lelong *et al*^[17] compared functional outcomes of TaTME and LaTME and demonstrated no significant difference in urinary complications and faecal incontinence between two groups. Rubinkiewicz *et al*^[22] also investigated functional outcomes in patients undergoing TaTME and LaTME for low rectal tumours and reported no significant differences in risk of low anterior resection syndrome between two groups and its severity. The authors found comparable median Wexner score in both groups^[22]. Considering the current limited evidence in the context of functional outcomes of TaTME compared with LaTME, no definitive conclusions can be made.

Although we were not able to analyse long term oncological outcomes including disease recurrence, the findings of one of our included RCTs in this context is important. After 5 years follow-up, Denost *et al*^[18] reported no significant differences in long-term outcomes between TaTME and LaTME. Although the authors found a significant association between CRM involvement and local recurrence ($P = 0.011$), the 5-year local recurrence rate was similar between two groups (3% vs 5%, $P = 0.30$). Moreover, the authors reported similar 5-year disease-free survival between two groups (72% vs 74%, $P = 0.351$). The rate of local recurrence in the aforementioned RCT is comparable with the recurrence rate of 4% reported in a review by Deijen *et al*^[26]. Undoubtedly, future high quality randomized studies with adequate follow-up periods are required to investigate long term oncological outcomes of transanal and laparoscopic approaches to TME.

This study has a number of limitations. Only two of the considered studies were RCTs. Most of the included studies were observational studies with their inherited

selection bias. Some of the included studies had small sample sizes which might have introduced Type 2 error to our findings. We were unable to conduct independent analyses ¹⁷ on length of hospital stay, functional outcomes or long term oncological outcomes as the data provided by the included studies on such outcomes was inadequate. Finally, there was moderate risk of bias in 3 of our included studies.

CONCLUSION

Our meta-analysis demonstrated that for low rectal tumours, TaTME is associated with better clinical and short term oncological outcomes compared to LaTME. More randomised controlled trials with adequate power and high quality are required to not only confirm these findings, but also ¹³ to evaluate long term oncological and functional outcomes.

29%

SIMILARITY INDEX

PRIMARY SOURCES

1 Shahin Hajibandeh, Shahab Hajibandeh, Alexios Dosis, Mohammed Kaif Qayum, Karim Hassan, Ambareen Kausar, Thomas Satyadas. "Level 2a evidence comparing robotic versus laparoscopic left lateral hepatic sectionectomy: a meta-analysis", *Langenbeck's Archives of Surgery*, 2021 168 words — 5%

Crossref

2 Shahin Hajibandeh, Shahab Hajibandeh, Syed Soulat Raza, David Bartlett, Bobby V.M. Dasari, Robert P. Sutcliffe. "Abdominal drainage is contraindicated after uncomplicated hepatectomy: Results of a meta-analysis of randomized controlled trials", *Surgery*, 2022 108 words — 3%

Crossref

3 Shahin Hajibandeh, Hannah Burton, Philippa Glead, Shahab Hajibandeh, Teun Wilmink. "Impact of arteriovenous fistulas versus arteriovenous grafts on vascular access performance in haemodialysis patients: A systematic review and meta-analysis", *Vascular*, 2021 77 words — 2%

Crossref

4 Shafquat Zaman, Shahin Hajibandeh, Shahab Hajibandeh, Ali Yasen Y Mohamedahmed et al. "Meta - analysis of the demographic and prognostic significance of gastrointestinal symptoms in - 19 patients ", *JGH Open*, 2022 74 words — 2%

Crossref

-
- 5 Hajibandeh, Shahin, Shahab Hajibandeh, Ashutosh O. Gumber, and Chee Siong Wong. "Laparoscopy versus laparotomy for the management of penetrating abdominal trauma: A systematic review and meta-analysis", International Journal of Surgery, 2016. 59 words — 2%
Crossref
-
- 6 pure.uva.nl 44 words — 1%
Internet
-
- 7 www.frontiersin.org 37 words — 1%
Internet
-
- 8 I. Alimova, S. Chernyshov, M. Nagudov, E. Rybakov. "Comparison of oncological and functional outcomes and quality of life after transanal or laparoscopic total mesorectal excision for rectal cancer: a systematic review and meta-analysis", Techniques in Coloproctology, 2021 36 words — 1%
Crossref
-
- 9 Yanlai Sun, Huirong Xu, Zengjun Li, Jianjun Han, Wentao Song, Junwei Wang, Zhongfa Xu. "Robotic versus laparoscopic low anterior resection for rectal cancer: a meta-analysis", World Journal of Surgical Oncology, 2016 35 words — 1%
Crossref
-
- 10 Zhiyuan Wu, Wenlong Zhou, Fu Chen, Wentao Wang, Yong Feng. "Short-term Outcomes of Transanal versus Laparoscopic Total Mesorectal Excision: A Systematic Review and Meta-Analysis of Cohort Studies", Journal of Cancer, 2019 33 words — 1%
Crossref
-
- 11 Shahin Hajibandeh, Shahab Hajibandeh, Stavros A. Antoniou, Francesco Torella, George A. Antoniou. "Meta-analysis of Left Subclavian Artery Coverage With and Without Revascularization in Thoracic Endovascular Aortic 26 words — 1%

12 &NA;, . "The American Society of Colon and Rectal Surgeons :", Diseases of the Colon & Rectum, 2013. 24 words — 1%

Crossref

13 Cristopher Varela, Nam Kyu Kim. "Surgical Treatment of Low-Lying Rectal Cancer: Updates", Annals of Coloproctology, 2021 23 words — 1%

Crossref

14 academic.oup.com 22 words — 1%

Internet

15 Hong Yang, Zhendan Yao, Ming Cui, Jiadi Xing, Chenghai Zhang, Nan Zhang, Maoxing Liu, Kai Xu, Fei Tan, Xiangqian Su. "Influence of Tumor Location on Short- and Long-Term Outcomes After Laparoscopic Surgery for Rectal Cancer: A Propensity Score Matched Cohort Study", Research Square Platform LLC, 2020 19 words — 1%

Crossref Posted Content

16 www.researchgate.net 19 words — 1%

Internet

17 www.wcb.ny.gov 19 words — 1%

Internet

18 www.science.gov 17 words — < 1%

Internet

19 "Transanal Minimally Invasive Surgery (TAMIS) and Transanal Total Mesorectal Excision (taTME)", Springer Science and Business Media LLC, 2019 13 words — < 1%

Crossref

20 journals.sagepub.com

13 words — < 1%

21 link.springer.com

Internet

13 words — < 1%

22 www.jmir.org

Internet

12 words — < 1%

23 Yueh-Chen Lin, Ya-Ting Kuo, Jeng-Fu You, Yih-Jong Chern et al. "Transanal Total Mesorectal Excision (TaTME) versus Laparoscopic Total Mesorectal Excision for Lower Rectal Cancer: A Propensity Score-Matched Analysis", Cancers, 2022

Crossref

11 words — < 1%

24 www.cmaj.ca

Internet

11 words — < 1%

25 "Free papers (posters)-pt.3", Journal of Gastrointestinal Surgery, 200411

Crossref

10 words — < 1%

26 Ellen Kapiteijn. "Preoperative Radiotherapy Combined with Total Mesorectal Excision for Resectable Rectal Cancer", New England Journal of Medicine, 08/30/2001

Crossref

9 words — < 1%

27 Sue J. Hahn, Patricia Sylla. "Technological Advances in the Surgical Treatment of Colorectal Cancer", Surgical Oncology Clinics of North America, 2022

Crossref

9 words — < 1%

28 www.js-premier.co.th

Internet

9 words — < 1%

- 29

Internet

9 words — < 1%
- 30

Ze Li, Jingkun Xiao, Yujie Hou, Xingwei Zhang, Haiqing Jie, Huashan Liu, Lei Ruan, Ziwei Zeng, Liang Kang. "Transanal versus Laparoscopic Total Mesorectal Excision in Male Patients with Low Tumor Location after Neoadjuvant Therapy: A Propensity Score-Matched Cohort Study", Gastroenterology Research and Practice, 2022
Crossref

8 words — < 1%
- 31

cyberleninka.org
Internet

8 words — < 1%
- 32

Loyola-Sanchez, A.. "Efficacy of ultrasound therapy for the management of knee osteoarthritis: a systematic review with meta-analysis", Osteoarthritis and Cartilage, 201009
Crossref

7 words — < 1%
- 33

Simillis, Constantinos, Roel Hompes, Marta Penna, Shahnawaz Rasheed, and Paris P Tekkis. "A systematic review of transanal total mesorectal excision. Is this the future of rectal cancer surgery?", Colorectal Disease, 2015.
Crossref

7 words — < 1%