

73769\_Auto\_Edited - check.docx

**Name of Journal:** *World Journal of Gastrointestinal Surgery*

**Manuscript NO:** 73769

**Manuscript Type:** LETTER TO THE EDITOR

**Imaging of acute appendicitis-advances**

Imaging of acute appendicitis-advances

Sonay Aydın, Erdal Karavas, Düzgün Can Şenbil

## **Abstract**

Abstract: We read with interest the review by Teng *et al*, who summarized the current approach to the diagnosis and treatment of acute appendicitis (AA). Also, the article summarizes the clinical scoring systems very effectively <sup>[1]</sup>. In one of the previous studies conducted by our research group, we showed that the use of Alvarado score, ultrasound and C-Reactive Protein values in combination provides a safe confirmation or exclusion of the diagnosis of acute appendicitis<sup>[2]</sup>. Computed tomography(CT) is particularly sensitive in detecting periappendiceal abscess, peritonitis, and gangrenous changes <sup>[1]</sup>. CT is not a good diagnostic tool in pediatric patients because of the ionizing radiation it produces. US is a valuable diagnostic tool to differentiate acute appendicitis from lymphoid hyperplasia. Presence of fluid collection in the periappendiceal and lamina propria thickness less than 1 mm are the most effective parameters in differentiating appendicitis from lymphoid hyperplasia. Although acute appendicitis is the most common cause of surgical acute abdomen, it remains an important diagnostic and clinical challenge. By combining clinical scoring systems, laboratory data and appropriate imaging methods, diagnostic accuracy and adherence to treatment can be increased. Lymphoid hyperplasia and perforated appendicitis present significant diagnostic challenges in children. Additional US findings are increasingly defined to differentiate acute appendicitis from these conditions.

**Key Words:** acute appendicitis; inflammation; acute abdomen; perforation

Aydın S, Karavas E, Şenbil DC. Imaging of acute appendicitis-advances. *World J Gastrointest Surg* 2022; In press

**Core Tip:** Core tip: Despite the fact that acute appendicitis is the most common cause of acute abdomen, it remains a diagnostic and clinical challenge today. When the US, Alvarado scoring, and CRP are used in conjunction to diagnose acute appendicitis, the diagnosis can be safely confirmed or ruled out. CT scans are extremely sensitive in

detecting complications from acute appendicitis. CT scans are especially effective at detecting periappendix abscesses, peritonitis, and gangrenous changes. Because of the ionizing radiation it emits, CT is not a good diagnostic tool in pediatric patients. In pediatric patients, US should be the preferred method.

## **TO THE EDITOR**

### **TO THE EDITOR**

We read with interest the review by Teng *et al*, who summarized the current approach to the diagnosis and treatment of acute appendicitis (AA). Also, the article summarizes the clinical scoring systems very effectively [1].

In one of the published studies of our research group, we have shown that <sup>1</sup> using Alvarado score, ultrasound (US) and C- reactive protein (CRP) levels in combination enables <sup>2</sup> to confirm or rule out acute appendicitis safely [2]. Alvarado scoring system is one of the most commonly used methods [1]. Even though the scoring system contains series of laboratory parameters, it do not contain CRP levels. Rather than using Alvarado system or US alone, combining these methods with CRP levels will increase diagnostic accuracy.

Teng et. al stated that computerized tomography (CT) scans have a well-established role in evaluating AA-related complications. CT is especially sensitive for detecting periappendiceal abscess, peritonitis, and gangrenous changes [1] (Figure 1). Pediatric patients are more likely to develop perforated appendicitis. Imaging is critical in diagnosing perforated appendicitis; clinical differentiation can be challenging, especially in younger children. CT is not a good diagnostic tool in pediatric patients due to the ionizing radiation it produces. According to our results, US can also be used as an effective diagnostic tool for the detection of pediatric perforated appendicitis cases. The most valuable US parameters are the <sup>2</sup> detection of loculated fluid in the periappendiceal area and fluid collection in all abdominal recesses. When these parameters are combined with CRP levels, diagnostic performance can be improved [3].

Teng *et al* emphasized that AA occurs when the appendiceal orifice is obstructed (for example, by lymphoid hyperplasia or fecaliths), resulting in inflammation. We have demonstrated that, in addition to causing AA, lymphoid hyperplasia can serve as a significant mimicker of acute appendicitis by forming an incompressible appendix larger than 6 mm in diameter, particularly in pediatric patients. US is a valuable diagnostic tool for differentiating AA from lymphoid hyperplasia. The presence of periappendiceal fluid collection and a lamina propria thickness of less than 1 mm are the most effective parameters for differentiating appendicitis from lymphoid hyperplasia [4] (Figure 2).

Portal vein can be affected from appendiceal inflammation and thrombosis might occur [1]. In addition to complications, according to our data, portal vein hemodynamic changes can help to confirm AA diagnosis in children. In equivocal cases, detecting an increase in portal vein diameter and/or flow velocity may corroborate other clinical signs of acute appendicitis [5].

To summarize, AA remains a significant diagnostic and clinical challenge despite being the most common cause of surgical acute abdomen. By combining clinical scoring systems, laboratory data, and appropriate imaging methods, diagnostic accuracy and treatment adherence can be increased. Lymphoid hyperplasia and perforated appendicitis present significant diagnostic challenges in children. Additional US findings are increasingly being defined for the purpose of distinguishing AA from these entities.

## ORIGINALITY REPORT

---

4%

SIMILARITY INDEX

---

## PRIMARY SOURCES

---

- 1

Sonay Aydin, Erdem Fatihoglu, Hayri Ramadan, Bilge S. Akhan, Enver N Koseoglu. "Alvarado Score, Ultrasound, and CRP: How to Combine Them for the Most Accurate Acute Appendicitis Diagnosis", Iranian Journal of Radiology, 2016

Crossref

18 words — 2%
- 2

dergipark.org.tr

Internet

14 words — 2%

---

EXCLUDE QUOTES ON

EXCLUDE MATCHES

< 12 WORDS

EXCLUDE BIBLIOGRAPHY ON