

85410_Auto_Edited.docx

Name of Journal: *World Journal of Gastrointestinal Surgery*

Manuscript NO: 85410

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Gene polymorphisms associated with sudden decreases in heart rate during extensive peritoneal lavage with distilled water after gastrectomy

SNPs associated with HRV after EIPL

Shuang Yao, Yan Yuan, Jun Zhang, Yang Yu, Guang-Hua Luo

Abstract

BACKGROUND

Our previous study found that the telomerase-associated protein 1 (*TEP1*, rs938886 and rs1713449) and homo sapiens RecQ like helicase 5 (*RECQL5*, rs820196) single nucleotide polymorphisms (SNPs) were associated with changes in heart rate (HR) $\geq 30\%$ during peritoneal lavage with distilled water after gastrectomy. This study established a single tube method for detecting these three SNPs using two-dimensional (2D) polymerase chain reaction (PCR), and investigated whether SNP-SNP and SNP-environment interactions increase the risk of high heart rate variability (HRV).

AIM

To investigate whether genotypes, genetic patterns, SNP-SNP and SNP-environment interactions were associated with HRV.

METHODS

2D PCR was used to establish a single-tube method to detect *TEP1* rs938886 and rs1713449 and *RECQL5* rs820196, and the results were compared with those of Sanger

sequencing. After adjusting for confounders such as age, sex, smoking, hypertension, and thyroid dysfunction, a nonconditional logistic regression model was used to assess the associations between the genotypes and the genetic patterns (codominant, dominant, overdominant, recessive, and additive) of the three SNPs and a risk $\geq 15\%$ or $\geq 30\%$ of a sudden drop in HR during postoperative peritoneal lavage in patients with gastric cancer. Gene-gene and gene-environment interactions were analyzed using generalized multifactor dimensionality reduction.

RESULTS

The coincidence rate between the 2D PCR and sequencing was 100%. When the HRV cutoff value was 15%, the patients with the *RECQL5* (rs820196) TC genotype had a higher risk of high HRV than those who had the TT genotype [odds ratio (OR)=2.834; 95% confidence interval (CI), 1.05-3.70; $P=0.045$]. Under the codominant and overdominant models, the TC genotype of *RECQL5* (rs820196) was associated with a higher risk of HR decrease relative to the TT and TT+CC genotypes ($P=0.0314$ and 0.0163 , respectively). When the HRV cutoff value was 30%, subjects carrying the GC-TC genotypes of rs938886 and rs820196 showed a higher HRV risk when compared with the GG-TT genotype carriers ($P=0.011$). In the three-factor model of rs938886, rs820196, and rs1713449, patients carrying the GC-TC-CT genotype had a higher risk of HRV compared with the wild-type GG-TT-CC carriers ($P=0.01$). For rs820196, nonsmokers with the TC genotype had a higher HRV risk compared with nonsmokers carrying the TT genotype ($P=0.044$). When the HRV cutoff value was 15%, patients carrying the TT-TT and the TC-CT genotypes of rs820196 and rs1713449 showed a higher HRV risk when compared with TT-CC genotype carriers ($P=0.035$ and 0.012 , respectively). Patients carrying the GC-CT-TC genotypes of rs938886, rs1713449, and rs820196 showed a higher HRV risk compared with GG-CC-TT genotype carriers ($P=0.02$). When the HRV cutoff value was 15%, the best-fitting models for the interactions between the SNPs and the environment were the rs820196-smoking ($P=0.022$) and rs820196-hypertension ($P=0.043$) models. Consistent with the results of the previous

grouping, for rs820196, the TC genotype nonsmokers had a higher HRV risk compared with nonsmokers carrying the TT genotype ($P=0.01$).

CONCLUSION

The polymorphism of the *RECQL5* and *TEP1* genes were associated with HRV during peritoneal lavage with distilled water after gastrectomy.

Key Words: Homo sapiens RecQ like helicase 5; Telomerase-associated protein 1; Polymorphism; Peritoneal lavage; Heart rate variability; Two-dimensional polymerase chain reaction.

Yao S, Yuan Y, Zhang J, Yu Y, Luo GH. Gene polymorphisms associated with sudden decreases in heart rate during extensive peritoneal lavage with distilled water after gastrectomy. *World J Gastrointest Surg* 2023; In press

Core Tip: Our previous study found that peritoneal lavage with distilled water may cause a sudden decrease in heart rate (HR) in some patients during clinical gastrectomy, which was related to telomerase-associated protein 1 (*TEP1*) and homo sapiens RecQ like helicase 5 (*RECQL5*) gene polymorphisms. Here, instead of Sanger sequencing, we developed a single-tube method using two-dimensional polymerase chain reaction to genotyping SNPs quickly and economically. We also investigated whether genotypes, genetic patterns and the interaction effects of SNP-SNP and SNP-environment were associated with a risk of high HR variability. This study helps clinicians to better assess the perioperative risk of patients undergoing gastrectomy.

INTRODUCTION

Gastric cancer currently ranks fifth in incidence and fourth in mortality worldwide^[1]. Peritoneal metastasis is the most frequent pattern of postoperative recurrence in patients with gastric cancer. In order to reduce abdominal metastasis and planting

caused by surgery, distilled water or normal saline are normally used to wash the peritoneum after radical distal gastrectomy. The Chinese SEIPLUS trial reported that, for Extensive Intraoperative Peritoneal Lavage (EIPL), a new prophylactic strategy for the prevention of the peritoneal metastasis of locally advanced gastric cancer, the short-term postoperative complication rate following surgery alone (17%) was significantly higher than that following combined surgery and multiple warm physiological saline lavages (11.1%)^[2]. However, they found that addition of EIPL with saline did not improve the 3-year survival rate of advanced gastric cancer patients compared with surgery alone^[3]. A latest randomized study evaluated the efficacy and long-term outcome of advanced gastric cancer patients with EIPL^[4]. It was found that EIPL can reduce the possibility of perioperative complications including ileus and abdominal abscess. The overall survival curve and recurrence-free survival curve were better in the EIPL group. A multicenter study found that EIPL with saline after surgery did not provide a survival benefit compared with surgery alone and should not be recommended for patients undergoing curative gastrectomy for gastric cancer^[5]. Distilled water can induce osmotic lysis in cancer cells. Takemoto *et al.*^[6] demonstrated the cytocidal effects of distilled-water-induced hypotonic shock on colorectal cancer cells, which supported the use of peritoneal lavage with distilled water to remove and kill colorectal cells during surgery. Recently, a randomized trial was performed to assess the survival impact of extensive peritoneal lavage using distilled water or saline at high volumes after pancreatic resection for pancreatic ductal adenocarcinoma, which suggested that lavage with distilled water or saline could become standard practice during surgery for pancreatic cancer if it proves to be beneficial to the long-term prognosis of the patient^[7].

Our previous study found that peritoneal lavage with distilled water may have caused a sudden decrease in heart rate in some patients during clinical gastrectomy. To investigate whether gene polymorphisms are associated with high heart rate variability (HRV), we genotyped 194 patients who underwent distal gastrectomy and identified three single nucleotide polymorphisms (SNPs) (*TEP1* rs938886 and rs1713449 and

RECQL5 rs820196) associated with a risk of high HRV using whole-exome sequencing (WES)^[8]. In this study, two-dimensional (2D) polymerase chain reaction (PCR) was used to establish a single-tube method to detect these three SNPs in 192 patients (two were excluded due to incomplete clinical data). Single-gene analysis may not be appropriate for the further study of complex traits because the main effect of a single locus may be too limited for observation. Therefore, SNP-SNP and SNP-environment interactions among the three variants from the two selected genes were tested using a generalized multifactor dimensionality reduction (GMDR) approach.

MATERIALS AND METHODS

Study Population

A total of 192 patients (137 males and 55 females) scheduled to undergo distal gastrectomy were enrolled. Patients were diagnosed according to clinical and pathological data, and those with a history of other cancers or arrhythmia were excluded. Participant demographic data (age, sex, hypertension, diabetes, thyroid dysfunction, and smoking status) were collected using a standard clinical information questionnaire. Evaluation of the genomic differences of these gastric cancer patients was conducted in a previous study^[8]. Patients were divided into two groups according to the changes in HR (using 30% and 15% as cutoffs). Change in HR = (ultimate HR before lavage - HR after lavage)/HR before lavage × 100%. This study received ethical approval from the institutional review board of the Third Affiliated Hospital of Soochow University and was conducted in accordance with the Helsinki Declaration on human medical research.

DNA Extraction and SNP Genotyping

Standard techniques were adopted for the collection of venous blood samples from the participants. Whole blood was stored at -80°C for subsequent SNP analysis. Genomic DNA was extracted using the TIANamp Blood DNA Kit (Tiangen Biotech, Beijing, China). According to the principles of 2D PCR and Amplification-Refractory Mutation System (ARMS) PCR, the mutation sites of the TEP1 (rs1713449), TEP1 (rs938886), and

RECQL5 (rs820196) genes were designed at the 3' ends of the specific forward primers. The respective specific reverse primers were designed according to the DNA sequence using the PrimerPremier5.0 software, wherein the primer for *TEP1* (rs1713449) was designed according to the reverse complementary strand. Three fluorescein amidite (FAM) tags and three hexachloro-fluorescein (HEX) tags were used to label the wild-type and mutant forward primers (respectively) of *TEP1* (rs1713449), *TEP1* (rs938886), and *RECQL5* (rs820196). The sequences of the primers and probes are listed in Supplementary Table 1, which were synthesized by Sangon Biotech (Shanghai, China). The optimized 2D multiplex PCR system included 2 μ L genomic DNA, 2.5 μ L 10 \times Immobuffer (Bioline, London, UK), 0.75 μ L 25 mmol/L MgCl₂ (Bioline), 0.7 μ L 10 mmol/L dNTPs (Takara Bio, Shiga, Japan), 0.5 μ L 5 U/ μ L IMMOLASE DNA polymerase (Bioline), 0.4 μ L each probe (10 μ M), 0.1 μ L each forward primer (10 μ M; 0.2 μ L for *TEP1* rs938886), 0.6 μ L reverse primer (10 μ M), and deionized water, to make a total volume of 25 μ L. The PCRs were carried out with the following thermal cycling conditions: initial denaturation of DNA at 95°C for 10 min; amplification for five cycles at 95°C for 20 s and 60°C for 15 s; and 35 cycles at 95°C for 20 s, 72°C for 1 s, and 60°C for 15 s. The fluorescence acquisition commenced with heating at 95°C for 15 s and 30°C for 4 min; the temperature was gradually increased from 30°C to 70°C with a ramp rate of 0.1°C/s, and the fluorescence signal was acquired continuously. The final step was cooling at 40°C for 30 s. The results of 2D PCR were compared with Sanger sequencing.

Statistical analysis

Statistical analyses were performed using SAS version 9.4 (Cary, NC, USA), and the nominal *P* value ≤ 0.05 was considered the significance threshold. Normally and non-normally distributed continuous variables were compared using Student's *t* test and Mann-Whitney *U* test, respectively, and the variables were expressed as mean \pm standard deviation. The genotype and allelic frequency distributions of polymorphisms between two groups were compared using the χ^2 test and the Hardy-Weinberg equilibrium. Associations between polymorphisms and HR change were assessed by calculating odds ratios (ORs) with 95% confidence intervals (CIs) using logistic

regression analysis adjusted for age, sex, hypertension, thyroid dysfunction, and smoking status. The pairwise linkage disequilibrium (LD) and frequency of haplotypes were calculated with Haploview 4.2. Generalized multifactor dimensionality reduction was used to analyze the SNP-SNP and SNP-environment interactions. $P < 0.05$ was considered statistically significant.

RESULTS

Baseline characteristics

We recruited 192 patients with gastric cancer. The basic characteristics are presented in Tables 1 and 2. The HRV cutoff values used were 30% and 15%. There were no significant differences in the baseline demography, smoking status, hypertension, diabetes, and thyroid dysfunction between the two groups ($P > 0.05$).

Genotyping by 2D PCR

Figure 1 shows the melting curves of six alleles of the three genes detected by 2D PCR. Six melting valleys were clearly identified (Figure 1A) and six different alleles were intuitively typed. The genotypes of *TEP1* (rs1713449), *TEP1* (rs938886), and *RECQL5* (rs820196) in 190 gastric cancer patients investigated by 2D PCR were completely consistent with those determined by Sanger sequencing (the remaining two cases were not validated because of sample loss, Figure 1A and 1B).

Genotypes and allele dissemination of SNPs

Genotypes and allele frequencies of *TEP1* (rs938886), *TEP1* (rs1713449) and *RECQL5* (rs820196) in patients with a 30% or 15% decrease in HR as the cutoff points were compared. When the patients were grouped according to the 30% decrease in HR cutoff, there were no significant differences in the distribution of heterozygous and homozygous mutant and wild-type genotypes (Table 3, $P > 0.05$). However, when the patients were grouped using the 15% decrease in HR cutoff, the distribution of the *RECQL5* (rs820196) genotype frequencies was significantly different between the two groups. For *RECQL5* (rs820196), compared with individuals with the TT genotype, patients with the TC genotype had a significantly increased risk of adverse HR decline,

with an OR of 1.97 (95%CI, 1.05-3.70), which implies that *RECQL5* (rs820196) was associated with HRV in the allelic models (Table 4, $P<0.05$).

Associations between SNPs and HRV risk under different inheritance model

We next applied five genetic models (codominant, dominant, overdominant, recessive, and additive) to further analyze the relationships between the three SNPs and HRV. No significant differences were found between the patients in the $HRV\geq 30\%$ group and the control group for the studied SNPs in all the genetic models ($P>0.05$, Table 5). However, when the patients were grouped according to the 15% HRV cutoff, *RECQL5* (rs820196) was associated with a higher risk of HR change in the codominant and overdominant models. In the codominant model, the TC genotype was associated with a higher risk relative to the TT genotype (OR=2.0; 95%CI, 1.06-3.76; $P=0.031$). In the overdominant model, the TC genotype was associated with a higher risk relative to the TT and CC genotypes (OR=2.10; 95%CI, 1.15-3.84; $P=0.016$, Table 6).

LD and haplotype analysis

The results of the pairwise LD analysis of the two *TEP1* SNPs are presented in Figure 2. We detected a haplotype block with a strong LD between the rs938886 and rs1713449 SNPs. The D' value was 1, indicating that these two SNPs tend to be co-inherited. The haplotype analysis indicated that, regardless of whether the patients were grouped according to the 30% or 15% HRV cutoff, the rs938886-C/rs1713449-T and rs938886-G/rs1713449-T haplotypes were not associated with HRV risk (Tables 7 and 8, $P>0.05$).

SNP-SNP and SNP-environment interaction analysis

The GMDR approach was used to evaluate the effect of SNP-SNP interactions among the three *TEP1* and *RECQL5* SNPs. The results obtained from the GMDR analysis of the one- to three-locus models in the patients with $HRV\geq 30\%$ are summarized in Table 9. The one-locus model of *RECQL5* rs820196 was the best model of SNP-SNP interaction, recording the highest cross-validation consistency (CVC) of 10/10 and a testing accuracy of 0.6127 (P value based on 1000 permutations, $P=0.032$).

In order to obtain an OR with a 95%CI for the joint effects of SNP-SNP interactions on HRV, we also conducted interaction analyses of the GMDR models using logistic

regression. After adjusting for the factors of age, gender, smoking, hypertension, and thyroid function by multivariate logistic regression analysis, the combination of the rs93886 and rs820196 SNPs (GC-TC) was correlated with a higher risk of decreased HR (Figure 3). Table 10 describes the results generated from the GMDR method for the two-way and three-way gene-gene interaction analyses using covariate adjustment. Compared with the wild type, individuals with the GC-TC (rs93886 and rs820196, respectively) genotype and the GC-TC-CT (rs93886, rs820196, and rs1713449, respectively) genotype showed a significantly increased HRV risk (OR=6.16 and 6.27 with $P=0.01$).

Table 11 describes the results generated with the GMDR method for the SNP-environment interaction analysis using covariate adjustment. The rs820196-smoking two-factor model was found to exhibit a statistically significant association with $HRV \geq 30\%$ ($P=0.045$) (Figure 4).

After adjusting for age, gender, hypertension, and thyroid function, logistics regression showed that the nonsmokers with the rs820196 TC genotype had a significantly higher risk of HRV than the nonsmokers with the TT genotype ($P=0.04$, Table 12).

The results obtained from the GMDR analysis of the one- to three-locus models in the patients with $HRV \geq 15\%$ are summarized in Table 13 and Figure 5. The one-locus model of *RECQL5* rs820196 was the best model of SNP-SNP interaction, recording the highest CVC of 10/10 and a testing accuracy of 0.5920 ($P=0.039$).

Table 14 summarizes the results obtained from the GMDR analysis for the two- to three-loci models of gene-gene interactions. Compared with the TT-CC (rs820196-rs1713449) genotype, the TT-TT and TC-CT genotypes were associated with a higher risk of $HRV \geq 15\%$ ($P=0.04$ and 0.01 , respectively). Compared with the GG-CC-TT (rs938886-rs1713449-rs820196) genotype, the GC-CT-TC genotype showed a significantly increased HRV risk ($P=0.02$).

Table 15 and Figure 6 describe the results generated with the GMDR method for the SNP-environment interaction analysis using covariate adjustment. The rs820196-

smoking and rs820196-hypertension two-factor models were found to exhibit a statistically significant association with HRV $\geq 15\%$ ($P < 0.05$).

After adjusting for age, gender, and thyroid function, logistics regression showed that the non-smokers with the rs820196 TC genotype had a significantly higher risk of HRV $\geq 15\%$ than the non-smokers with the TT genotype ($P = 0.01$, Table 16).

DISCUSSION

RECQL5 protein is involved in the regulation of transcription elongation, the DNA damage response and DNA replication^[9, 10]. Mutations in RECQ helicases are associated with the genetic disorders Bloom syndrome, Werner syndrome, and Rothmund-Thomson syndrome, which are characterized by chromosomal instability, premature aging, and a predisposition to cancer^[11-13]. *Recql5*-knockout mice are more likely to develop cancer, and human cells deficient in RECQL5 display chromosomal instability and elevated sister chromatid exchange events, similar to cells deficient in any of the other RecQ helicases^[14]. According to a recent study, RECQL5 promotes metastasis and resistance to cisplatin in non-small cell lung cancer^[15]. A large case-control study suggests that RECQL5 is a new moderate-risk breast cancer gene^[16]. RECQL5 protein overexpression in breast cancer is strongly correlated with poor prognosis and survival, and with therapeutic resistance. A small molecule targeting RECQL5 was able to preferentially kill RECQL5-expressing breast cancers and led to the efficient sensitization of cisplatin-resistant breast cancers^[17]. Philip *et al.*^[18] identified RECQL5 as a novel pharmacological target for expanding Poly (ADPRibose) Polymerase inhibitor based treatment horizon for homologous recombination-proficient cancers. Low RECQL5 expression indicates poor prognosis in gastric carcinoma and is an independent prognostic factor^[19]. To date, no study has shown that RECQL5 is associated with arrhythmias, although a variant in chromodomain helicase DNA-binding protein 4 associated with childhood idiopathic epilepsy with sinus arrhythmia has been reported^[20]. Given the cardinal role of the RECQL5 protein in genome stability^[21], one might speculate that DNA disrepair caused by RECQL5 mutations is

the probable cause of myocardial apoptosis and arrhythmia.⁷ Further investigations are required to determine the risk of sudden cardiogenic arrhythmia caused by RECQL5 mutations.

In this study, patients with gastric cancer were divided into two groups according to two different cutoffs for HRV: 30% and 15%. There were no significant differences in gender, age, smoking history, hypertension, diabetes, blood type, and thyroid function between the two groups regardless of the cutoff used, indicating that these factors had no effect on HRV. The distribution of the *RECQL5* (rs820196) genotype frequency in patients with HRV \geq 15% differed from that in the control group. The relationships between these SNPs and HRV risk were evaluated by the inheritance models. Under the co-dominant and overdominant models, the TC genotype was associated with a higher risk of HR decrease relative to the TT and TT+CC genotypes.

⁴ TEP1 is a component of the telomerase ribonucleoprotein complex, and is responsible for catalyzing the addition of new synthetic telomere sequences to chromosome ends^[22]. Genetic variations¹¹ in telomere-associated pathway genes might affect telomere length and chromosomal stability, and subsequently disease susceptibility. In this study, the LD analysis showed that *TEP1* rs938886 and *TEP1* rs1713449 had a strong linkage relationship. Unexpectedly, there were no differences among the three haplotypes ($P>0.05$).

³ GMDR is a novel and powerful statistical tool for detecting and modeling epistasis. It is a non-parametric and model-free alternative to logistic regression for detecting and characterizing non-linear interactions among discrete genetic and environmental attributes^[23]. It is mainly used to detect gene–gene and gene–environment interactions underlying complex traits in epidemiological and genetic research. In this study, after adjusting for a multitude of covariates, we found that the best one-factor model was rs820196, whether the HRV cutoff was 30% or 15%. Logistic regression analysis was performed for better risk assessment. When the HRV cutoff value was 30%, there was a significant gene–gene interaction between rs938886 and rs820196. The subjects carrying the GC-TC genotypes of rs938886 and rs820196 showed a higher HRV risk when

compared with the GG-TT genotype carriers. In the three-factor model of rs938886, rs820196, and rs1713449, the patients carrying the GC-TC-CT genotype had a higher risk of HRV compared with the GG-TT-CC wild-type genotype. We also found a potential gene-environment interaction between rs820196 and smoking, such that the nonsmokers carrying the TC genotype of rs820196 had a higher HRV risk compared with the nonsmokers carrying the TT genotype. When the HRV cutoff was 15%, patients carrying the TT-TT and TC-CT genotypes of rs820196 and rs1713449 showed a higher HRV risk when compared with the TT-CC genotype carriers. Patients carrying the GC-CT-TC genotypes of rs938886, rs1713449, and rs820196 showed a higher HRV risk when compared with the GG-CC-TT genotype carriers. When the HRV cutoff was 15%, the best-fitting models for SNP-environment interactions were rs820196-smoking and rs820196-hypertension. Consistent with the results of the previous grouping, the non-smokers carrying the TC genotype of rs820196 had a higher HRV risk compared with the non-smokers carrying the TT genotype.

Although several positive associations were observed, some limitations of this study should be considered. First, the sample size was small, with only 192 patients being enrolled. Second, all the participants were recruited from the same hospital; hence, inherent selection bias was unavoidable. Third, was the prognosis of patients with high HRV poorer than that of those with low HRV? Did the patients with high HRV ever have arrhythmias in daily life? These questions need to be investigated. Given these limitations, further studies with larger sample sizes and more comprehensive clinical information will be required to confirm our findings.

This study on *RECQL5* and *TEP1* genetic polymorphisms may help uncover the underlying mechanisms of arrhythmia phenotypic variation. The 2D PCR used in this study helped to screen the three SNP loci more quickly and economically. In the future, when performing tumor resection and peritoneal lavage with distilled water, we suggest anesthesiologists assess the risk of sudden HR drop based on the genetic polymorphisms of *RECQL5* (rs820196) and *TEP1* (rs938886 and rs1713449), and medical history. If patients are at high risk and the baseline HR is <40 beats/min, vasopressors

such as norepinephrine, epinephrine, dopamine, and phenylephrine will be recommended before surgery. During the perioperative period, all patients are routinely monitored for arterial blood pressure by electrocardiography. An anesthetic machine is used to support breathing and monitor end-expiratory carbon dioxide partial pressure. Once the HR decreases by 30% or <40 beats/min after lavage, vasopressors should be used immediately. If cardiac arrest occurs, cardiac compression should be performed immediately, so that the heartbeat pause time is strictly limited to 1 min. Extensive peritoneal lavage with warm distilled water is widely used in surgery for breast cancer, lung cancer and gastrointestinal cancer. The purpose of this study was to screen high-risk groups through the SNP detection of high-risk genes, and focus on improving safety during the perioperative period.

CONCLUSION

³In conclusion, our results showed, for the first time, that polymorphisms of the *RECQL5* and *TEP1* genes were associated with sudden decreases in HR during abdominal lavage in patients with gastric cancer. Non-smokers carrying the TC genotype of rs820196 and the GC-CT-TC genotype carriers of rs938886, rs1713449 and rs820196 were found to have a higher HRV risk.

ARTICLE HIGHLIGHTS

Research background

Peritoneal lavage with distilled water to kill residual tumor cells is a routine procedure in gastrectomy, but this procedure often causes a sudden decrease in heart rate (HR) in some patients.

Research motivation

To investigate whether there are differences in genetic background between patients with discordant HR changes and help clinicians to better assess the perioperative risk of patients undergoing gastrectomy.

Research objectives

To investigate whether genotypes, genetic patterns, single nucleotide polymorphism (SNP)–SNP and SNP–environment interactions were associated with HRV.

13

Research methods

A total of 192 patients who underwent distal gastrectomy were divided into two groups according to changes in HR (using 30% and 15% as cutoffs). Two-dimensional polymerase chain reaction was used to establish a single-tube method to detect *TEP1* rs938886 and rs1713449 and *RECQL5* rs820196. Genotypes, genetic patterns and the interaction of SNP–SNP and SNP–environment were analyzed by non-conditional logistic regression model and generalized multifactor dimensionality reduction.

Research results

The polymorphism of the *RECQL5* gene (rs820196) was associated with a sudden decrease in HR during abdominal lavage in patients with gastric cancer. Rs820196–smoking and rs820196–hypertension were associated with HRV $\geq 15\%$. Nonsmokers carrying the TC genotype of rs820196 and patients carrying the GC–CT–TC genotype of rs938886, rs1713449 and rs820196 had higher HRV risk.

Research conclusions

The polymorphisms of *RECQL5* (TC genotype of rs820196) and *TEP1* (GC–CT genotype of rs938886 and rs1713449) genes were associated with HRV.

Research perspectives

HRV risk assessment in patients who are about to undergo peritoneal lavage is helpful for perioperative safety. This cost-effective SNP screening method can be extended to other patients undergoing tumor resection (such as breast cancer, lung cancer and other gastrointestinal cancer) and multicenter studies.

ACKNOWLEDGEMENTS

The authors thank Professor Yueping Shen from Department of Epidemiology and Biostatistics, Medical College of Soochow University for providing data statistical analysis support.

14%

SIMILARITY INDEX

PRIMARY SOURCES

- | | | |
|----------|--|-----------------------|
| 1 | www.researchgate.net
<small>Internet</small> | 127 words — 3% |
| <hr/> | | |
| 2 | www.dovepress.com
<small>Internet</small> | 100 words — 2% |
| <hr/> | | |
| 3 | Yan Wang, Fei Yu, Songcheng Yu, Dongdong Zhang, Jun Wang, Han Han, Hualei Sun, Yuan Xue, Yue Ba, Chongjian Wang, Wenjie Li. "Triangular relationship between CYP2R1 gene polymorphism, serum 25(OH)D 3 levels and T2DM in a Chinese rural population", Gene, 2018
<small>Crossref</small> | 97 words — 2% |
| <hr/> | | |
| 4 | pubmed.ncbi.nlm.nih.gov
<small>Internet</small> | 53 words — 1% |
| <hr/> | | |
| 5 | jamanetwork.com
<small>Internet</small> | 50 words — 1% |
| <hr/> | | |
| 6 | Rui Luo, Xingyuan Li, Shasha Qin, Zhenzhou Luo, Xi Luo, Peiwu Hu, Zhisheng Liu, Xiangling Feng, Xingli Li. " Impact of SNP-SNP interaction among , , and on methadone dosage requirement in Han Chinese patients ", Pharmacogenomics, 2017
<small>Crossref</small> | 38 words — 1% |
| <hr/> | | |
| 7 | www.ncbi.nlm.nih.gov
<small>Internet</small> | 33 words — 1% |

8 www.semanticscholar.org

Internet

21 words — < 1%

9 link.springer.com

Internet

15 words — < 1%

10 www.wjgnet.com

Internet

15 words — < 1%

11 zero.sci-hub.se

Internet

15 words — < 1%

12 Dao-Di Peng, Wei Xie, Zai-Xin Yu. "Impact of interaction between CYP1A1 genetic polymorphisms and smoking on coronary artery disease in the Han of China", Clinical and Experimental Hypertension, 2017

Crossref

14 words — < 1%

13 H. Tsujioka, S. Matsuoka, S. Sorano, H. Yamamoto, S. Nakamura, A. Tohyama, T. Fukami, M. Goto, R. Matsuoka, F. Eguchi. "Follow-up study of symptomatic submucous fibroids after hysteroscopic myomectomy", Clinical and Experimental Obstetrics & Gynecology, 2017

Crossref

13 words — < 1%

14 Jun-hong Li, Ning Wu, Hong-mei Yang, Hong-bing Tang, Dong-ping Bao, Jun-min Ji. "Interaction between STAT3 gene polymorphisms and smoking on Crohn's disease susceptibility: a case-control study in a Chinese Han population", Inflammation Research, 2016

Crossref

13 words — < 1%

15 Wan-Ju Annabelle Lee, Wei-Pang Chung, Shih-Chieh Shao, Edward Chia-Cheng Lai, Yi-Chen Chen, Chung-Han Ho. "Risk of retinal vein occlusion in colorectal cancer patients receiving anti-vascular endothelial

13 words — < 1%

16

Kenichi Takemoto, Atsushi Shiozaki, Daisuke Ichikawa, Shuhei Komatsu et al. "Evaluation of the efficacy of peritoneal lavage with distilled water in colorectal cancer surgery: in vitro and in vivo study", Journal of Gastroenterology, 2014

Crossref

12 words — < 1%

17

www.science.gov

Internet

12 words — < 1%