



Name of Journal: *World Journal of Gastrointestinal Surgery*

ESPS manuscript NO: 35571

Manuscript type: Original Article

Clinical Practice Study

Risk factors for pancreatic fistula following pancreaticoduodenectomy: A retrospective study in a Thai tertiary center

Narongsak Rungsakulkij, Somkit Mingphruedhi, Pongsatorn Tangtawee, Chonlada Krutsri, Paramin Muangkaew, Wikran Suragul, Penampai Tannaphai, Suraida Aeesoa

Abstract

AIM

The aim of this study was to analyze the risk factors of postoperative pancreatic fistula following pancreaticoduodenectomy in a Thai tertiary care center.

Match Overview

- | Rank | Source | Words | Similarity |
|------|---|----------|------------|
| 1 | Crossref Nobuhisa Akamatsu. "Risk factors for postoperative pancreatic fistula after pancreaticoduodenectomy: the significance | 42 words | 1% |
| 2 | Crossref Ji-Ye Chen, Jian Feng, Xian-Qiang Wang, Shou-Wang Cai, Jia-Hong Dong, Yong-Liang Chen. "Risk scoring system an | 29 words | 1% |
| 3 | Crossref Manabu Kawai. "How Do We Predict the Clinically Relevant Pancreatic Fistula After Pancreaticoduodenectomy?—A ... | 22 words | 1% |
| 4 | Internet crawled on 31-Jul-2016
www.jultrasoundmed.org | 21 words | 1% |
| 5 | Crossref N. A. Gaag. "Preoperative Biliary Drainage in Patients with Obstructive Jaundice: History and Current Status", Journal | 16 words | <1% |
| 6 | Crossref C. Ansore. "Structured intraoperative assessment of panc | 11 words | <1% |

[全部](#) [图片](#) [视频](#) [新闻](#) [地图](#) [图书](#)

找到约 17,500 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[Google 学术: Risk factors for pancreatic fistula following pancreaticoduodenectomy: A retrospective study in a Thai tertiary center](#)

[... of pancreatoduodenectomy in a tertiary cancer center ...](#) - Shrikhande - 被引用次数: 28

[... who are candidates for pancreatoduodenectomy or ...](#) - Iacono - 被引用次数: 95

[... Two-Center Case-Case-Control Study of Risk Factors ...](#) - Logan - 被引用次数: 10

[Risk factors for postoperative pancreatic fistula: Analysis of 539 ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5016380/>

14 Sep 2016 ... The risk factors for pancreatic fistula were analyzed by univariate analysis and ... In this study, we conducted a retrospective analysis of 539 successive Pancreatic fistula after pancreaticoduodenectomy is a common and serious Care must be taken in patients with any of these risk factors, and an ...

[Risk factors affecting pancreatic fistulas after ... - NCBI - NIH](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2773861/>

7 Dec 2008 ... METHODS: We retrospectively reviewed 172 consecutive patients ...

CONCLUSION: Our study demonstrated that pancreatic fistulas are ...

Pancreaticoduodenectomy (PD) is a commonly performed surgical ... The aim of this study was to analyze the independent risk factors for pancreatic fistulas after PD.

[\[PDF\] Pancreatic fistula following pancreaticoduodenectomy](#)

https://openarchive.ki.se/xmlui/bitstream/.../Thesis_Ansorge.pdf?...

indicators of pancreatic fistula after pancreaticoduodenectomy, ... 'intermediate' (softer PC or smaller PDD, 26%) or 'low' (no risk factors, 2%). ... In study III, a prospective observational cohort study on 195 consecutive PD patients ... high-volume centers with surgical and oncologic expertise may show promise to improve.

[\[PDF\] ผลลัพธ์ระหว่างการผ่าตัด Pancreaticoduodenectomy ในโรงพยาบาล](#)

www.kmutt.ac.th/jif/public_html/Download/Format.../48.pdf

Material and Method: Retrospective study of data from 26 patients who ... study, preoperative and operative factors had no significant effect on morbidity rate. ... achieved in high volume centers, mortality rates in low ... fistula, Pancreatic leakage rate of 10-20% after PD have



找到约 67 条结果 (用时0.14秒)

时间不限

2017以来

2016以来

2013以来

自定义范围...

按相关性排序

按日期排序

不限语言

中文网页

简体中文网页

☒ 包括专利

☐ 包含引用

☒ 创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

Evolution of **pancreatoduodenectomy** in a **tertiary** cancer center in India: improved results from service reconfiguration

SV Shrikhande, [SG Barreto](#), BA Somashekar... - *Pancreatology*, 2013 - Elsevier

... Despite this progress, PD carries a significant morbidity (38–52%) and **risk** of mortality (0–9%) even in the developed ... hemorrhage (PPH) are as per our previous publications and by the International **Study** Group of **Pancreatic** Surgery (ISGPS ... **Pancreas**, 13 (9%), 27 (13%), 37 (25 ...

☆ 99 被引用次数: 29 相关文章 所有 8 个版本

Role of preoperative biliary drainage in jaundiced patients who are candidates for **pancreatoduodenectomy** or hepatic resection: highlights and drawbacks

C Iacono, [A Ruzzenente](#), T Campagnaro... - *Annals of ...*, 2013 - journals.lww.com

Introduction: In this review of the literature, we analyze the indications for preoperative drainage.

☆ 99 被引用次数: 100 相关文章 所有 5 个版本

[HTML] ... -Spectrum β -Lactamase-Producing Enterobacteriaceae Infections in Children: A Two-**Center** Case–Case–Control **Study** of **Risk Factors** and Outcomes in ...

LK Logan, LA Meltzer, JB McAuley... - *Journal of the ...*, 2014 - academic.oup.com

... **Study** period prevalence remained stable (1.7 ... The incidence of pediatric infection due to ESBL-positive Enterobacteriaceae was stable at 2 large **tertiary-care** medical **centers** over a 4 ... **Risk factors** for infection due to ESBL-producing bacteria include neurologic medical conditions ...

☆ 99 被引用次数: 10 相关文章 所有 3 个版本

[HTML] Advances in diagnosis, treatment and palliation of **pancreatic** carcinoma: 1990-2010

C Sharma, KM Eltaib, PD Banerjee... - *World Journal of ...*, 2011 - wjg.com

[全部](#)[图片](#)[新闻](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 21,500 条结果（用时 0.68 秒）

小提示： 仅限搜索简体中文结果。您可以在设置中指定搜索语言

Google 学术: Risk factors for pancreatic fistula following pancreaticoduodenectomy: A retrospective study in a Thai tertiary center

... of pancreatoduodenectomy in a tertiary cancer center ... - Shrikhande - 被引用次数: 29

... who are candidates for pancreatoduodenectomy or ... - lacono - 被引用次数: 100

... Two-Center Case–Case–Control Study of Risk Factors ... - Logan - 被引用次数: 10

Pancreatic fistula following pancreaticoduodenectomy: clinical ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/19461951> - 翻译此页

作者: CM Schmidt - 2009 - 被引用次数: 74 - 相关文章

2009年5月18日 - Pancreatic fistula following pancreaticoduodenectomy: clinical predictors and ... This study seeks to identify clinical factors which may predict pancreatic fistula (PF) ... We performed a retrospective analysis of a clinical database at an academic tertiary care hospital with a high volume of pancreatic surgery.

Risk factors for postoperative pancreatic fistula: Analysis of 539 ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: BY Hu - 2016 - 被引用次数: 1 - 相关文章

2016年9月14日 - The risk factors for pancreatic fistula were analyzed by univariate analysis ... In this study, we conducted a retrospective analysis of 539 The dilemma of pancreatic fistula after pancreaticoduodenectomy has not yet been resolved[27]. Care must be taken in patients with any of these risk factors, and an ...

缺少字词: thai tertiary

Predictive risk factors for clinically relevant pancreatic fistula analyzed ...