

国内版

国际版

Clinical outcomes of patients with duodenal adenocarcinoma ;



All

Images

Videos

关闭取词

27,500 Results

Any time ▾

## Outcomes and Treatment Options for Duodenal ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6097725>

Jun 26, 2018 · Duodenal adenocarcinoma (DA) is a rare tumor representing approximately 0.5% of all gastrointestinal tumors, although it accounts for more than 50% of small bowel adenocarcinomas (SBAs), and in many studies outcomes for all SBAs are grouped together.<sup>1–3</sup> Due to the location of the duodenum in the gastrointestinal tract, molecular similarities, and possibly similar phenotypic carcinogenesis, DA is often compared with colorectal cancer ...

## Clinicopathologic Features and Prognosis of Duodenal ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4378233>

For the diagnosis of AA, we used the criteria proposed by Adsay et al. and defined AA as a tumor epicentered in the lumen or walls of the distal ends of the common bile duct and/or pancreatic duct; at the "papilla of Vater", or at the duodenal surface of the papilla . Patients with AA or PDA, who received any form of neoadjuvant therapy, were excluded from this retrospective study.

Cited by: 13

Author: Maryam Zenali, Michael James Overman,...

Publish Year: 2013

## Long-term outcomes in patients with duodenal ...

[https://www.researchgate.net/publication/236664503\\_Long-term\\_outcomes\\_in\\_patients\\_with...](https://www.researchgate.net/publication/236664503_Long-term_outcomes_in_patients_with...)

Segmental duodenal resection for patients with duodenal adenocarcinoma is associated with acceptable postoperative morbidity and long-term survival. The ...

## Clinical relevance of pancreatobiliary and intestinal ...

<https://www.sciencedirect.com/science/article/pii/S1424390319300201>

Clinical relevance of pancreatobiliary and intestinal subtypes of ampullary and duodenal adenocarcinoma: Pattern of recurrence, chemotherapy, and survival after pancreatoduodenectomy

Author: Inger Marie Bowitz Lothe, Inger Marie ...

Publish Year: 2019

## Intestinal-type of differentiation predicts favourable ...

<https://bmccancer.biomedcentral.com/articles/10.1186/1471-2407-13-428> ▾

Sep 22, 2013 · In univariate analysis, intestinal subtype was associated with better survival in ampullary, pancreatic ductal and duodenal adenocarcinoma. The intestinal type of pancreatic ductal adenocarcinoma was not associated with intraductal papillary mucinous neoplasm and could not be reliably diagnosed by immunohistochemical staining pattern alone

**Name of Journal:** *World Journal of Gastrointestinal Oncology*

**Manuscript NO:** 52190

**Manuscript Type:** ORIGINAL ARTICLE

*Observational study*

**Clinical outcomes of patients with duodenal adenocarcinoma and intestinal-type papilla of Vater adenocarcinoma**

Meijer LL *et al.* Clinical outcome of DA

Laura L Meijer, Marin Strijker, Jacob K de Bakker, Jurgen GJ Toennaer, Barbara M Zonderhuis, Hans J van der Vliet, Hanneke Wilmink, Joanne Verheij, Freek Daams,

### Match Overview

1	<b>Internet</b> 38 words crawled on 15-Dec-2019 <a href="http://link.springer.com">link.springer.com</a>	1%
2	<b>Internet</b> 18 words crawled on 11-Mar-2015 <a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a>	<1%
3	<b>Internet</b> 14 words crawled on 03-Jan-2020 <a href="http://hbsn.amegroups.com">hbsn.amegroups.com</a>	<1%
4	<b>Internet</b> 13 words crawled on 10-Oct-2019 <a href="http://www.karger.com">www.karger.com</a>	<1%



国内版 国际版

Clinical outcomes of patients with duodenal adenocarcinoma and intestinal



登录



网页 图片 视频 学术 词典 地图

试用 Microsoft Edge >

检测到您输入了英文，试试切换到国际版？搜英文结果更丰富更准确 >

7,310 条结果 时间不限 ▾

### Long-term outcomes in patients with duodenal ... 翻译此页

The aims of the study were to evaluate the rate of POPF and long-term outcomes for patients with duodenal adenoma and adenocarcinoma who underwent curative resection. ... papilla and ampulla of ... [https://www.researchgate.net/publication/236664503\\_Long-term\\_outcomes\\_in\\_patients\\_with...](https://www.researchgate.net/publication/236664503_Long-term_outcomes_in_patients_with...)

### Clinical relevance of pancreatobiliary and intestinal ... 翻译此页

Author: Inger Marie Bowitz Lothe, Inger Marie ... Publish Year: 2019  
Introduction. Periapillary adenocarcinomas originate from the duodenum, ampulla of Vater, bile duct or the pancreas, and pancreatoduodenectomy is the standard procedure for surgical treatment of these tumours []. Following curative surgery, ampullary and duodenal carcinomas have a better prognosis than pancreatic ductal adenocarcinoma and distal bile duct cancer [2,3].  
<https://www.sciencedirect.com/science/article/pii/S1424390319300201>

### Prognostic impact of immunohistochemical ... 翻译此页

Cited by: 7 Author: Sung Pil Yun, Hyung Il Seo  
Publish Year: 2015 位置: 8600 Rockville Pike, Bethesda, MD  
2015-11-24 · Prognostic impact of immunohistochemical expression of CK7 and CK20 in curatively resected ampulla of Vater cancer ... Intestinal type adenocarcinomas of the duodenal papilla originate from the intestinal mucosa covering the papilla and evolve through an adenoma-dysplasia-carcinoma sequence. ... Kim S-J, Roh M-S, Jeong J-S, et al. The ...  
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4657248>

### Intestinal-type and Pancreatobiliary-type ... 翻译此页

Cited by: 76 Author: Arne Westgaard, Arne Westgaard, Ewa P.



27,000 Results

Any time ▾

## Outcomes and Treatment Options for Duodenal ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6097725>

Jun 26, 2018 · Duodenal adenocarcinoma (DA) is a rare tumor representing approximately 0.5% of all gastrointestinal tumors, although it accounts for more than 50% of small bowel adenocarcinomas (SBAs), and in many studies outcomes for all SBAs are grouped together.<sup>1–3</sup> Due to the location of the duodenum in the gastrointestinal tract, molecular similarities, and possibly similar phenotypic carcinogenesis, DA is often compared with colorectal cancer ...

Cited by: 6

Author: Laura L. Meijer, Anna J. Alberga, Jacob K...

Publish Year: 2018

## Exceptional Response to Nanoparticle Albumin-Bound ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4748796>

In conclusion, chemotherapy with nanoparticle albumin-bound paclitaxel plus gemcitabine was an effective and well-tolerated palliative treatment in a patient with metastatic adenocarcinoma of the ampulla of Vater. It is worthwhile to reassess the histopathological diagnosis to clarify the reasons for an ineffective treatment strategy.

## Long-term outcomes in patients with duodenal ...

[https://www.researchgate.net/publication/236664503\\_Long-term\\_outcomes\\_in\\_patients\\_with...](https://www.researchgate.net/publication/236664503_Long-term_outcomes_in_patients_with...)

Long-term outcomes in patients with duodenal adenocarcinoma. Background: Because of the rarity of duodenal adenocarcinoma, little is known regarding its natural history or prognostic factors for survival. We therefore evaluated surgical treatment, and prognostic factors for survival in patients with duodenal adenocarcinoma.

## Different Clinicopathologic Findings in Two Histologic ...

[onlinelibrary.wiley.com/doi/10.1111/j.1349-7006.1994.tb02077.x/full](https://onlinelibrary.wiley.com/doi/10.1111/j.1349-7006.1994.tb02077.x/full)

Aug 24, 2005 · The aim of this study was to investigate the differences between the clinicopathological findings in two histologic types of carcinoma of the papilla of Vater. We histologically classified carcinoma of the papilla into two types: 1) an intestinal type that resembles tubular adenocarcinoma of the stomach or colon, and 2) a pancreaticobiliary type that is characterized by papillary ...

Cited by: 240

Author: Wataru Kimura, Noriaki Futakawa, Seiich...

Publish Year: 1994

## Histomolecular Phenotypes and Outcome in ...

<https://ascopubs.org/doi/full/10.1200/JCO.2012.46.8868>