

1 / 20

Name of Journal: *World Journal of Gastrointestinal Oncology*

Manuscript NO: 55153

Manuscript Type: ORIGINAL ARTICLE

Basic Study

Clinical pathological characteristics and prognosis of 132 cases of rectal neuroendocrine tumors

Yong-Jun Yu, Yu-Wei Li, Yang Shi, Zhao Zhang, Min-Ying Zheng, Shi-Wu Zhang

Abstract

BACKGROUND

Neuroendocrine tumors (NETs) frequently occur in the gastrointestinal tract, lung, and pancreas, and the rectum and appendix are the sites with the highest incidence. Epidemiology statistics show that an estimated 8000 people every year in the United

Match Overview

1	Internet 42 words crawled on 17-May-2020 www.wjgnet.com	1%
2	Internet 35 words crawled on 11-Sep-2017 digitalcommons.wustl.edu	1%
3	Crossref 13 words Shunsuke Tsukamoto. "Clinicopathological characteristics and prognosis of rectal well-differentiated neuroendocrine tumors",	<1%
4	Internet 12 words crawled on 09-Jun-2018 mdanderson.influent.utsystem.edu	<1%
5	Crossref 12 words Yihebalu Chi, Feng Du, Hong Zhao, Jin-Wan Wang, Jian-Qiang Cai. "Characteristics and long-term prognosis of patients with"	<1%

国内版

国际版

Clinical pathological characteristics and prognosis of 132 cases of



ALL

IMAGES

VIDEOS

151,000 Results

Any time ▼

Characteristics and long-term prognosis of patients with ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4239514>

Nov 21, 2014 · INTRODUCTION. **Rectal neuroendocrine tumors** are rare with an incidence of about 3.08 per 100000 person-years[], but there has been a slow upward trend in their incidence since the 1990s[2,3]. They are considered to be a type of tumor with indolent biological behavior and a relatively favorable **prognosis** []. In 2003, Modlin et al[] reported that the 5-year **survival** rate was 88.3%.

Cited by: 14

Author: Yihebaoli Chi, Feng Du, Hong Zhao, Jin-W...

Publish Year: 2014

The Clinical Pathological Characteristics and Prognostic ...

<https://www.enets.org/the-clinical-pathological...> ▼

The **Clinical Pathological Characteristics** and Prognostic Factors of **Rectal Neuroendocrine Tumors**- A Retrospective Analysis Based on Multi Center Data Abstract #1583 Introduction: The **pathological characteristics and prognosis** factors of rectum **neuroendocrine** tumor still have a lot of uncertainty.

Diagnosis and Management of Rectal Neuroendocrine Tumors

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5719921>

Nov 30, 2017 · **CLINICAL FEATURES**. Macroscopically, **rectal** NETs appear as smooth, round, polypoid lesions with normal overlying mucosa that are usually found within 5–10 cm from the anal verge (Fig. 1) []. According to SEER data, there is a slight male predominance and mean age at **diagnosis** is 57 years [1,4]. **Rectal** NETs have a higher incidence and prevalence in both African American and Asian ...

Search To

Turn off Hover



Clinical pathological characteristics and prognosis of 132 cases of re



ALL

IMAGES

VIDEOS

MAPS

NEWS

SHOPPING

159,000 Results

Any time ▾

[Characteristics and long-term prognosis of patients with ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4239514>

Nov 21, 2014 · INTRODUCTION. **Rectal neuroendocrine tumors** are rare with an incidence of about 3.08 per 100000 person-years[], but there has been a slow upward trend in their incidence since the 1990s[2,3]. They are considered to be a type of tumor with indolent biological behavior and a relatively favorable **prognosis** []. In 2003, Modlin et al[] reported that the 5-year **survival** rate was 88.3%.

Cited by: 14

Author: Yihebaoli Chi, Feng Du, Hong Zhao, Jin-W...

Publish Year: 2014

[The Clinical Pathological Characteristics and Prognostic ...](#)

<https://www.enets.org/the-clinical-pathological...> ▾

The **Clinical Pathological Characteristics** and Prognostic Factors of **Rectal Neuroendocrine Tumors**- A Retrospective Analysis Based on Multi Center Data Abstract #1583 Introduction: The **pathological characteristics and prognosis** factors of rectum **neuroendocrine** tumor still have a lot of uncertainty.

[Diagnosis and Management of Rectal Neuroendocrine Tumors](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5719921>

Nov 30, 2017 · **CLINICAL FEATURES**. Macroscopically, **rectal** NETs appear as smooth, round, polypoid lesions with normal overlying mucosa that are usually found within 5–10 cm from the anal verge (Fig. 1) []. According to SEER data, there is a slight male predominance and mean age at **diagnosis** is 57 years [1,4]. **Rectal** NETs have a higher incidence and prevalence in both African American and Asian ...

Cited by: 9

Author: Shreya Chablaney, Zachary A. Zator, Nik...

Publish Year: 2017

[Rectal neuroendocrine tumor with uncommon metastatic ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4753174>

Feb 15, 2016 · Core tip: **Rectal neuroendocrine tumors** consist approximately 5%-14% of all **neuroendocrine** neoplasms in Europe. Distant metastases from **rectal neuroendocrine tumors** are not very common. Herein we describe a **case** of a **rectal neuroendocrine** tumor with an uncommon natural