

Feedback

[PDF] Liver tumors in children with metabolic disorders

https://tp.amegroups.com/article/viewFile/8124/8905

Can a child with NAFLD develop liver cancer?

children dying before the age of 2 years, and a chronic form consisting of chronic liver disease, renal tubular dysfunction, and hypophosphatemia with rickets (25). The neurologic crises (resulting from δ-aminolevulinic acid toxicity) consist of painful paresthesias and autonomic signs similar to those seen in acute porphyria (17).

Pediatric Hepatoblastoma (Liver Cancer) | Children's ...

https://www.chp.edu/.../hepatoblastoma-liver-cancer •

Hepatoblastoma is the most common liver cancer of childhood, although it affects only 2 to 3 people in a million. It affects children during the first three years of life and usually presents as an abdominal mass which causes pain and discomfet. A large percentage of those tumors again in

Quotes Included Bibliography Included

13%

Name of Journal: World Journal of Gastrointestinal Oncology

Manuscript NO: 65218

Manuscript Type: MINIREVIEWS

Liver tumors in children with chronic liver diseases


Sintusek P et al. Pediatric liver tumors

Palittiya Sintusek, Teerasak Phewplung, Anapat Sanpawat, Yong Poovorawan

Abstract

Liver tumors are rare in children, albeit the incidence may increase in some circumstances and particularly in chronic liver diseases. Most liver tumors consequent to chronic liver diseases are malignant hepatocellular carcinoma. Other liver tumors include hepatoblastoma, focal nodular hyperplasia, adenoma, pseudotumor, and nodular regenerative hyperplasia. Screening of suspected cases is beneficial. Imaging and surrogate markers of alpha-fetoprotein are used initially as noninvasive tools for

		2 2 2	SIMILAR
	Ma	tch Overview	
	4		
	1	Internet 59 words crawled on 15-Feb-2017 www.wjgnet.com	2%
	2	Crossref 44 words Yueh-Wei Liu. "Hepatic pseudotumor in long-standing b ary atresia patients undergoing liver transplantation", Live	1%
	3	Crossref 43 words Nicolaj Laugesen, Christian Nolsoe, Jacob Rosenberg. "C linical Applications of Contrast-Enhanced Ultrasound in th	1%
	4	Crossref 36 words Zimmermann, Arthur. "Liver Tumors of Childhood", Practic al Hepatic Pathology A Diagnostic Approach, 2011.	1%
	5	Crossref 33 words Milton J. Finegold. "Liver tumors: Pediatric population", Li ver Transplantation, 11/2008	1%
	6	Internet 30 words crawled on 20-Aug-2020 www.intechopen.com	1%
	7	Internet 25 words crawled on 24-Jun-2020 journals.lww.com	190
) 4 9	8	Crossref 23 words Bioulac-Sage, Paulette, Hervé Laumonier, and Charle Paul Balabaud. "Benign Hepatocellular Tumors", Practical	1%



Chronic Liver Diseases in Children: Clinical Profile and ...

https://pubmed.ncbi.nlm.nih.gov/26393179

国内版

Another aim being to assess the prognosis and to know the outcome and the effects of treatment in chronic liver diseases in paediatric age group. Materials and methods: It was a prospective study, included the clinical profile of Chronic Liver Diseases (CLD) in children and the histopathological correlation. A total of 55 children were ...

Cited by: 10 Author: Sachin Devidas Dhole, Archana S Kher, Rad...

Publish Year: 2015

Liver Conditions in Children | Johns Hopkins Medicine

https://www.hopkinsmedicine.org/health/conditions... •

Cirrhosis or Chronic Liver Failure. Cirrhosis or chronic liver failure can be caused by a number of the liver and biliary diseases listed in Clinical Services or sometimes appear without known cause. Our approach to pediatric patients with this problem is to start with a thorough history and physical examination and nutritional assessment. We also have a frank discussion with the patient and family about what is involved in the evaluation for a life-saving liver transplant. Liver Cyst or Tumor

Definition & Facts of NAFLD & NASH in Children | NIDDK

https://www.niddk.nih.gov/health-information/liver-disease/nafld-nash-children/...
NAFLD is the most common cause of chronic liver disease in children in the United States. Researchers estimate that close to 10 percent of U.S. children ages 2 to 19 have NAFLD. 1 NAFLD has become more

common in children in recent decades, in part because childhood obesity has become more common.

Pediatric Liver Tumors: Overview, Benign Tumors, Malignant ...

https://emedicine.medscape.com/article/940516-overview

Sep 15, 2016 - Most children with liver tumors present with abdominal distension, a palpable abdominal mass, or both. Anemia, thrombocytopenia, and leukocytosis are sometimes present. Children ...

Neurologic Manifestations of Chronic Liver Disease and ...

https://pubmed.ncbi.nlm.nih.gov/25908229

Neurologic damage in chronic liver disease and liver cirrhosis seems to be multifactorial primarily attributable to the following: brain accumulation of ammonia, manganese, and lactate; altered permeability of the blood-brain barrier; recruitment of monocytes after microglial activation; and neuroinflammation, that is, direct effects of circulating systemic proinflammatory cytokines such as tumor ...

Cited by: 17 Author: Binit Sureka, Kalpana Bansal, Yashwant Pa...

Publish Year: 2015




Feedback

Liver Disease in Children | SSM Health

https://www.ssmhealth.com/.../liver-diseases •

When the liver becomes severely damaged it can result in acute or chronic liver disease, or liver failure both of which are life-threatening conditions. Our team of SLUCare Physician Group hepatologists (liver doctors) at SSM Health Cardinal Glennon Children's Hospital are leaders in the care of children with liver disease.

Some results are removed in response to a notice of local law requirement. For more information, please see here.



A highly contagious liver infection caused by hepatitis A virus that cause

A condition in which amyloid proteins build up on organs like heart, kidney

and liver. Amyloid is an abnormal protein, and its deposit on the organ can

Data from Focus Medica. Reviewed by a panel of doctors.Learn more

Chat with Bing

Search Tools

Suggest an edit

Amyloidosis

eventually damage them.

Turn off Hover Translation (关闭取词)

inflammation and affect liver function.

Consult a medical professional for advice