

74638\_Auto\_EditedC.docx

**Name of Journal:** *World Journal of Gastrointestinal Oncology*

**Manuscript NO:** 74638

**Manuscript Type:** ORIGINAL ARTICLE

### *Retrospective Study*

**Clinicopathological differences, risk factors and prognostic scores for western patients with intestinal and diffuse-type gastric cancer**

**Díaz del Arco C *et al.* Clinicopathological prognostic scores for Laurén subtypes**

**Cristina Díaz del Arco, Lourdes Estrada Muñoz, Luis Ortega Medina, Elena Molina Roldán, Ma Ángeles Cerón Nieto, Soledad García Gómez de las Heras, M. Jesús Fernández Aceñero**

### **Abstract**

#### **BACKGROUND**

In the molecular era, the Laurén system is still a cost-effective and widely implemented classification for gastric cancer (GC), and it has been recently associated with clinical, histological, and molecular features of these tumors. Despite recent advances in the understanding of the molecular biology of GC, there is a need to develop new prognostic tools for patient stratification in clinical practice. Thus, the identification of easily available prognostic factors in patients with intestinal and diffuse-type tumors can significantly improve risk assessment and patient stratification in GC.

#### **AIM**

To identify clinicopathological differences, risk factors, and to develop cost-effective prognostic scores for patients with intestinal and diffuse-type GC.

#### **METHODS**

Retrospective study of all patients undergoing surgery for GC at a tertiary referral center, from 2001 to 2019. 286 cases met inclusion criteria (intestinal: 190, diffuse: 96). Clinical data and gross findings were collected. All specimens were reviewed by two independent pathologists and a detailed protocol for histologic evaluation was followed. Five tissue microarrays (TMAs) were constructed and sections of the TMA block were immunostained for HERCEPTEST, MSH2, MSH6, MLH1 and PMS2. Statistical analyses were performed, and prognostic scores were developed based on hazard ratios.

## RESULTS

Intestinal and diffuse-type GC showed different epidemiological, clinicopathological and prognostic features. Diffuse tumors were significantly associated with younger age, less symptomatology, flat morphology, deeper invasion, perineural infiltration, advanced stage at diagnosis, administration of adjuvant therapy and poorer prognosis. Intestinal lesions were fungoid or polypoid, showed necrosis, desmoplasia, microsatellite instability and HERCEPTEST positivity, and were diagnosed at earlier stages. Tumor depth, desmoplasia, macroscopic type and lymph node involvement were independently related to the Laurén subtype. Furthermore, intestinal and diffuse GC were associated with different risk factors for progression and death. Vascular invasion, perineural infiltration and growth pattern were important prognostic factors in intestinal-type GC. On the contrary, tumor size and necrosis were significant prognosticators in diffuse-type GC. Our recurrence and cancer-specific death scores for patients with intestinal and diffuse-type GC showed an excellent patient stratification into three (diffuse GC) or four (intestinal) prognostic groups.

## CONCLUSION

Our findings support that Laurén subtypes represent different clinicopathological and biological entities. The development of specific prognostic scores is a useful and cost-effective strategy to improve risk assessment in GC.

**Key Words:** Gastric cancer; Clinicopathological; Score; Prognosis; Laurén; Molecular

<sup>7</sup> Díaz del Arco C, Estrada Muñoz L, Ortega Medina L, Molina Roldán E, Cerón Nieto MÁ, García Gómez de las Heras S, Fernández Aceñero MJ. Clinicopathological differences, risk factors and prognostic scores for western <sup>16</sup> patients with intestinal and diffuse-type gastric cancer. *World J Gastrointest Oncol* 2022; In press

**Core Tip:** In the molecular era, the Laurén system is a cost-effective and widely implemented classification. The identification of easily available prognostic factors in intestinal and diffuse-type tumors may significantly improve patient stratification in gastric cancer (GC). In this study, the authors found that intestinal and diffuse-type GC show different epidemiological, clinical and prognostic features. Laurén subtypes were also associated with different risk factors for tumor progression and death. Finally, separate clinicopathological scores for patients with intestinal and diffuse-type GC showed an excellent prognostic stratification. The development of specific prognostic scores is a useful, cost-effective strategy to improve risk assessment in GC.

## INTRODUCTION

Gastric cancer (GC) is an aggressive tumor which is usually diagnosed at advanced stages in western countries<sup>[1,2]</sup>. It can be classified according to its location, macroscopical or microscopical features<sup>[3,4]</sup>. Although several histological-based classifications have been proposed, only the Laurén and the most recent WHO classification are currently widely used<sup>[5]</sup>. The Laurén system has been extensively adopted by clinicians and pathologists since its publication in 1965, and it can be easily evaluated in conventional paraffin-embedded hematoxylin-eosin stained slides<sup>[6]</sup>. This classification divides GC into intestinal, diffuse and mixed types, depending on the tumor architecture, growth pattern and cell morphology. Intestinal-type GC is composed of glandular structures accompanied by papillary or solid components. On the other hand, diffuse-type GC is composed of loosely attached cells, growing as small clusters or scattered cells with an infiltrative pattern. This classification has been variably associated with clinicopathological features<sup>[7,8]</sup>. Intestinal tumors occur more frequently in older men, and they are related to *Helicobacter pylori* (*H. pylori*) infection and environmental factors. Furthermore, most studies have identified Laurén subtype as an independent prognosticator in GC<sup>[9-11]</sup>.

Recent technological advances have allowed us to improve the understanding of the molecular biology of GC<sup>[12]</sup>. In 2014, The Cancer Genome Atlas (TCGA) Research Network defined four molecular subtypes of GC: tumors positive for Epstein-Barr virus, microsatellite unstable tumors (MSI), genomically stable tumors (GS) and tumors with chromosomal instability (CIN)<sup>[13]</sup>. Laurén classification has also been correlated with these molecular groups<sup>[14,15]</sup>.

The only curative treatment for GC is surgery, and localized tumors are treated by total or subtotal gastrectomy. However, patient prognosis is poor, with estimated 5-year survival rates lower than 30%<sup>[16]</sup>. There is an urgent need to identify potential therapeutic targets and prognostic factors in GC, in order to improve patient subclassification and response to therapy. It has been suggested that diffuse GC may benefit from broader surgical margins, extended lymphadenectomy and hyperthermic

intraperitoneal chemotherapy (HIPEC) for peritoneal metastasis<sup>[17,18]</sup>. Interestingly, diffuse GC may be more resistant to standard chemotherapeutic regimens than intestinal GC, and several investigators have recommended the use of the Laurén classification in clinical trials<sup>[19]</sup>.

In summary, in the era of molecular medicine, the Laurén system is a cost-effective and widely implemented classification, which has been associated with clinical, pathological, prognostic and molecular features. Thus, Laurén subtypes can be considered distinct entities that differ in their histologic, biological, and clinical behaviour, and the identification of easily available prognostic factors in patients with intestinal and diffuse-type tumors may significantly improve risk assessment and patient stratification in GC.

In this study, our objectives were to: (1) Assess the clinicopathological differences between Laurén subtypes; (2) Identify and compare the clinicopathological risk factors for recurrence and cancer-specific death of patients with intestinal and diffuse-type GC; and (3) Develop specific cost-effective prognostic scores for overall survival (OS) and disease-free survival (DFS) for patients with intestinal and diffuse GC. As far as we know, no other study has developed specific clinicopathological prognostic scores for patients with intestinal and diffuse-type GC.

## **MATERIALS AND METHODS**

### ***Study design***

This is a retrospective study including all patients undergoing surgery for GC at a tertiary referral hospital in Madrid (Spain), from 2001 to 2019. All tumors were treated by total or subtotal gastrectomy with D1 or D2 Lymphadenectomy. Clinical records were reviewed and demographic, clinical and follow-up information was retrieved, including age at diagnosis, sex, symptoms (local and systemic), smoking and drinking habits, treatment, tumor recurrence and cause of death. Gross findings (tumor size, tumor location, macroscopic type) were collected from the database of the Department



of Surgical Pathology (PatWin). The study<sup>8</sup> was reviewed and approved by the institutional review board of the hospital.

### *Histopathological study*

All tumors<sup>1</sup> were formalin-fixed and embedded in paraffin. Slides were reviewed by two independent pathologists<sup>1</sup>, following a detailed protocol for histologic evaluation. Discordant cases were conjointly reviewed. Main microscopic features were assessed, including tumor type (Laurén classification), histologic grade, presence of signet-ring cells (independently of the percentage of signet-ring cells), tumor budding, perineural infiltration, lymphovascular invasion, growth pattern (expansive or infiltrative), desmoplasia, necrosis, surgical margins, tumor depth (T stage), number of lymph node<sup>4</sup> (LN) dissected, number of metastatic LN and lymph node ratio (LNR).<sup>1</sup> The LNR was defined as the ratio between the number of metastatic LN and the total number of LN retrieved from the resection specimen. LNR was treated as a quantitative variable. All cases<sup>12</sup> were staged according to the 8th edition of the American Joint Committee on Cancer tumor, node, metastasis (TNM) classification of GC<sup>[20]</sup>.

### *Immunohistochemical study*

Five tissue microarrays (TMAs) including a subgroup of cases from the GC cohort were assembled for immunohistochemical (IHC) analysis. TMAs were<sup>8</sup> constructed using the MTA-1 tissue arrayer (Beecher Instruments, Sun Prairie, United States), and contained 2 cores per case (diameter: 1 mm). Representative areas were pre-selected by a pathologist<sup>2</sup>, corresponding to the center and leading edge of each tumor. Cores were punched out from the donor block and transferred into a TMA. 2 µm sections were obtained from the TMA block for IHC study.<sup>6</sup> Slides were deparaffinised by incubation at 60°C for 10 min and incubated with Dako PT-Link for 20 min at 95°C in a high pH buffered solution. Holders were incubated with peroxidase blocking reagent (Dako, Denmark) to block endogenous peroxidase.<sup>2</sup> Sections were incubated for 20 min with the primary antibodies, followed by incubation with the appropriate anti-Ig horseradish

peroxidase-conjugated polymer (EnVision, Dako, Denmark) to detect anti-gen-antibody reaction. Then, biopsies were visualized with 3,3'-diaminobenzidine as a chromogen for 5 min and counterstained with haematoxylin. Sections of the TMA block were immunostained for HERCEPTEST, MSH2, MSH6, MLH1 and PMS2 (all antibodies prediluted; Dako, Denmark). Positive and negative controls were included. IHC markers were evaluated by two experienced pathologists. Staining intensity, location, and percentage of cells stained were assessed for all antibodies. For the aims of this study, HERCEPTEST was evaluated according to the CAP recommendations. MSI tumors showed complete loss of MLH1, MSH2, MSH6 and/or PMS2 IHC expression.

### *Inclusion criteria*

We reviewed all GC resected in our institution between 2001 and 2019. After data collection, patients receiving neoadjuvant therapy, metastatic tumors at diagnosis, patients with R1 or R2 resections and tumors of the mixed type were excluded from the study.

### *Statistical analysis*

All data was stored in an anonymized Excel file and analyzed using IBM SPSS statistics (V20). Qualitative variables were described as frequencies. Quantitative variables were expressed as mean  $\pm$  SD or median with range, as appropriate. For the analysis of the association between variables, we have performed  $\chi^2$  (chi)-squared test for qualitative data, and Student's *t*-test for dichotomic quantitative variables. Statistical significance was settled at a *P* value < 0.05.

Multivariate analyses were performed by Cox regression (backward stepwise method), and regression models were adjusted for potential confounders. Two models were calculated for each Laurén subtype (OS and DFS).

Prognostic scores for tumor progression and death were developed based on the hazard ratios of significant independent prognostic factors, as seen in other studies<sup>[21]</sup>.



Two prognostic scores (one for recurrence and one for cancer-specific death) were constructed for each Laurén subtype.

OS and DFS curves according to the prognostic scores were estimated by Kaplan Meier analysis, and significance was tested with the log-rank test. Receiver operating characteristic curves for cancer-specific death and progression were plotted. The area under the curve (AUC) was calculated for each prognostic score.

## RESULTS

A total of 377 GC were resected in our institution between 2001 and 2019. Final analyses included 286 patients with pure intestinal-type GC ( $n = 190$ ) and diffuse-type GC ( $n = 96$ ). Clinicopathological features of our cases are presented as Supplementary Table 1. Mean age at diagnosis was 72 years, and most patients were symptomatic (90.2%). Mean tumor size was 43 mm, and most lesions were located in the gastric antrum (56.1%) and body (34.5%). According to their macroscopic appearance, tumors were mainly fungoid (36%) or ulcerative (31.8%). Microscopically, 66.4% of GC were intestinal ( $n = 190$ ) and 33.6% were diffuse ( $n = 96$ ). 35.1%, 35.1% and 50.5% of cases showed vascular invasion, perineural infiltration and desmoplasia, respectively. IHC was performed in 172 GC (intestinal  $n = 107$ , diffuse  $n = 65$ ): 28.5% were microsatellite unstable and most cases were HERCEPTEST 0 (91.9%). Patients were diagnosed at stages I (27.2%), II (31.8%) and III (40.5%). 18% of patients received adjuvant therapy. Mean follow-up was 46.5 mo (0-208 mo). During follow-up, 36.6% of tumors recurred and 26.8% of patients died due to tumor.

### *Clinicopathological differences between Laurén subtypes*

Clinicopathological features of our cases depending on the Laurén subtype are summarized in Supplementary Table 1. Univariate analysis (Table 1) showed a significant association between Laurén subtypes and patient age, tumor depth, macroscopic type, local symptoms, necrosis, perineural infiltration, intratumoral inflammatory infiltration, desmoplasia, MSI, HERCEPTEST, T, N, LNR, adjuvant

therapy, tumor recurrence and patient death. Diffuse tumors were diagnosed at advanced stages, in younger patients with less local symptoms. They infiltrated deeper into the gastric wall, had flat morphology and higher rates of perineural infiltration. MSI was infrequent and HERCEPTEST was negative (0) in all cases. Adjuvant treatment was administered more frequently to patients with diffuse GC. Intestinal lesions were more frequently fungoid or polypoid, showed necrosis and desmoplasia, and were diagnosed at earlier stages. In respect of patient prognosis, diffuse GC was significantly associated with higher rates of tumor recurrence and cancer-specific death. DFS and OS curves according to the Laurén classification are presented as Supplementary Figures 1 and 2, respectively.

#### *Prognostic factors in intestinal and diffuse-type GC*

**Intestinal GC:** Univariate analysis is summarized in Supplementary Table 2. Tumor recurrence was significantly associated with perineural infiltration, vascular invasion, T, N, TNM stage and LNR. The relationship between tumor budding or MSI and recurrence tended to be significant ( $P = 0.057$  and  $0.084$ , respectively). Death due to GC was significantly related to infiltrative growth pattern, vascular invasion, T, N, TNM stage and LNR. Presence of necrosis and MSI approached significance ( $P = 0.088$  and  $0.096$ , respectively). Multivariate analysis is presented in Table 3: LNR, vascular invasion and T stage were independent risk factors for tumor recurrence, whereas LNR and growth pattern were independently associated with tumor death.

**Diffuse GC:** Univariate analysis is summarized in Supplementary Table 2. GC recurrence was significantly related to tumor size, T, N, TNM stage and LNR; and death due to GC was significantly associated with tumor necrosis, presence of systemic symptoms, N, TNM stage and LNR. The association between cancer-specific death and vascular invasion, T stage and tumor size tended to be significant ( $P = 0.059$ ,  $0.058$  and  $0.08$ , respectively). Multivariate analysis (Table 3) identified tumor size and LNR as

independent predictors of tumor recurrence. Necrosis and LNR were independent risk factors for death due to GC.

### *Prognostic scores for patients with intestinal and diffuse-type GC*

**Intestinal GC:** Two prognostic scores were constructed based on hazard ratios (Table 4). The recurrence score included T stage, LNR and vascular invasion; total score ranged from 0 to 9. Kaplan-Meier curves showed an excellent patient stratification into four prognostic groups (S1-S4,  $P < 0.001$ , Figure 1). Mean DFS times were 161, 129, 83 and 61 mo for S1-S4 cases, respectively. The risk score for predicting cancer-specific death included LNR and growth pattern; total score ranged from 0 to 37. Cases were divided into four categories (S1-S4). This score showed a good prognostic performance by Kaplan-Meier analysis ( $P < 0.001$ , Figure 2). Mean OS was 170, 132, 77 and 67 mo for S1-S4 patients. AUC values of the prognostic scores for recurrence and cancer-specific death were 0.745 and 0.763, respectively (Supplementary Table 3).

**Diffuse GC:** Prognostic scores for diffuse-type GC are presented in Table 5. The score for predicting tumor recurrence included tumor size and LNR; total score ranged from 0 to 120. The score for cancer-specific death included tumor necrosis and LNR; total score ranged from 0 to 10. Both prognostic scores showed an excellent risk stratification of patients into three groups (S1-S3,  $P < 0.005$ , Figures 3 and 4). Mean DFS was 93, 90 and 33 mo (S1-S3 patients) and mean OS was 145, 86 and 16 mo (S1-S3 patients). AUC values of the prognostic scores for recurrence and cancer-specific death were 0.674 and 0.710, respectively (Supplementary Table 3).

## **DISCUSSION**

The global incidence of GC has been decreasing in recent years, and this fact may be due to the detection and eradication of *H. pylori* and improvements in food preservation<sup>[22-24]</sup>. However, the relative incidence of diffuse GC is consequently increasing<sup>[25]</sup>. As previously mentioned, the Laurén system was developed in 1965 as a

“histo-clinical classification”. After this first description of intestinal and diffuse-type GC, several studies have variably associated Laurén subtypes with clinicopathological features of GC, including patient age, sex or macroscopic morphology<sup>[7,8,26]</sup>. In our series, Laurén subtypes showed significant differences in age at diagnosis, tumor depth, macroscopic type, local symptoms, necrosis, perineural infiltration, intratumoral inflammatory infiltration, desmoplasia, T, N, LNR and administration of adjuvant therapy.

In respect of GC prognosis, the relationship between Laurén subtypes and patient outcomes is still controversial. In this study, we observed that diffuse GC showed higher rates of recurrence and cancer-specific death than intestinal tumors. Furthermore, in a previous study, we identified Laurén subtype as an independent prognostic factor for both DFS and OS in a subgroup of patients with GC from our institution<sup>[27]</sup>. Most authors have found that diffuse tumors are significantly and independently related to poor prognosis<sup>[10,28,29]</sup>, but other studies have not confirmed these findings<sup>[30,31]</sup>. A recent meta-analysis including 73 publications and more than 61000 patients further confirmed the prognostic value of the Laurén classification<sup>[32]</sup>.

The huge impact of technological advances on GC diagnosis and pathogenesis has led to the development of new molecular-based classifications<sup>[33]</sup>. However, molecular studies are expensive and these classifications have not been implemented in practice. The most important systems have been published by TCGA in 2014 and the Asian Cancer Research Group (ACRG) in 2015<sup>[13,34]</sup>. TCGA defined four subtypes: tumors positive for Epstein-Barr virus, MSI tumors, GS tumors and tumors with CIN<sup>[13]</sup>. ACRG divided GC into *p53* active, *p53* inactive, mesenchymal and MSI GC<sup>[34]</sup>. Most GS and mesenchymal tumors are diffuse and most cases of MSI and CIN GC are intestinal-type tumors<sup>[35]</sup>. Intestinal tumors are associated with MSI and show higher mutation rates and more copy-number alterations than diffuse-type GC. On the other hand, diffuse GC is related to *CDH1* mutation, and approximately 9% of these tumors present MSI<sup>[36]</sup>. Recent studies have also shown that HER2 positivity is more frequently seen in intestinal-type GC<sup>[37,38]</sup>. Our results support these findings: we observed that intestinal-



type tumors are associated with higher rates of MSI (34.6% *vs* 18.5%) and HERCEPTEST positivity. 6.7%, 3.8% and 3.8% of intestinal cases were 1+, 2+ and 3+, respectively, whereas all diffuse tumors were HERCEPTEST negative (0).

Laurén classification may also play a role in patient management and response to therapy<sup>[39]</sup>. Early GC can be treated by endoscopic resection, and standard criteria include well or moderately differentiated GC confined to the mucosa, size  $\leq 20$  mm and absence of lymphatic or venous invasion<sup>[40,41]</sup>. Current expanded criteria for endoscopic submucosal dissection include the resection of high grade GC  $\leq 2$  cm in size<sup>[42]</sup>. GC in stages IB-III is treated by gastrectomy with lymphadenectomy. Diffuse tumors may benefit from more aggressive surgical options and prevention or treatment of peritoneal metastases by HIPEC<sup>[43,44]</sup>. Regarding chemotherapeutic and radiotherapeutic regimens, most treatment protocols are based on the TNM classification. However, several authors have observed that treatment response may vary depending on the Laurén subtype, and have suggested that diffuse tumors may be more resistant to standard chemotherapeutic agents than intestinal-type GC<sup>[39]</sup>. In a recent literature review, we summarized the main findings of clinical trials and comparative studies analyzing treatment regimens depending on the Laurén subtypes<sup>[11]</sup>. According to this review, studies on adjuvant therapy showed that intestinal-type GC is more chemo-sensitive than diffuse GC, and treatment response may also vary depending on geographical features. The benefit of neoadjuvant therapy seems to be limited to patients with intestinal GC.

Clinicopathological prognostic features of GC patients have been previously analyzed in the literature<sup>[45]</sup>. Some authors have studied specific prognostic factors in younger patients or proximal tumors<sup>[46,47]</sup>. However, as far as we know, no other study has evaluated the prognostic features of patients with intestinal and diffuse-type GC. Thus, in this study, our second objective was to identify clinicopathological risk factors for tumor recurrence and cancer-specific death depending on the Laurén subtype. As might be expected, the TNM system was associated with tumor recurrence and death in both GC subtypes. But several differences were observed: vascular invasion, perineural

infiltration and infiltrative growth pattern were important prognostic features in intestinal-type GC. On the contrary, tumor size and necrosis were significant prognosticators in diffuse-type GC. As for molecular features, we found that the relationship between MSI and prognosis tended to be significant only in intestinal-type tumors.

Finally, we constructed prognostic scores for predicting tumor recurrence and cancer-specific survival in patients with intestinal and diffuse-type GC. Our scores included only clinicopathological variables and can be easily calculated in clinical practice. All scores showed an excellent patient stratification into three (diffuse GC) or four (intestinal GC) prognostic groups by Kaplan-Meier analyses. Previous studies have developed predictive scores for GC patients, and most of them included nutritional and laboratory findings<sup>[41,48]</sup>. Other authors have developed molecular signatures or scores including immunohistochemical parameters<sup>[49]</sup>. Recently, Bao *et al*<sup>[50]</sup> developed a three-gene signature for prognostic prediction in diffuse-type GC. Clinicopathological prognostic scores, although easy to apply, have been less frequently published<sup>[51]</sup>. As far as we know, no other study has developed separate clinicopathological risk scores for patients with intestinal and diffuse-type GC.

### **Strengths and limitations of our study**

The results of this study should be interpreted in the context of its strengths and limitations. Strengths: This study includes patients with pure intestinal or diffuse GC treated by curative gastrectomy. Cases with neoadjuvant therapy, mixed tumors and R1-2 resections were excluded. All patients were diagnosed and treated in a western tertiary hospital. All tumors were reviewed and pathological features were independently assessed by two pathologists, following a detailed protocol. Limitations: Retrospective study. GC is not frequent in western countries, so this study includes less patients than Asian studies. IHC markers were performed in TMA sections, and they may not represent the full heterogeneity of the tumor. In an attempt to overcome this limitation, cores were selected from the center and the leading edge of each case.



Furthermore, <sup>20</sup> no significant differences were observed between the two cores of each tumor.

## **CONCLUSION**

In our series, intestinal and diffuse-type GC showed different epidemiological, clinical and prognostic features, and they were associated with different risk factors for progression and death. Our specific prognostic scores for predicting tumor recurrence and cancer-specific survival in patients with intestinal and diffuse-type GC showed an excellent patient stratification into three (diffuse GC) or four (intestinal GC) prognostic groups.

Laurén classification is a cost-effective and widely implemented tool in GC, and it has regained importance in the last years due to its correlation to the molecular groups of GC. Our findings support the notion that Laurén subtypes may represent different clinicopathological and biological entities, and the development of specific prognostic scores could be a useful and cost-effective strategy to improve risk assessment and patient stratification in GC. Our scores include clinicopathological variables easily available in practice, and patients can be stratified according to their risk without complementary tests. However, our results should be externally validated and refined in other western and eastern cohorts of patients. Thus, more studies with larger number of patients and other ethnic groups are <sup>3</sup> needed, in order to confirm the prognostic validity of the proposed prognostic scores and the current role of Laurén classification in GC.

## **ARTICLE HIGHLIGHTS**

### ***Research background***

In the molecular era, the Laurén system is still a cost-effective and widely implemented classification for gastric cancer (GC), and it has been recently associated with clinical, histological, and molecular features of these tumors. Laurén subtypes have also shown differences in response to systemic therapy.

### *Research motivation*

Despite recent advances in the understanding of the molecular biology of GC, there is a need to develop new prognostic tools for patient stratification in clinical practice. The implementation of specific scores for patients with intestinal and diffuse-type GC may significantly improve risk assessment and management of GC.

### *Research objectives*

Our aims were to: (1) evaluate the clinicopathological differences between Laurén subtypes; (2) identify specific risk factors for these subtypes; and (3) develop prognostic scores for patients with intestinal and diffuse-type GC.

### *Research methods*

This is a retrospective study of all patients undergoing surgery for GC at a tertiary referral center, from 2001 to 2019. Clinical data and gross findings were collected. Histological and immunohistochemical features were assessed by two independent pathologists, and prognostic scores were developed based on hazard ratios.

### *Research results*

In our series of western patients with GC, intestinal and diffuse-type tumors showed distinctive epidemiological, clinical and prognostic features. In addition, Laurén subtypes were associated with different risk factors for tumor progression and cancer-specific death. Our prognostic scores for predicting overall survival and disease-free survival in patients with intestinal and diffuse-type GC included clinicopathological variables that can be easily calculated in clinical practice, and showed an excellent patient stratification into three (diffuse GC) or four (intestinal GC) prognostic groups.

### *Research conclusions*

The stratification of GC patients depending on Laurén subtypes and the implementation of specific clinicopathological prognostic scores in intestinal and diffuse-type tumors can be useful for patient stratification, risk assessment and treatment selection.

### *Research perspectives*

Our prognostic scores should be externally validated in patients from both western and eastern countries, due to the geographical variation of GC. In addition, this study opens a door to the development and implementation of cost-effective and specific clinicopathological prognostic scores in patients with GC in different contexts.

22%

SIMILARITY INDEX

---

PRIMARY SOURCES

---

1 Cristina Díaz del Arco, Lourdes Estrada Muñoz, Andrés Sánchez Pernaute, Luis Ortega Medina et al. "Development of a simplified tumor-lymph node ratio classification system for patients with resected gastric cancer: A western study", Annals of Diagnostic Pathology, 2021 219 words — 5%

Crossref

2 link.springer.com 141 words — 3%

Internet

3 www.ncbi.nlm.nih.gov 113 words — 3%

Internet

4 Cristina Díaz del Arco, Lourdes Estrada Muñoz, Luis Ortega Medina, Ángela Chávez et al. "Update, validation and comparison of three different clinicopathological scores for patients with resected gastric cancer: A western experience", Annals of Diagnostic Pathology, 2020 97 words — 2%

Crossref

5 Cristina Díaz del Arco, Lourdes Estrada Muñoz, Andrés Sánchez Pernaute, Luis Ortega Medina et al. "Which Lymph Node Staging System Better Predicts Prognosis in Patients With Gastric Carcinoma? A Comparative Study Between 3 Different Lymph Node Classifications for Resected Gastric Cancer in a Western Tertiary Center", American Journal of Clinical Oncology, 2021 59 words — 1%

Crossref

6	Cristina Díaz del Arco, Lourdes Estrada Muñoz, Rodrigo Barderas Manchado, Alberto Peláez García et al. "Prognostic Role of Aryl Hydrocarbon Receptor Interacting Protein (AIP) Immunohistochemical Expression in Patients with Resected Gastric Carcinomas", Pathology & Oncology Research, 2020 Crossref	54 words — 1%
7	www.wjgnet.com Internet	51 words — 1%
8	journals.plos.org Internet	31 words — 1%
9	Hyeyoon Chang, Woon Yong Jung, Youngran Kang, Hyunjoo Lee, Aeree Kim, Han Kyeom Kim, Bong Kyung Shin, Baek-hui Kim. "Programmed death-ligand 1 expression in gastric adenocarcinoma is a poor prognostic factor in a high CD8+ tumor infiltrating lymphocytes group", Oncotarget, 2016 Crossref	28 words — 1%
10	C. de la Fouchardiere, L. Decoster, E. Samalin, C. Terret, C. Kenis, J.P. Droz, C. Coutzac, E. Smyth. "Advanced oesophago-gastric adenocarcinoma in older patients in the era of immunotherapy. A review of the literature", Cancer Treatment Reviews, 2021 Crossref	23 words — 1%
11	www.science.gov Internet	18 words — < 1%
12	www.springermedizin.de Internet	18 words — < 1%
13	Zhang, Yilong, Rajul Jain, and Min Zhu. "Recent Progress and Advances in HGF/MET-Targeted	15 words — < 1%

---

14 " Week 2020 Poster Presentations ", United European Gastroenterology Journal, 2020 13 words — < 1 %

Crossref

---

15 Cristina Díaz del Arco, Luis Ortega Medina, Lourdes Estrada Muñoz, Soledad García Gómez de las Heras et al. "Pathologic Lymph Node Staging of Gastric Cancer", American Journal of Clinical Pathology, 2021 12 words — < 1 %

Crossref

---

16 Si-Chun Ming. "Gastric carcinoma:A pathobiological classification", Cancer, 1977 12 words — < 1 %

Crossref

---

17 [www.dovepress.com](http://www.dovepress.com) 11 words — < 1 %

Internet

---

18 Mengwei Li, Yanzhen Han, Haoze Zhou, Xin Li, Chenyu Lin, Erhao Zhang, Xiaowei Chi, Jialiang Hu, Hanmei Xu. "Transmembrane protein 170B is a novel breast tumorigenesis suppressor gene that inhibits the Wnt/ $\beta$ -catenin pathway", Cell Death & Disease, 2018 10 words — < 1 %

Crossref

---

19 Ji-Hye Choi, Young-Bae Kim, Ji Mi Ahn, Min Jae Kim, Won Jung Bae, Sang-Uk Han, Hyun Goo Woo, Dakeun Lee. "Identification of genomic aberrations associated with lymph node metastasis in diffuse-type gastric cancer", Experimental & Molecular Medicine, 2018 9 words — < 1 %

Crossref

---

20 "Abstracts", Virchows Archiv, 2018 8 words — < 1 %

Crossref



---

21 Cristina Díaz del Arco, Lourdes Estrada Muñoz, Elena Molina Roldán, M<sup>a</sup> Ángeles Cerón Nieto et al. 8 words — < 1%  
"Immunohistochemical classification of gastric cancer based on new molecular biomarkers: a potential predictor of survival", Virchows Archiv, 2018  
Crossref

---

22 Seokha Yoo, Han-Byoel Lee, Wonshik Han, Dong-Young Noh, Sun-Kyung Park, Won Ho Kim, Jin-Tae Kim. " Total Intravenous Anesthesia Inhalation Anesthesia for Breast Cancer Surgery ", Anesthesiology, 2019 8 words — < 1%  
Crossref

---

23 [www.nejm.org](http://www.nejm.org) 8 words — < 1%  
Internet

---

EXCLUDE QUOTES OFF  
EXCLUDE BIBLIOGRAPHY OFF

EXCLUDE SOURCES OFF  
EXCLUDE MATCHES OFF