

81934\_Auto\_Edited-check.docx

3

**Name of Journal:** *World Journal of Gastrointestinal Oncology*

**Manuscript NO:** 81934

**Manuscript Type:** ORIGINAL ARTICLE

***Basic Study***

**Circ\_0003356 suppresses gastric cancer growth through targeting the miR-668-3p/SOCS3 axis**

Li WD *et al.* Circ\_0003356 suppresses the malignancy of GC

## Abstract

### BACKGROUND

Circular RNAs (circRNAs) have attracted extensive attention as therapeutic targets in gastric cancer (GC). Circ\_0003356 is known to be downregulated in GC tissues, but its cellular function and mechanisms remain undefined.

### AIM

To investigate the role of circ\_0003356 in GC at the molecular and cellular level.

### METHODS

Circ\_0003356, miR-668-3p, and SOCS3 expression were assessed *via* quantitative real time-polymerase chain reaction (qRT-PCR). Wound healing, EdU, CCK-8, flow cytometry and transwell assays were used to analyze the migration, proliferation, viability, apoptosis and invasion of GC cells. The subcellular localization of circ\_0003356 was monitored using fluorescence in situ hybridization. The interaction of circ\_0003356 with miR-668-3p was confirmed using RIP-qRT-PCR, RNA pull-down, and dual luciferase reporter assays. We observed protein levels of genes *via* western blot. We injected AGS cells into the upper back of mice and performed immunohistochemistry staining for examining E-cadherin, N-cadherin, Ki67, and SOCS3 expressions. TUNEL staining was performed for the assessment of apoptosis in mouse tumor tissues.

### RESULTS

Circ\_0003356 and SOCS3 expression was downregulated in GC cells, whilst miR-668-3p was upregulated. Exogenous circ\_0003356 expression and miR-668-3p silencing suppressed the migration, viability, proliferation, epithelial to mesenchymal transition (EMT) and invasion of GC cells and enhanced apoptosis. Circ\_0003356 overexpression impaired tumor growth in xenograft mice. Targeting of miR-668-3p by circ\_0003356 was confirmed through binding assays and SOCS3 was identified as a downstream target of

miR-668-3p. The impacts of circ\_0003356 on cell proliferation, apoptosis, migration, invasion and EMT were reversed by miR-668-3p up-regulation or SOCS3 down-regulation in GC cells.

## CONCLUSION

Circ\_0003356 impaired GC development through its interaction with the miR-668-3p/SOCS3 axis.

**Key Words:** Epithelial-mesenchymal transition; Circ\_0003356; Gastric cancer; Invasion; Proliferation; Migration

Li WD, Wang HT, Huang YM, Cheng BH, Xiang LJ, Zhou XH, Deng QY, Guo ZG, Yang ZF, Guan ZF, Wang Y. Circ\_0003356 suppresses gastric cancer growth through targeting the miR-668-3p/SOCS3 axis. *World J Gastrointest Oncol* 2023; In press

**Core Tip:** We observed the low level of circ\_0003356 expression in gastric cancer (GC) tissues and cells. Circ\_0003356 expression was positively related to GC patient prognosis. Exogenous circ\_0003356 overexpression and/or miR-668-3p suppression enhanced apoptosis in GC cells and suppressed GC cell proliferation, migration, invasion, and epithelial to mesenchymal transition. The overexpression of circ\_0003356 also prevented tumor growth in mice. At the mechanistic level, circ\_0003356 was found to interact with the miR-668-3p/SOCS3 axis to impair GC development. Together, we reveal new and important molecular details highlighting circ\_0003356 as a novel cancer target.

## INTRODUCTION

Gastric cancer (GC) encompasses a range of gastrointestinal malignancies<sup>[1,2]</sup>, and ranks as the 2<sup>nd</sup> main cause of global cancer-associated deaths<sup>[3,4]</sup>. Chemotherapy, radiotherapy, and surgical resection have progressed GC treatment<sup>[5,6]</sup>, but five-year

survival rates remain low<sup>[7,8]</sup>. Personalized therapy has shown promise for GC treatment<sup>[9]</sup>, but more effective anti-GC targets are required. Circular RNAs (circRNAs) regulate GC occurrence and development. The downregulation of circ\_SKA3 inhibits the colony formation, proliferation, migration, and invasion of GC cells *in vitro* and suppresses tumorigenesis *in vivo*<sup>[10]</sup>. Circ\_0005758 up-regulation can inhibit GC cell proliferation, migration and invasion, and suppress tumor growth in xenograft mice<sup>[11]</sup>. Up-regulation of circ\_0021087 impairs proliferation, epithelial to mesenchymal transition (EMT), and invasion of GC cells and overpowers tumor development in xenograft mice<sup>[12]</sup>. CircRNA analysis in GSE184882 indicates that circ\_0003356 is lowly expressed in GC tissues, but the molecular effects governing its role in GC suppression have not been investigated at the molecular level.

MicroRNAs (miRNAs) regulate the post-transcriptional expression of a number of genes<sup>[13,14]</sup> and play a key role in GC progression. The overexpression of miR-548 promotes proliferation, migration and invasion of GC cells *in vitro*<sup>[15]</sup>. Silencing of miR-762 reduces GC cell viability and facilitates apoptosis<sup>[16]</sup>. Inhibition of miR-4742-5p suppresses the proliferation, migration and invasion of GC cells<sup>[17]</sup>. Especially, miR-668-3p is oncogenic in hepatocellular carcinoma (HCC)<sup>[18]</sup> and colorectal cancer (CRC) cells<sup>[19]</sup>.

SOCS3 is a negative regulator of hormone and cytokine signaling and plays a key role in cancer development<sup>[20,21]</sup>. Available studies have highlighted the importance of SOCS3 in the manipulation of cancer development. A study from Yu *et al*<sup>[22]</sup> have revealed that the downregulation of SOCS3 can reverse the suppressive effects of miR-3173-5p inhibitor on colony formation, cell proliferation, migration, invasion and EMT in prostate cancer cells. Another study from Li *et al*<sup>[23]</sup> have reported that SOCS3 overexpression also suppresses spheroid formation and CRC cell proliferation. In particular, SOCS3 is proven to exert an important role against tumor in GC. For instance, inhibition of miR-340 suppresses cell cycle progression through enhancing the expression of SOCS3 in GC cells<sup>[24]</sup>. MiR-665 also serves as an oncogene in GC through the downregulation of SOCS3<sup>[25]</sup>. The regulatory effects of miR-668-3p on SOCS3 in GC

cells has not been investigated. Here, we show for the first time that circ\_0003356 is downregulated in GC cells and tumor tissues and reveal its association with GC progression.

## **MATERIALS AND METHODS**

### ***Sample collection***

GC and normal tissues ( $n = 80$  pairs) were collected from surgical specimens of patients with GC at Zhongshan City People's Hospital. Patients were diagnosed with GC *via* histopathology. No patients had received prior immunotherapy, adjuvant chemotherapy, or radiation therapy before surgery. GC patients were allocated into circ\_0003356 low or high-expression groups based on the median expression value in tumor tissues. The Ethics Committee of Zhongshan City People's Hospital provided approval (K2017-182). All participants agreed to the study and provided informed written consent.

### ***Cell culture***

AGS and HGC-27 cell lines (human) and GES-1 (non-GC) were cultured in Roswell Park Memorial Institute 1640 medium + 10% fetal bovine serum (FBS) + 1% streptomycin-penicillin. Cells were maintained at 37 °C 5% CO<sub>2</sub>.

**Quantitative real time-polymerase chain reaction:** Cells were lysed in Trizol and total RNA was quantitated on a NanoDrop 2000c spectrophotometer. For circ\_0003356 and SOCS3, PrimeScript™ RT reagent kits were used for complementary DNA (cDNA) synthesis. For miR-668-3p, a TaqMan miRNA Assay kit was used for cDNA synthesis. cDNA was amplified using BeyoFast™ SYBR Green qPCR Mix. Primer sequences as shown in Table 1 were bought from TaKaRa. Relative mRNA expression of circ\_0003356 and SOCS3 were normalised to GAPDH and calculated using the  $2^{-\Delta\Delta Ct}$  method. miR-668-3p was normalised to U6.

### ***Cell transfection***

The empty pcDNA3.1 vector, circ\_0003356 overexpression vector (pcDNA3.1-circ\_0003356), miR-668-3p inhibitor, negative control (NC) inhibitor, miR-668-3p mimic, NC mimic, small interfering RNA (siRNA) against SOCS3 (si-SOCS3) and si-NC were bought from GenePharma (Shanghai, China). Above factors were then transfected into AGS and HGC-27 cells using Lipofectamine 3000 (Invitrogen, Carlsbad, California, United States) for 48 h.

**CCK-8 assay:** The proliferation of HGC-27 and AGS cells was evaluated *via* a cell counting kit-8 (Dojindo, Osaka, Japan). HGC-27 and AGS cells ( $1 \times 10^4$  cells/well) were plated in 96-well plates for incubating 24 h, 48 h, 72 h and 96 h. After that, each well was treated with CCK-8 reagent (10  $\mu$ L) for another 1 h. Ultimately, a microplate reader (Bio-Rad, CA, United States) was used for measurement of optical density at 450 nm.

**EdU assays:** A Cell-Light EdU DNA Cell Proliferation Kit (RiboBio, Guangzhou, China) was used to perform EdU assay. Cells were labeled with 50  $\mu$ M EdU for 2 h and fixed with 4 % paraformaldehyde (PFA) (4%). Cells were washed in phosphate buffered saline (PBS) containing glycine (2 mg/mL) for 10 min to remove artefacts and permeabilized. Cells were stained with Apollo dye liquor in the dark followed by DAPI. Cells were imaged on a fluorescence microscope.

**Flow cytometry analysis:** Apoptotic rates were measured by flow cytometry (FACSCalibur, BD, United States). Cells were resuspended in 300  $\mu$ L binding buffer and incubated with + Annexin V-FITC (5  $\mu$ L) for 10 min. Cells were labeled with propidium iodide (5  $\mu$ L) in the dark for 5 min. Apoptotic cells were quantified using FlowJo V10 software.

**Wound healing assay:** HGC-27 and AGS cells ( $5 \times 10^4$  cells/well) were put to 6-well plates coated with extracellular matrix molecule, which were cultured in RPMI1640

including 10% FBS. When the cell monolayer was formed, a sterile pipette tip was utilized for scratching the monolayer and scratch width was recorded under an inverted microscope (TE2000; Nikon, Tokyo, Japan). After being washed *via* PBS, FBS-free medium was added to culture cells for 24 h, followed by record of scratch wound. At last, the migration rate was calculated according to the following formula:  $(1 - 24 \text{ h scratch width} / 0 \text{ h scratch width}) \times 100$ .

**Transwell assay:** Matrigel-coated transwell chambers (8  $\mu\text{m}$ ; BD Biosciences, Franklin Lakes, NJ, United States) were utilized for evaluating invasion of HGC-27 and AGS cells. In detail,  $3 \times 10^3$  cells were re-suspended in FBS-free RPMI1640 medium (200  $\mu\text{L}$ ), which were appended to the upper chamber. In contrast, RPMI1640 medium containing 10% FBS was appended to bottom chambers. Following 24 h of incubation, cells in upper chambers were removed *via* a cotton swab. Cells in bottom chambers were fixed *via* 4% PFA and stained *via* 0.1% crystal violet. At last, the number of these cells was counted through inverted fluorescence microscopy (Leica Microsystems GmbH, Wetzlar, Germany).

**Cytoplasmic and nuclear fractions:** AGS and HGC-27 cells were lysed on ice in lysis buffer + protease inhibitors and centrifuged for 3 min. The subsequent pellets and supernatants were harvested as nuclear and cytoplasmic fractions. RNA was extracted from each fraction using Buffer SK. Quantitative real-time polymerase chain reaction (qRT-PCR) analysis was performed for the detection of circ\_0003356 expression. U6 and 18S rRNA were used as controls of cytoplasmic and nuclear transcripts, respectively.

**Fluorescence *in situ* hybridization (FISH):** Fluorescence *in situ* hybridization (FISH) was performed using specific probes against circ\_0003356 and miR-668-3p. Briefly, cells were fixed in 4% PFA for 15 min and washed with a 70%, 95%, and 100% gradient of ethanol. Cell hybridization was performed at 37 °C in a dark incubator. Cells were blocked in 3% normal goat serum and 1% bovine serum albumin for 1 h. Cells were

cultured with HRP-conjugated anti-biotin antibodies at 4 °C for 1 d. Cells were imaged on a fluorescence microscope.

**Dual luciferase reporter assay:** The online database circinteractome and StarBase v2.0 were used to identify predicted targets of circ\_0003356 and miR-668-3p. The interaction of circ\_0003356 and miR-668-3p or SOCS3 and miR-668-3p were verified *via* dual luciferase reporter (DLR) assays. Briefly, the 3'-UTR sequences of WT circ\_0003356 or SOCS3 comprising the miR-668-3p binding sites were cloned into pRL-CMV. 3'-UTR sequences of circ\_0003356 or SOCS3 containing mutant sites for miR-668-3p were also generated. MiR-668-3p mimic or NC mimic were transfected into HGC-27 and AGS cells and luciferase activity was measured 2 d post-transfection.

**RNA binding protein immunoprecipitation assay:** Cells were lysed in RNA binding protein immunoprecipitation (RIP) buffer and magnetic beads conjugated with anti-Ago2 and anti-immunoglobulin G (IgG) were added to cell lysates. MiR-668-3p and circ\_0003356 levels in the beads were analyzed.

**RNA pull-down assays:** RNA pull-downs were performed using commercial Magnetic RNA-Protein Pull-Down Kits (Pierce). Cells were transfected with 3'-biotinylated miR-668-3p (Bio-miR-668-3p) or Bio-NC mimics. After 2 d, cells were lysed and streptavidin-coupled beads added to pull-down biotin-coupled RNA complexes. Circ\_0003356 enrichment in the bound fractions was subsequently assayed.

**Western blot analysis:** Cells were lysed in RIPA buffer and protein content assessed *via* BCA assays. A total of 20 µg of protein was resolved on 10% sodium dodecyl sulphate-polyacrylamide gel electrophoresis (SDS-PAGE) gels and transferred to on polyvinylidene difluoride membranes. Membranes were blocked in non-fat milk for 1 h and probed with anti-N-cadherin (1:6000, ab76011), anti-E-cadherin (1:10000, ab40772), anti-SOCS3 (1:1000, ab280884) and β-actin (1:1000, ab8224) primary antibodies at 4 °C

for 24 h. Cells were subsequently labeled with anti-rabbit secondary antibodies (1:2000, ab205718) and proteins were visualized using chemiluminescence.

**Establishment of xenograft tumor models:** Mice (4-5 wk-old; BALB/c nude; male) were housed at room temperature with 50% humidity and 12 h light/dark cycle. All experiments were approved by the Animal Care and Use Committee of Zhongshan City People's Hospital (K2017-182).

Mice were injected into the upper back with  $1 \times 10^7$  AGS cells transfected with circ\_0003356 or control vector<sup>[26]</sup>. Mice were assigned into 2 groups ( $n = 6$  mice/group): The circ\_0003356 group and the vector group. After injection for five weeks, mice were anesthetized with pentobarbital sodium and sacrificed *via* dislocation of cervical vertebrae. Tumors were dissected and tumor volumes calculated ( $\text{length} \times \text{width}^2$ )/2.

**Immunohistochemistry staining:** Mice tumor tissues were fixed in formalin and paraffinized. Tumors were cut into 4  $\mu\text{m}$  sections, dewaxed with xylene, and hydrated in ethanol. Sections were boiled in citrate buffer (10 mmol/L) and cultured using  $\text{H}_2\text{O}_2$  solution (3%) to inhibit endogenous peroxidase activity. Sections were probed with anti-Ki67 (1:200, ab16667), anti-SOCS3 (1:1000, ab280884), anti-N-cadherin (1:500, ab76011), and anti-E-cadherin (1:500, ab40772) primary antibodies at 4 °C overnight and labeled with HRP-conjugated secondary antibodies (1:500, ab6112) 30 min. Cells were stained using DAB substrate and counterstained using hematoxylin. Cells were imaged on a light microscope.

**TUNEL staining:** TUNEL assay was performed using the commercially available *in situ* Cell Death Detection Kit as per the manufactures recommendations.

### *Statistical analysis*

GraphPad Prism 8.0 was used for data analysis. Experiments were performed on a minimum of 3 occasions and are shown as the mean  $\pm$  SD. Survival curves were plotted using the Kaplan-Meier method. Data were compared using a student's *t*-test.

## **RESULTS**

### ***Circ\_0003356 expression is down-regulated in GC tissues***

Published GEO and GSE184882 datasets were analyzed<sup>[27,28]</sup>. Five circRNAs with significant expression differences in GC tissues were identified and selected for heat map analysis (Figure 1A). Then their expression was further determined *via* qRT-PCR. Five circRNAs with low relative expression in tumor tissues compared to normal tissues were identified ( $n = 20$ ), amongst which circ\_0003356 expression ( $0.2 \pm 0.06532$ ) was the lowest ( $P = 0.0002$ , Figure 1B). Circ\_0003356 was selected for further analysis.

Circ\_0003356 expression was validated in GC normal and tumor tissues ( $n = 80$ ). As expected, circ\_0003356 expression was markedly downregulated in tumor tissues ( $0.4908 \pm 0.02457$ ) compared to normal adjacent tissues ( $0.99 \pm 0.0243$ ;  $P < 0.0001$ , Figure 1C). Patients with low circ\_0003356 expression correlated with a short survival time ( $P < 0.0001$ , Figure 1D). Moreover, circ\_0003356 displayed lower expression in AGS ( $0.3833 \pm 0.06012$ ) and HGC-27 cells ( $0.4967 \pm 0.06012$ ) than GES-1 cells ( $P = 0.0004$ ,  $P = 0.0001$ , Figure 1E).

### ***Circ\_0003356 overexpression promotes cell apoptosis and impairs invasion, proliferation, migration and EMT of AGS and HGC-27 cells***

Circ\_0003356 was overexpressed in AGS and HGC-27 cells to further explore its influence on GC cell proliferation, apoptosis, migration, invasion, and EMT. Circ\_0003356 expression in AGS cells ( $2.613 \pm 0.05207$ ) and HGC-27 cells ( $2.5 \pm 0.05774$ ) was dramatically increased following transfection of pcDNA3.1-circ\_0003356 ( $P < 0.0001$ ,  $P < 0.0001$ , Figure 2A). Functional analyses showed that the viability of AGS ( $1.183 \pm 0.06155$ ) and HGC-27 cells ( $1.247 \pm 0.0636$ ) at 96 h were attenuated after circ\_0003356 overexpression ( $P = 0.0002$ ,  $P = 0.0002$ , Figure 2B). EdU assays also

revealed that the proliferative ability of AGS ( $22.33 \pm 1.764$ ) and HGC-27 cells ( $20 \pm 1.732$ ) was attenuated by circ\_0003356 overexpression ( $P = 0.0025$ ,  $P = 0.0043$ , Figure 2C). Apoptotic rates in AGS ( $22.68 \pm 0.2887$ ) and HGC-27 cells' ( $18.38 \pm 0.2887$ ) were increased by circ\_0003356 overexpression ( $P < 0.0001$ ,  $P < 0.0001$ , Figure 2D). In terms of the migration and invasion of AGS ( $24.33 \pm 2.728$ ) and HGC-27 cells ( $20.33 \pm 1.764$ ), decreased numbers of invading AGS ( $168 \pm 11.27$ ) and HGC-27 cells ( $161.7 \pm 12.35$ ) were observed in response to circ\_0003356 overexpression ( $P = 0.0001$ ,  $P = 0.0031$ ,  $P = 0.0002$ ,  $P = 0.0016$ , Figures 2E and F). Circ\_0003356 overexpression also attenuated the EMT of AGS ( $1.07 \pm 0.6531$ ,  $0.57 \pm 0.5168$ ) and HGC-27 ( $0.93 \pm 0.06533$ ,  $0.39 \pm 0.06137$ ) cells evidenced by reduced N-cadherin expression and the enhanced expression of E-cadherin ( $P = 0.0006$ ,  $P = 0.0023$ ,  $P = 0.0021$ ,  $P = 0.0007$ , Figure 2G).

#### ***Circ\_0003356 directly binds to miR-668-3p***

The subvocalization of circ\_0003356 in AGS and HGC-27 cells was confirmed through FISH, which showed that circ\_0003356 located in AGS and HGC-27 cells' cytoplasm (Figure 3A). The findings were confirmed by qRT-PCR, showing that the circ\_0003356 level was in AGS and HGC-27 cells markedly higher in the cytoplasm (85%) than the nuclear (15%) (Figure 3B).

Target predictions were performed using circinteractome, indentifying binding sites between miR-668-3p and circ\_0003356 (Figure 3C). We verified this relationship *via* RIP-qRT-PCR, RNA pull-down, and DLR assays. The Ago2 group showed enrichment for circ\_0003356 and miR-668-3p in AGS ( $12.3 \pm 0.7024$ ,  $15.33 \pm 0.3801$ ) and HGC-27 cells ( $11.3 \pm 0.6272$ ,  $14.33 \pm 0.6912$ ) compared to the IgG group ( $P = 0.000087$ ,  $P = 0.0000029$ ,  $P = 0.00008$ ,  $P = 0.00004$ , Figure 3D). The biotinylated miR-668-3p probe enriched more circ\_0003356 than the biotinylated NC probe in AGS ( $0.9033 \pm 0.03756$ ) and HGC-27 cells ( $0.9167 \pm 0.0491$ ) ( $P = 0.0002$ ,  $P = 0.0004$ , Figure 3E). Relative luciferase activity in AGS ( $0.35 \pm 0.06532$ ) and HGC-27 ( $0.31 \pm 0.06478$ ) were reduced in cells co-transfected with miR-668-3p mimic and circ\_0003356 WT relative to miR-668-3p mimic and circ\_0003356 MUT ( $P = 0.00057$ ,  $P = 0.00041$ , Figure 3F). High expression of miR-668-3p

was observed in tumor tissues ( $2.006 \pm 0.02872$ ) compared to healthy tissues ( $0.9984 \pm 0.02829$ ) ( $P < 0.0001$ , Figure 3G). An inverse relationship between the expression of miR-668-3p and circ\_0003356 in tumor tissues was also observed ( $r = -0.8783$ ,  $P = 0.0007$ , Figure 3H).

***Suppression of miR-668-3p promotes apoptosis and reduces proliferation, migration, invasion, and EMT in AGS and HGC-27 cells***

Accordingly, the influences of miR-668-3p on cell migration, apoptosis, proliferation, invasion, and EMT were explored. Firstly, miR-668-3p expression in HGC-27 ( $2.8 \pm 0.05508$ ) and AGS ( $3 \pm 0.1155$ ) cells was found to be remarkably higher than GES-1 cells ( $1.007 \pm 0.05207$ ) ( $P < 0.0001$ ,  $P < 0.0001$ , Figure 4A). When we determined its specific functions, we found that the viability ( $1.393 \pm 0.06128$ ,  $1.342 \pm 0.0636$ ) and proliferative capacity ( $12.33 \pm 1.453$ ,  $14 \pm 2.082$ ) of AGS and HGC-27 cells were attenuated by miR-668-3p suppression ( $P = 0.00037$ ,  $P = 0.00039$ ,  $P = 0.0033$ ,  $P = 0.0020$ , Figures 4B and C). Apototic rates in AGS ( $21.29 \pm 0.8838$ ) and HGC-27 cells' ( $21.09 \pm 0.9074$ ) were raised following miR-668-3p suppression ( $P < 0.0001$ ,  $P = 0.0001$ , Figure 4D). The migration ( $25.67 \pm 2.603$ ,  $41.67 \pm 2.906$ ) and invasion ( $174.7 \pm 9.262$ ,  $175.7 \pm 9.025$ ) of AGS and HGC-27 cells were also repressed following miR-668-3p down-regulation ( $P = 0.0001$ ,  $P = 0.0019$ ,  $P = 0.0026$ ,  $P = 0.0023$ , Figures 4E and F). The EMT of AGS ( $1.09 \pm 0.06348$ ,  $0.4 \pm 0.06137$ ) and HGC-27 cells ( $0.9 \pm 0.05627$ ,  $0.31 \pm 0.06137$ ) was similarly attenuated by miR-668-3p suppression, evidenced by decreased N-cadherin and increased E-cadherin expression ( $P = 0.00089$ ,  $P = 0.00051$ ,  $P = 0.00066$ ,  $P = 0.00054$ , Figure 4G).

***SOCS3 is targeted by miR-668-3p***

MiR-668-3p targets were assessed using the starbase2.0 database. SOCS3 contained binding sequences (AGUGAC) of miR-668-3p (Figure 5A). DLR assay was performed to validate this interaction. Relative luciferase activity in AGS ( $0.37 \pm 0.06532$ ) and HGC-27 cells ( $0.32 \pm 0.06532$ ) transfected with SOCS3 WT reporter was reduced in response to transfection of miR-668-3p mimic ( $P = 0.00041$ ,  $P = 0.00065$ , Figure 5B). The SOCS3 MUT

reporter showed no such changes following miR-668-3p mimic addition (Figure 5B), suggestive of a direct interaction between miR-668-3p and SOCS3. The regulatory relation between miR-668-3p and SOCS3 was assessed through western blotting. SOCS3 expression in AGS and HGC-27 cells was reduced through the addition of miR-668-3p mimic ( $0.5833 \pm 0.03756$ ,  $0.6633 \pm 0.03528$ ) and boosted by the miR-668-3p inhibitor ( $1.03 \pm 0.04619$ ,  $0.92 \pm 0.03786$ ), implying an inverse regulatory relationship between miR-668-3p and SOCS3 ( $P = 0.0076$ ,  $P = 0.0036$ ;  $P = 0.0038$ ,  $P = 0.0047$ , Figures 5C and D). Lower expression of SOCS3 in tumor tissues ( $0.5001 \pm 0.02447$ ) compared to normal tissues ( $0.9979 \pm 0.02526$ ) ( $P < 0.0001$ , Figure 5E) was also observed. Upon correlation analysis between circ\_0003356 and SOCS3 expression ( $P < 0.0001$ ,  $r = 0.9364$ ; Figure 5F), the expression of SOCS3 was negatively associated with miR-668-3p in tumor tissues ( $P < 0.0001$ ,  $r = -0.8438$ ; Figure 5G).

#### *Circ\_0003356 inhibits malignant behavior of GC cells via sponging miR-668-3p to target SOCS3*

SOCS3 was silenced *via* transfection of si-SOCS3 and its reduced expression confirmed in AGS ( $0.7267 \pm 0.05044$ ) and HGC-27 cells ( $0.5933 \pm 0.05207$ ) ( $P = 0.0086$ ,  $P = 0.0065$ , Figure 6A). miR-668-3p overexpression ( $0.6467 \pm 0.06394$ ,  $0.6633 \pm 0.04322$ ) or SOCS3 downregulation ( $0.6433 \pm 0.06394$ ,  $0.6267 \pm 0.04527$ ) reversed the promoting effect of circ\_0003356 on SOCS3 expression level in AGS and HGC-27 cells ( $P = 0.0012$ ,  $P = 0.0032$ ,  $P = 0.0030$ ,  $P = 0.0022$ ,  $P = 0.0031$ ,  $P = 0.0031$ , Figure 6B). The viability and proliferative capacity of AGS and HGC-27 cells were attenuated by circ\_0003356 overexpression, whilst miR-668-3p upregulation or SOCS3 downregulation recovered the circ\_0003356-mediated decrease in cell viability and proliferative capacity ( $P = 0.00039$ ,  $P = 0.0025$ ,  $P = 0.0037$ ,  $P = 0.00042$ ,  $P = 0.0035$ ,  $P = 0.0051$ ;  $P < 0.0001$ ,  $P = 0.0002$ ,  $P < 0.0001$ ,  $P < 0.0001$ ,  $P = 0.0044$ ,  $P = 0.0012$ ; Figures 6C and D). The effects of circ\_0003356 on the apoptotic rates of AGS and HGC-27 cells were similarly reversed by miR-668-3p upregulation or SOCS3 downregulation ( $P < 0.0001$ , Figure 6E). When it came to circ\_0003356-mediated inhibition of migration, invasion, and EMT, miR-668-3p

up-regulation or SOCS3 down-regulation could partially eliminate them ( $P < 0.0001$ ,  $P = 0.0017$ ,  $P = 0.0007$ ,  $P < 0.0001$ ,  $P = 0.0039$ ,  $P = 0.0011$ ;  $P < 0.0001$ ,  $P = 0.0009$ ,  $P = 0.0004$ ,  $P < 0.0001$ ,  $P = 0.0006$ ,  $P = 0.0003$ ;  $P = 0.0028$ ,  $P = 0.010$ ,  $P = 0.0164$ ,  $P = 0.0029$ ,  $P = 0.023$ ,  $P = 0.0278$ ;  $P = 0.00314$ ,  $P = 0.0213$ ,  $P = 0.0144$ ,  $P = 0.00285$ ,  $P = 0.0267$ ,  $P = 0.00914$ ; Figures 6F-H).

### *Circ\_0003356 impedes GC tumorigenesis in vivo*

A xenograft mouse model was constructed to confirm the anti-tumor role of circ\_0003356 in GC. As depicted in Figures 7A-C, tumor size, volume ( $110 \pm 5.859$ ,  $P = 0.0001$ ), and weight ( $0.2833 \pm 0.03528$ ,  $P = 0.0007$ ) were reduced when circ\_0003356 was overexpressed. Decreased expressions of Ki67 (proliferative marker protein) and N-cadherin were observed following circ\_0003356 overexpression, while SOCS3 and E-cadherin expression increased, suggestive of inhibitory effects on cell proliferation and EMT (Figure 7D). TUNEL staining indicated that circ\_0003356 upregulation increased apoptosis in tumor tissues of mice (Figure 7E). Circ\_0003356 expression was found to be boosted ( $2.423 \pm 0.09135$ ) and expression of miR-668-3p ( $0.5 \pm 0.06531$ ) was diminished after circ\_0003356 was overexpressed in mouse tumor tissues ( $P = 0.000099$ ,  $P = 0.0016$ , Figure 7F).

## **DISCUSSION**

Due to a lack of effective treatment strategies and prognostic indicators, GC remains a threat to human health<sup>[29,30]</sup>. Emerging evidence has implicated circRNAs during GC development<sup>[31,32]</sup>. Thence, great efforts should be made to comprehensively understand circRNAs. To date, many circRNAs show low expression in GC, such as circ\_0021087<sup>[12]</sup>, hsa\_circ\_0005556<sup>[33]</sup>, and circPSMC3<sup>[34]</sup>. Here, we observed low expression levels of circ\_0003356 in GC cells and tissues. This is the first report to show an association of circ\_0003356 with GC. Moreover, increasing findings have indicated that several circRNAs act as prognostic indicators for GC, including hsa\_circ\_0005556<sup>[33]</sup> and

circ\_0000260<sup>[35]</sup>. Here, based on Kaplan-Meier analysis, patients with low circ\_0003356 had poor survival rates.

Accumulating data reveal that circRNAs have an anti-tumor function in GC. For instance, circ\_0021087 overexpression suppresses the proliferation, migration, invasion, and EMT of GC cells and impairs tumorigenesis in xenograft mice<sup>[12]</sup>. CircPSMC3 up-regulation prevents GC cells from migrating, proliferating, and invading, thereby suppressing tumorigenesis in xenograft mice<sup>[36]</sup>. Similarly, we found that circ\_0003356 overexpression repressed proliferation, migration, invasion, and EMT of GC cells and enhanced GC cell apoptosis. *In vivo*, circ\_0003356 overexpression suppressed tumorigenesis. These results suggest an anti-tumor role for circ\_0003356's in GC, highlighting its potential in the clinical management of GC.

To our knowledge, circRNAs affect gene expression *via* sponging downstream miRNAs in cancer cell<sup>[37,38]</sup>. Inspired by this, we predicted downstream miRNAs of circ\_0003356 using circinteractome. MiR-668-3p was identified as one such target. We performed RIP-qRT-PCR, RNA pull-downs, and DLR assays to verify our prediction. In previous reports, miR-668-3p has shown high expression and carcinogenesis in HCC and CRC<sup>[18,19]</sup>. Consistent with these studies, we discovered high miR-668-3p expression in GC tissues and cells. The down-regulation of miR-668-3p enhanced cell apoptosis and impaired the proliferation, migration, invasion, and EMT of cells (AGS and HGC-27). These outcomes suggest that miR-668-3p is oncogenic in GC. Several circRNAs, including circ\_0014717 and circTMEM59, target miR-668-3p to suppress tumor development<sup>[18,19]</sup>. Here, an inverse relationship was observed between miR-668-3p and circ\_0003356 expression in tumor tissues of GC patients. MiR-668-3p was negatively regulated by circ\_0003356 in mouse tumor tissues. In particular, the inhibitory influence of circ\_0003356 on the proliferation, migration, invasion. EMT of GC cells were reversed by miR-3619-5p overexpression. We deduced that circ\_0003356 prevents GC malignancy through sponging miR-668-3p.

It is acknowledged that the miRNA-circRNA-mRNA network is closely related to the regulation of GC progression<sup>[39]</sup>. We therefore predicted target genes of miR-668-3p and

found that SOCS3 was targeted by miR-668-3p. SOCS3 has been shown to display low expression in GC tissues and to participate in tumor suppression in GC<sup>[24,40]</sup>. In this study, we observed that the downregulation of SOCS3 expression in GC tissues was regulated by miR-668-3p. An inverse correlation between SOCS3 and miR-668-3p expression levels in tumor tissues of GC patients was observed. Thus, we deduced that miR-668-3p acts as a tumor promoter, targeting SOCS3 in GC. Additionally, we identified a positive relationship between SOCS3 and circ\_0003356 and found that the suppressive effects of circ\_0003356 on cell proliferation, migration, invasion, and EMT were reversed by SOCS3 upregulation in GC cells. These findings strongly support the notion that circ\_0003356 has an anti-tumor role through sponging miR-668-3p to target SOCS3 in GC.

## **CONCLUSION**

In summary, we show that circ\_0003356 expression is low in GC tissues and cells and positively related to GC patient prognosis. Circ\_0003356 up-regulation or miR-668-3p inhibition facilitated cell apoptosis and suppressed proliferation, migration, invasion and EMT in cells (AGS and HGC-27). Circ\_0003356 overexpression also prevented tumor growth in mice. Mechanistically, circ\_0003356 was found to interact with the miR-668-3p/SOCS3 axis to impair GC development. Overall, this investigation provides a new perspective on the therapeutic targets of GC.

## **ARTICLE HIGHLIGHTS**

### ***Research background***

Gastric cancer (GC) is a common malignant tumor with high prevalence and mortality. Circular RNA (circRNA) analysis in GSE184882 has indicated that circ\_0003356 shows abnormal expression in GC tissues, but the function of circ\_0003356 remains unclear in GC.

### ***Research motivation***

To seek the prognostic biomarker and therapeutic target for GC.

### ***Research objectives***

In order to explore the function and mechanism of circ\_0003356 in GC.

### ***Research methods***

Quantitative real-time polymerase chain reaction was used to detect expression of circ\_0003356. The clinical value of circ\_0003356 for GC prognosis was evaluated. The role of circ\_0003356 in GC was assessed *via in vitro* and *in vivo* experiments. The molecular mechanism of circ\_0003356 on miR-668-3p/SOCS3 axis was validated.

### ***Research results***

Circ\_0003356 expression was markedly decreased in GC tissues and cells, and it was positively associated with survival time of GC patients. Moreover, functional analyses showed that circ\_0003356 overexpression inhibited malignant behaviors of GC cells *via* targeting the miR-668-3p/SOCS3 axis.

### ***Research conclusions***

Circ\_0003356 is lowly expressed in GC tissues and cells, displaying potential as a prognostic biomarker for GC. Circ\_0003356 up-regulation represses the malignancy of GC *via* targeting the miR-668-3p/SOCS3 axis.

### ***Research perspectives***

Other mechanisms of circ\_0003356 in GC may be probed in future researches, and its application in treatment of GC will be extended.

# 3%

SIMILARITY INDEX

### PRIMARY SOURCES

- 1

Yunlong Wu, Yuying Guo, Qiang Wang. "USP21 accelerates the proliferation and glycolysis of esophageal cancer cells by regulating the STAT3/FOXO1 pathway", Tissue and Cell, 2022  
Crossref

48 words — 1%
- 2

Xufeng Guo, Ming Li. "LINC01089 is a tumor-suppressive lncRNA in gastric cancer and it regulates miR-27a-3p/TET1 axis", Cancer Cell International, 2020  
Crossref

20 words — < 1%
- 3

f6publishing.blob.core.windows.net  
Internet

16 words — < 1%
- 4

www.wjgnet.com  
Internet

15 words — < 1%
- 5

jocr-online.biomedcentral.com  
Internet

13 words — < 1%
- 6

Jia Cao, Rong Ma, Jia Wang, Bin Shi, Jufen Zhao, Jinhai Tian, Qi Huang, Jingjing Yu, Xiang Tao, Libin Wang. "NKD1 targeting PCM1 regulates the therapeutic sensitivity of HHT on colorectal cancer cells", Research Square Platform LLC, 2023  
Crossref Posted Content

12 words — < 1%

---

7 Jinzhu Han, Zixin Yang, Shan Zhao, Likang Zheng, Yanhua Tian, Yingqian Lv. "Circ\_0027599 elevates RUNX1 expression via sponging miR - 21 - 5p on gastric cancer progression", European Journal of Clinical Investigation, 2021  
Crossref 12 words — < 1%

---

8 Rong He, Heng Zhang, Lian He, Jiabin Huang, Zhe Dai, Shouyan Xiang, Yongjing Zhou, Yakun Lang, Dandan Gong, Yu Fan. "HOXA3-Induced Exosomal lncRNA RNF144A-AS1 Promotes Liver Metastasis of Gastric Cancer via Interacting with PUF60 Protein and Sponging miR-361-3p", Research Square Platform LLC, 2021  
Crossref Posted Content 12 words — < 1%

---

9 [www.celljournal.org](http://www.celljournal.org)  
Internet 12 words — < 1%

---

EXCLUDE QUOTES ON  
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE SOURCES < 12 WORDS  
EXCLUDE MATCHES < 12 WORDS