

87998\_Auto\_Edited-check.docx

**Name of Journal:** *World Journal of Gastrointestinal Oncology*

**Manuscript NO:** 87998

**Manuscript Type:** META-ANALYSIS

**Effectiveness and tolerability of programmed cell death protein-1 inhibitor + chemotherapy compared to chemotherapy for upper gastrointestinal tract cancers**

Zhang XM *et al.* Meta-analysis of PD-1 inhibitor' effectiveness

## Abstract

### BACKGROUND

The combination of programmed cell death protein-1 (PD-1) inhibitor and chemotherapy is approved as a standard first- or second-line treatment in patients with advanced oesophageal or gastric cancer. However, it is unclear whether this combination is superior to chemotherapy alone.

### AIM

To assess the comparative effectiveness and tolerability of combining PD-1 inhibitors with chemotherapy *vs* chemotherapy alone in patients with advanced gastric cancer, gastroesophageal junction (GEJ) cancer, or oesophageal carcinoma.

### METHODS

We searched the PubMed and Embase databases for studies that compared the efficacy and tolerance of PD-1 inhibitors in combination with chemotherapy *vs* chemotherapy alone in patients with advanced oesophageal or gastric cancer. We employed either random or fixed models to analyze the outcomes of each clinical trial, encompassing data on overall survival (OS), progression-free survival (PFS), objective response rate, and adverse events (AEs).

### RESULTS

Nine phase 3 clinical trials (7016 advanced oesophageal and gastric cancer patients) met the inclusion criteria. Our meta-analysis demonstrated that the pooled PD-1 inhibitor + chemotherapy group had a significantly longer OS than the chemotherapy-alone group [hazard ratio (HR) = 0.76, 95% confidence interval (CI): 0.71-0.81]; the pooled PFS result was consistent with that of OS (HR = 0.67, 95%CI: 0.61-0.74). The count of patients achieving an objective response in the PD-1 inhibitor + chemotherapy group surpassed that of the chemotherapy-alone group [odds ratio (OR) = 1.86, 95%CI: 1.59-2.18]. AE incidence was also higher in the combination-therapy group than in the

chemotherapy-alone group, regardless of whether  $\geq$  grade 3 only (OR = 1.30, 95%CI: 1.07-1.57) or all AE grades (OR = 1.88, 95%CI: 1.39-2.54) were examined. We performed a subgroup analysis based on the programmed death-ligand 1 (PD-L1) combined positive score (CPS) and noted extended OS and PFS durations within the CPS  $\geq$  1, CPS  $\geq$  5, and CPS  $\geq$  10 subgroups of the PD-1 inhibitor + chemotherapy group.

## CONCLUSION

In contrast to chemotherapy alone, the combination of PD-1 inhibitor and chemotherapy appears to present a more favorable option for initial or subsequent treatment in patients with gastric cancer, GEJ tumor, or oesophageal cancer. This holds true particularly for individuals with PD-L1 CPS scores of  $\geq$  5 and  $\geq$  10.

**Key Words:** Programmed cell death protein-1 inhibitor; Chemotherapy; Oesophageal squamous cell carcinoma; Gastric/gastroesophageal junction adenocarcinoma; Overall survival; Progression-free survival; Objective response rate; Adverse event

Zhang XM, Yang T, Xu YY, Li BZ, Shen W, Hu WQ, Yan CW, Zong L. Effectiveness and tolerability of programmed cell death protein-1 inhibitor + chemotherapy compared to chemotherapy for upper gastrointestinal tract cancers. *World J Gastrointest Oncol* 2024; In press

**Core Tip:** The combination of programmed cell death protein-1 (PD-1) inhibitor and chemotherapy is approved as a standard first- or second-line treatment in patients with advanced oesophageal or gastric cancer. However, it is unclear whether this combination is superior to chemotherapy alone. We assessed the comparative effectiveness and tolerability of combining PD-1 inhibitors with chemotherapy *vs* chemotherapy alone in patients with advanced gastric cancer, gastroesophageal junction cancer, or oesophageal carcinoma. Our analysis showed that immunotherapy combined with chemotherapy significantly prolonged patients' overall survival and

progression-free survival relative to the chemotherapy group, both in the overall population and in the combined positive score (CPS)  $\geq 1$ , CPS  $\geq 5$ , and CPS  $\geq 10$  subgroup.

## **INTRODUCTION**

Upper gastrointestinal tract (UGT) cancers stand as the prevailing form of malignancy, encompassing oesophageal squamous cell carcinoma (ESCC) and gastric/gastroesophageal junction (G/GEJ) adenocarcinomas<sup>[1]</sup>. According to Global Cancer Statistics 2020, an estimated 1089103 and 604100 new cases of gastric and oesophageal cancers, respectively, occur each year, and the annual numbers of new deaths are 768793 and 544076<sup>[2]</sup>. The presently employed methods for treating UGT cancers are surgery, chemotherapy, and radiotherapy. Even with continuous enhancements in chemoradiotherapy, the clinical results for ESCC and G/GEJ adenocarcinoma have not exhibited substantial advancement over time, underscoring the persisting gap in effective treatment<sup>[3]</sup>.

Immunotherapy is an evolving approach to cancer treatment and is designed to destroy cancer cells by strengthening patients' immune capabilities. Various immunotherapies encompass the utilization of immune checkpoint inhibitor (ICIs), chimeric antigen receptor T-cell therapy, cancer vaccines, and adoptive cell therapy<sup>[4]</sup>. Programmed cell death protein-1 (PD-1) functions as a checkpoint receptor expressed on the surface of diverse immune cells. Blocking the interaction between PD-1 and programmed death-ligand 1 (PD-L1) serves as an effective means to bolster the immune response<sup>[5]</sup>. Anti-PD-1 antibodies such as nivolumab, pembrolizumab, sintilimab, camrelizumab, and tislelizumab demonstrated improved antitumor activity alongside manageable safety profiles in patients with oesophageal or gastric cancer<sup>[6]</sup>.

Combinations of a PD-1 inhibitor and chemotherapy have gained approval as standard first or second-line treatment choices for patients with advanced oesophageal and gastric cancer in clinical settings<sup>[7]</sup>. While various studies have outlined the advantages of pairing PD-1 inhibitor therapy with chemotherapy, it remains

undetermined whether the combination of a PD-1 inhibitor + chemotherapy surpasses the efficacy of chemotherapy alone. Tabernero *et al*<sup>[8]</sup> revealed that the combination of pembrolizumab with chemotherapy did not exhibit superiority over chemotherapy alone in terms of the tested overall survival (OS) and progression-free survival (PFS) endpoints. In a separate study involving patients with gastric or GEJ cancer, the addition of nivolumab to chemotherapy did not lead to a significant improvement in OS compared to the group treated with chemotherapy alone<sup>[9]</sup>. Therefore, it remains essential to ascertain whether chemotherapy alone or the combination of PD-1 inhibitors with chemotherapy yields a more favorable curative effect. We performed this current meta-analysis to assess the comparative efficacy and tolerability of PD-1 inhibitors combined with standard chemotherapy in contrast to chemotherapy alone, specifically in patients diagnosed with advanced gastric cancer, GEJ cancer, or oesophageal adenocarcinoma. Our aim was to establish a foundation for clinical treatment choices for future patients afflicted with UGT cancer.

## **MATERIALS AND METHODS**

### ***Search strategy***

To identify relevant studies, we performed a comprehensive literature search of the electronic databases PubMed (<https://pubmed.ncbi.nlm.nih.gov/>) and Embase ([embase.com/](https://www.embase.com/)). Filter articles published before July 2023. We used search terms pertaining to diseases (*e.g.*, GEJ adenocarcinoma, gastric, gastric cancer, gastro-oesophageal junction cancer, oesophageal squamous cell carcinoma, oesophageal cancer, oesophageal adenocarcinoma, oesophageal squamous cell carcinoma) or the therapies established for those diseases (*e.g.*, chemotherapy, nivolumab, pembrolizumab, camrelizumab, sintilimab, tislelizumab). We also set up filters for clinical trials. The search results from each database were combined in Endnote X9, and duplicates were removed. For the rest of the literature, we evaluate whether it is qualified by reading the full text. In addition, we also searched the relevant literature

manually to find other potentially relevant articles. The literature selection process was completed independently by two authors (Zhang XM and Yang T).

### ***Meta-analysis inclusion and exclusion criteria***

Eligible studies were selected following inclusion and exclusion criteria (Figure 1). A study was considered eligible for inclusion in the present meta-analysis if it met the following inclusion criteria: (1) The study was limited to oesophageal squamous cell carcinoma and/or G/GEJ adenocarcinoma; (2) The study was a phase III clinical trial; (3) The study included a sample of more than 500 patients; (4) The study encompassed at least one group receiving a combination of PD-1 inhibitor and chemotherapy, alongside one group receiving chemotherapy alone; (5) The study's basic data (*e.g.*, trial year, sample size, patient sex distribution, tumor types, trial abbreviation, registry number) were complete. Additionally, it was essential for the study to acquire the hazard ratio (HR) along with its corresponding 95% confidence interval (CI) for OS and PFS, as well as comparable data for objective response rate (ORR) and adverse events (AEs); and (6) The study was reported in English.

The exclusion criteria were as follows: (1) The literature type was meta-analysis, review or letter; (2) Lack of HR values and 95% CIs for OS and PFS comparisons between the two groups, or the number of patients who achieved ORRs and the number of AEs between the two groups were not reported; (3) The study was a non-randomized clinical trial; (4) The study concerned the results of radiotherapy; (5) The study included only results from special patient populations, such as elderly patients; and (6) Research lacking sufficient published data suitable for analysis.

### ***Data extraction***

Data were extracted independently by two of the authors (Zhang XM and Yang T). The extraction process encompassed the following items in accordance with the inclusion criteria: (1) General characteristics of the studies: Name of the first author, publication year, trial abbreviation, and registry number; (2) The patients' baseline characteristics:

Age, sex, sample size, tumor type, intervention arm, chemotherapy regimens, and PD-L1 combined positive scores (CPSs); and (3) Outcomes: We extracted the HRs and their 95% CIs for each study's OS, PFS, the number of patients who achieved an objective response, all AEs, grade > 3 AEs, and the counts of patients with a PD-L1 CPS of  $\geq 5$  or  $\geq 10$ .

### ***Quality assessment of the included studies***

We evaluated the potential bias in each study using Cochrane Collaboration's tool (Review Manager 5.3). The primary criteria for this assessment included random sequence generation, allocation concealment, participant blinding, outcome assessment blinding, incomplete outcome data, and selective reporting, as well as other biases. This assessment assigned grades according to levels of risk, *i.e.*, high, low, or unclear risk.

### ***Statistical analyses***

Stata ver. 12.0 software was used to perform all statistical tests. For the quantitative aggregation of the survival outcomes, the HRs and their 95% CIs were combined as the effective value. The significance of the pooled OR or HR was assessed using the Z-test, where  $P$ -values < 0.05 were deemed statistically significant. To evaluate the global heterogeneity between studies, we used the  $\chi^2$ -based Q-test ( $P > 0.10$  was considered a lack of heterogeneity) and the  $I^2$  test ( $I^2 \leq 50\%$  indicated low heterogeneity, while  $I^2 > 50\%$  indicated high heterogeneity). We employed the fixed-effects model if the heterogeneity was low; otherwise, we used the random-effects model. Funnel plots were generated to assess the presence of publication bias in the outcomes. An asymmetrical plot indicated possible publication bias. Review of the funnel plots could not rule out the potential for publication bias for the efficacy of PD-1 inhibitors in the treatment of UGT cancers. Further, asymmetry in the funnel plot was evaluated through Egger's test and Begg's test, with  $P$ -values < 0.05 indicating significance. If publication bias existed, we assessed the effect of publication bias by using the trim-fill method. Additionally, sensitivity analyses were conducted to assess the stability of the



results. The statistical methods of this study were reviewed by Guo-Lin Han from Medical records statistics room of Changzhi People's Hospital.

## **RESULTS**

### ***Search results***

A total of 310 articles were retrieved from the PubMed and Embase databases; one duplicate paper was removed, leaving 309 articles remaining. We initially screened the articles according to their titles and abstracts. As a result, we excluded 254 articles as irrelevant and 15 as either meta-analyses, case reports, guidelines, or reviews. We then screened the remaining 41 articles by reading them in full, resulting in the exclusion of 22 articles with the same trial registration number, 3 incomplete clinical trials, and 7 articles lacking the OS, PFS, ORR, and/or AE data required by this study. The final total was 9 articles<sup>[10-18]</sup>. Figure 1 is a flow diagram illustrating the inclusion and exclusion of articles.

### ***Study and patient characteristics***

All 9 of the articles included in this meta-analysis were reports of phase 3 clinical trials. Among them, 8 were multicenter studies and the remainder was a single-center study. The 9 articles were published between 2020 and 2022 and included a total of 7016 patients, with sample sizes ranging from 596 to 1581. Three articles compared nivolumab in combination with chemotherapy *vs* chemotherapy alone, while two articles compared pembrolizumab + chemotherapy *vs* chemotherapy alone. Two articles examined sintilimab + chemotherapy *vs* chemotherapy alone, and the remaining two articles investigated chemotherapy alone *vs* camrelizumab and tislelizumab, respectively. Table 1 summarizes the patients' clinical characteristics.

Next, we evaluated the risk of bias for each study, and the outcomes are illustrated in Figure 2. One of the sources of bias originated from the absence of blinding, which notably impacted the study quality. The precision of results in another study could have been influenced due to participant withdrawals or losses. Additional biases could have

arisen from factors such as an uneven geographic distribution of participants, poor or incomplete patient compliance, or other underlying reasons.

### ***Primary outcomes (OS and PFS)***

All 9 of the clinical trials provided HR and 95%CI values for the OS and PFS outcomes of patients treated with a PD-1 inhibitor + chemotherapy compared to chemotherapy alone. We pooled the HR values from the 9 clinical trials using randomized- or fixed-effects models, and the results showed that the pooled PD-1 inhibitor + chemotherapy group had longer OS than the pooled chemotherapy-alone group (HR = 0.76, 95%CI: 0.71-0.81). Figure 3A shows the combined forest plot of the OS data. The combined outcome for patients' PFS corresponded with that of their OS (HR = 0.67, 95%CI: 0.61-0.74), as depicted in Figure 3B.

### ***Secondary outcomes: ORR and AEs***

Of the 9 studies, 8 provided ORR data, and we gathered the count of patients who obtained an objective response in each study. The findings suggested that the count of individuals achieving an objective response in the PD-1 inhibitor + chemotherapy group exceeded that in the chemotherapy-alone group (OR = 1.86, 95%CI: 1.59-2.18) (Figure 3C). Regarding the AEs observed in the studies, we aggregated both the AEs of all grades and those rated  $\geq$  grade 3. As shown in Figure 3D, the meta-analysis of AEs indicated that the frequency of AEs in the PD-1 inhibitor + chemotherapy group exceeded that in the chemotherapy-alone group. This held true for AEs of all grades (OR = 1.88, 95%CI: 1.39-2.54) as well as for AEs rated  $\geq$  grade 3 (OR = 1.30, 95%CI: 1.07-1.57) in Figure 4C.

### ***Subgroup analyses of OS and PFS***

We employed the patients' PD-L1 CPS scores as a measure to assess the efficacy of immunotherapy against tumors. Of the 9 clinical trials, 5 reported treatment outcomes in different PD-L1 CPS subgroups. Consequently, we conducted a subgroup analysis of

OS and PFS based on the patients' PD-L1 CPS scores. The subgroup analysis of OS (Figure 4A) indicated that patients treated with a PD-1 inhibitor + chemotherapy exhibited prolonged OS compared to those treated with chemotherapy alone, irrespective of the subgroup CPS  $\geq 1$  (HR = 0.73, 95%CI: 0.60-0.90,  $P = 0.002$ ), CPS  $\geq 5$  (HR = 0.68, 95%CI: 0.60-0.78,  $P < 0.001$ ), or CPS  $\geq 10$  (HR = 0.68, 95%CI: 0.57-0.82,  $P < 0.001$ ). The outcomes of our subgroup analysis for PFS (Figure 4B) aligned with those for OS across all three groups: CPS  $\geq 1$  (HR = 0.70, 95%CI: 0.56-0.88,  $P = 0.002$ ), CPS  $\geq 5$  (HR = 0.63, 95%CI: 0.56-0.72,  $P < 0.001$ ), and CPS  $\geq 10$  (HR = 0.54, 95%CI: 0.46-0.64,  $P < 0.001$ ).

### ***Publication bias***

To minimize the effects of publication bias, we expanded database searches and performed a thorough search for unpublished studies, and used funnel plot to quantify the potential presence of publication bias. We utilized funnel plots of the OS and PFS data to evaluate potential publication bias (Supplementary Figure 1). The outcomes of the Begg's test (OS:  $P = 0.25$ ; PFS:  $P = 0.532$ ) and Egger's test (OS:  $P = 0.336$ ; PFS:  $P = 0.512$ ) for both the OS and PFS funnel plots indicated the absence of substantial publication bias in this meta-analysis.

### ***Sensitivity analysis***

The Stata sensitivity analysis (Supplementary Figure 2) revealed that the exclusion of any of the 9 articles had a minimal impact on the overall effect size. This suggested that the sensitivity of this meta-analysis was low and that the removal of a single article did not significantly alter the results. In order to further explore the source of heterogeneity of OS and PFS, we used meta-regression to analyze whether the year of publication, sample size, and tumor type were the source of heterogeneity of OS and PFS. The results of meta-regression showed that the year of publication, sample size, and tumor type were not sources of heterogeneity in OS (year of publication:  $P = 0.167$ ; sample

size:  $P = 0.233$ ; tumor type:  $P = 0.247$ ) and PFS (year of publication:  $P = 0.574$ ; sample size:  $P = 0.609$ ; tumor type:  $P = 0.914$ ).

## **DISCUSSION**

The combination of ICIs with chemotherapy has emerged as a potent targeted therapy for advanced or metastatic gastrointestinal tumors. Across various studies, the utilization of a PD-1 inhibitor + chemotherapy notably enhanced both patients' OS and PFS, including the PD-1 inhibitors nivolumab<sup>[12]</sup>, pembrolizumab<sup>[10]</sup>, camrelizumab<sup>[13]</sup>, and sintilimab<sup>[14]</sup>. Guo *et al*<sup>[19]</sup> performed an independent meta-analysis concerning PD-1 inhibitor + oxaliplatin or cisplatin-based chemotherapy. Thus far, however, there hasn't been a meta-analysis comparing the effectiveness of PD-1 inhibitors + chemotherapy and chemotherapy alone. We therefore conducted a meta-analysis to assess and contrast the efficacy and safety of PD-1 inhibitors + chemotherapy *vs* chemotherapy alone.

Several clinical trials of PD-1 inhibitors in combination with chemotherapy *vs* chemotherapy alone have been completed or are under way. In terms of efficacy, however, it remains unknown whether a PD-1 inhibitor + chemotherapy can significantly prolong patients' OS and PFS compared with chemotherapy alone. CheckMate 649<sup>[12]</sup> is a multicenter, randomized, phase 3 trial that compared nivolumab + chemotherapy *vs* chemotherapy alone as a first-line treatment for advanced gastric cancer, GEJ cancer, and oesophageal adenocarcinoma. The results revealed that nivolumab + chemotherapy substantially extended patients' OS (HR = 0.8,  $P < 0.0002$ ) and PFS (HR = 0.8,  $P < 0.0002$ ) in comparison to chemotherapy alone. Nevertheless, in the KEYNOTE-062 trial<sup>[10]</sup>, which compared pembrolizumab + chemotherapy to chemotherapy alone, the combination of pembrolizumab + chemotherapy did not extend patients OS compared to chemotherapy alone (HR = 0.85,  $P = 0.05$ ). Similar results were obtained in the ATTRACTION-4 trial<sup>[11]</sup> (nivolumab + chemotherapy *vs* placebo + chemotherapy) (HR = 0.90,  $P = 0.26$ ). Our present meta-analysis of 9 clinical trials revealed that treatment with a PD-1 inhibitor + chemotherapy significantly

<sup>3</sup> prolonged both OS (HR = 0.76, 95%CI: 0.71-0.81) and PFS (HR = 0.68, 95%CI: 0.64-0.72) in patients with gastric cancer, GEJ cancer, or oesophageal cancer.

The ORR is the proportion of patients who respond to treatment, which is a key measure of a drug therapy's effectiveness. The results of our meta-analysis demonstrated that the PD-1 inhibitor + chemotherapy group had superior ORR to that of the chemotherapy-alone group. We also conducted an analysis of the incidence of AEs between the two treatment approaches. This revealed a heightened occurrence of AEs in patients receiving immunotherapy combined with chemotherapy compared to those undergoing chemotherapy alone. This trend was consistent whether we examined AEs of all grades (OR = 1.88, 95%CI: 1.39-2.54) or evaluated AEs rated  $\geq$  grade 3 (HR = 1.30, 95%CI: 1.07-1.57). The AEs of immunotherapy are referred to as immune-related (ir)AEs. These irAEs have been documented to be generally mild and manageable, often not necessitating specific interventions. More importantly, they are noted to have minimal impact on treatment efficacy<sup>[20,21]</sup>.

PD-L1 expression levels are frequently employed as predictive biomarkers to ascertain whether patients are likely to derive benefits from immunotherapy. The CPS score has been indicated to be a more effective method of assessing PD-L1 expression than the tumor proportion score (TPS)<sup>[22]</sup>. In the present meta-analysis, we therefore conducted subgroup analyses of OS and PFS based on the CPS scores for PD-L1. The outcomes demonstrated that within the subgroup with a CPS  $\geq$  5 or CPS  $\geq$  10, patients in the immunotherapy + chemotherapy group exhibited longer OS periods. The PFS subgroup analysis revealed that the patients in the subgroup with a CPS  $\geq$  10 were more likely to benefit from immunotherapy + chemotherapy. Based on the results of our analysis, PD-1 inhibitor + chemotherapy can be recommended as a suitable first-line treatment for patients with PD-L1 CPS scores of  $\geq$  5 or  $\geq$  10. This study has two limitations. First, not all of the included articles provided survival data for CPS subgroups, opening the possibility of bias in our CPS subgroup analysis. Second, the limited number of articles focused on the same tumor type precluded the possibility of conducting a subgroup analysis based on tumor type.

## **CONCLUSION**

The outcomes of this meta-analysis illustrated that patients receiving a PD-1 inhibitor + chemotherapy experienced notably prolonged OS and PFS compared to patients treated with chemotherapy alone. The ORR and the rate of AEs were both higher in the group treated with a PD-1 inhibitor + chemotherapy than in the chemotherapy-alone group. For the subgroups with a CPS  $\geq 5$  or CPS  $\geq 10$ , the patients in the immunotherapy + chemotherapy group had longer OS and PFS. Considering both efficacy and safety, PD-1 inhibitor + chemotherapy could potentially emerge as a preferable first-line treatment option for patients diagnosed with gastric cancer, GEJ tumor, or esophageal cancer, particularly for those with a PD-L1 CPS score of  $\geq 5$  or  $\geq 10$ .

## ORIGINALITY REPORT

# 4%

SIMILARITY INDEX

### PRIMARY SOURCES

- |          |   |                 |
|----------|---|-----------------|
| <b>1</b> | <a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a><br><small>Internet</small>   | 45 words — 1%   |
| <hr/>    |   |                 |
| <b>2</b> | <a href="http://www.omicsdi.org">www.omicsdi.org</a><br><small>Internet</small>   | 23 words — 1%   |
| <hr/>    |   |                 |
| <b>3</b> | <a href="#">Jiwei Huang, Yueming Wang, Haoran Zhang, Xiaoyi Hu et al. "Clinical outcomes of second-line treatment following first-line VEGFR-TKI failure in patients with metastatic renal cell carcinoma: a comparison of axitinib alone and axitinib plus anti-PD-1 antibody", Cancer Communications, 2021</a><br><small>Crossref</small> | 20 words — 1%   |
| <hr/>    |   |                 |
| <b>4</b> | <a href="#">Zanzan Wang, Shuna Luo, Xiaoying Zhao. "The Prognostic Impact of Body Mass Index in Patients with Diffuse Large B-Cell Lymphoma: A Meta-Analysis", Nutrition and Cancer, 2020</a><br><small>Crossref</small>  | 20 words — 1%   |
| <hr/>    |   |                 |
| <b>5</b> | <a href="http://cdn.techscience.cn">cdn.techscience.cn</a><br><small>Internet</small>   | 18 words — < 1% |
| <hr/>    |   |                 |
| <b>6</b> | <a href="http://bmjopen.bmj.com">bmjopen.bmj.com</a><br><small>Internet</small>   | 16 words — < 1% |

EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES

< 15 WORDS