

22

Name of Journal: *World Journal of Gastrointestinal Oncology*

ESPS Manuscript NO: 21417

Manuscript Type: ORIGINAL ARTICEL

Basic Study
 Plasma chitinase 3-like 1 is ¹⁸persistently elevated during first month after minimally invasive colorectal cancer resection

 HMC ³Shantha Kumara, David Gaita, Hiromichi Miyagaki, Xiaohong Yan, Sonali AC Hearsh, Linda Njoh, Vesna Cekic, Richard L Whelan
Abstract
 AIM: To assess blood chitinase 3-like 1 (CHI3L1) levels for 2 mo ¹²after minimally invasive colorectal resection (MICR) for colorectal cancer (CRC). ⁷

 Method: CRC Patients in an Institutional review Board approved data/plasma bank who underwent elective MICR for whom Preoperative (PreOp), early postoperative (postop) and 1 or more late postop samples (POD 7-27) available were included. Plasma ¹⁰CHI3L1 levels (ng/ml) were determined in duplicate by ELISA.

 Results: PreOp and Postop plasma sample were available for 80 MICR cancer patients ⁸for the study. The median PreOp CHI3L1 level was 56.8 CI 41.9- 78.6 ng/ml (n=80). Significantly elevated (p<0.001) median plasma levels (ng/ml) over PreOp levels were detected on POD1 (667.7 CI: 495.7, 771.7; n=79), POD 3 (132.6 CI: 95.5,173.7 ; n=76),

Match Overview

Rank	Source	Words	Similarity
1	CrossCheck	94 words	3%
	Shantha Kumara, H. M. C., Elizabeth A. Myers, Sonali A. C. Herath, Linda Njoh, Xiaohong Yan, Daniel Kirchoff, Nadav		
2	CrossCheck	70 words	2%
	Kumara, H. M. C. S., S. T. Tohme, I. Y. Kim, D. G. Kim, M. F. Kalady, M. Luchtefeld, K. Hoffman, V. DiMaggio, and R.		
3	CrossCheck	70 words	2%
	"2013 Scientific Session of the Society of American Gastrointestinal and Endoscopic Surgeons (SAGES) Baltimore,		
4	CrossCheck	58 words	2%
	H. M. C. Shantha Kumara. "Plasma levels of angiopoietin-like protein 4 (ANGPTL4) are significantly lower preoperativ		
5	CrossCheck	53 words	2%
	H. M. C. Shantha Kumara. "Plasma soluble vascular adhesion molecule-1 levels are persistently elevated during the...		
6	CrossCheck	53 words	2%
	Michael J. Grieco. "Minimally invasive colorectal resection is associated with a rapid and sustained decrease in plas...		



全部

图片

新闻

视频

更多 ▾

搜索工具

已启用

找到约 1,870 条结果 (用时 0.80 秒)

Plasma Levels of Human Chitinase 3-like 1 (chi3l1), a Proangiogenic ...

www.sages.org/.../plasma-levels-of-human-chitinase-3-like-1-chi3l1-a-proan... ▾ 翻译此页

Plasma Levels of Human Chitinase 3-like 1 (chi3l1), a Proangiogenic Protein, Are ... During the First Month After Minimally Invasive Colorectal Cancer Resection ... Persistently elevated CHI3L1 levels, together with the similarly increased ...

Xiaohong Yan - Publications - ResearchGate

https://www.researchgate.net/profile/Xiaohong_Yan/publications ▾ 翻译此页

Article: Minimally invasive colorectal resection is associated with significantly elevated levels of plasma matrix metalloproteinase 3 (MMP-3) during the first month after ... Colorectal Resection for Benign Pathology Is Associated With Persistent ... Cancer Is Associated With Elevated Plasma Levels of Chitinase 3-Like-1.

Richard L Whelan - Publications - ResearchGate

https://www.researchgate.net/profile/Richard_Whelan/publications ▾ 翻译此页

Article: Minimally invasive colorectal resection is associated with significantly elevated levels of plasma matrix metalloproteinase 3 (MMP-3) during the first ...

Chandana M C Shantha Kumara - Publications - ResearchGate

https://www.researchgate.net/profile/Chandana_Kumara/publications ▾ 翻译此页

Article: Minimally invasive colorectal resection is associated with significantly ... Cancer Is Associated With Elevated Plasma Levels of Chitinase 3-Like-1 ... levels are persistently elevated during the first month after colorectal cancer resection.

Reduced NGF Secretion by HT-29 Human Colon Cancer Cells ...

connection.ebscohost.com/.../reduced-ngf-secretion-by-ht-29-human-colon-... ▾ 翻译此页

Chitinase 3-like 1 promotes macrophage recruitment and angiogenesis in ... levels are persistently elevated during the first month after colorectal cancer resection. ... Plasma from the second and third

[全部](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 1,870 条结果 (用时 0.80 秒)

Plasma Levels of Human Chitinase 3-like 1 (chi3l1), a Proangiogenic ...

www.sages.org/.../plasma-levels-of-human-chitinase-3-like-1-chi3l1-a-proan... ▾ [翻译此页](#)

Plasma Levels of Human Chitinase 3-like 1 (chi3l1), a Proangiogenic Protein, Are ... During the First Month After Minimally Invasive Colorectal Cancer Resection ... Persistently elevated CHI3L1 levels, together with the similarly increased ...

Xiaohong Yan - Publications - ResearchGate

https://www.researchgate.net/profile/Xiaohong_Yan/publications ▾ [翻译此页](#)

Article: Minimally invasive colorectal resection is associated with significantly elevated levels of plasma matrix metalloproteinase 3 (MMP-3) during the first month after ... Colorectal Resection for Benign Pathology Is Associated With Persistent ... Cancer Is Associated With Elevated Plasma Levels of Chitinase 3-Like-1.

Richard L Whelan - Publications - ResearchGate

https://www.researchgate.net/profile/Richard_Whelan/publications ▾ [翻译此页](#)

Article: Minimally invasive colorectal resection is associated with significantly elevated levels of plasma matrix metalloproteinase 3 (MMP-3) during the first ...

Chandana M C Shantha Kumara - Publications - ResearchGate

https://www.researchgate.net/profile/Chandana_Kumara/publications ▾ [翻译此页](#)

Article: Minimally invasive colorectal resection is associated with significantly ... Cancer Is Associated With Elevated Plasma Levels of Chitinase 3-Like-1 levels are persistently elevated during the first month after colorectal cancer resection.

Reduced NGF Secretion by HT-29 Human Colon Cancer Cells ...

connection.ebscohost.com/.../reduced-ngf-secretion-by-ht-29-human-colon-... ▾ [翻译此页](#)

Chitinase 3-like 1 promotes macrophage recruitment and angiogenesis in ... levels are persistently elevated during the first month after colorectal cancer resection. ... Plasma from the second and third



全部 图片 新闻 视频 更多 ▾ 搜索工具

找到约 1,830 条结果 (用时 0.91 秒)

Plasma Levels of Human Chitinase 3-like 1 (chi3l1), a Proangiogenic ...

www.sages.org/.../plasma-levels-of-human-chitinase-3-like-1-chi3l1-a-proan... ▾ 翻译此页

Plasma Levels of Human Chitinase 3-like 1 (chi3l1), a Proangiogenic Protein, Are ... During the First Month After Minimally Invasive Colorectal Cancer Resection ... Persistently elevated CHI3L1 levels, together with the similarly increased ...

Xiaohong Yan - Publications - ResearchGate

https://www.researchgate.net/profile/Xiaohong_Yan/publications ▾ 翻译此页

Article: Minimally invasive colorectal resection is associated with significantly elevated levels of plasma matrix metalloproteinase 3 (MMP-3) during the first month after ... Colorectal Resection for Benign Pathology Is Associated With Persistent ... Cancer Is Associated With Elevated Plasma Levels of Chitinase 3-Like-1.

Chandana M C Shantha Kumara - Publications - ResearchGate

https://www.researchgate.net/profile/Chandana_Kumara/publications ▾ 翻译此页

Article: Minimally invasive colorectal resection is associated with significantly ... Cancer Is Associated With Elevated Plasma Levels of Chitinase 3-Like-1 levels are persistently elevated during the first month after colorectal cancer resection.

Richard L Whelan - Publications - ResearchGate

https://www.researchgate.net/profile/Richard_Whelan/publications ▾ 翻译此页

Article: Minimally invasive colorectal resection is associated with significantly elevated levels of plasma matrix metalloproteinase 3 (MMP-3) during the first ...

Express Submission and Peer-review System

www.wjgnet.com/esps/ManuscriptList.aspx ▾ 翻译此页

Manuscript title: Plasma chitinase 3-like 1 is persistently elevated during first month after minimally