

**Name of journal:** *World Journal of Gastrointestinal Oncology*

**ESPS Manuscript NO:** 24692

**Manuscript Type:** Original Article

*Randomized Clinical Trial*

**Utility of different serum fibrosis markers in diagnosis patients with chronic pancreatitis and pancreatic adenocarcinoma**

**Anna Kozak, Renata Talar-Wojnarowska, Aleksandra Kaczka, Anna Borkowska, Leszek Czupryniak, Ewa Małecka-Panas, Anita Gąsiorowska**

## Match Overview

There are no matching sources for this report.

Utility of serum fibrosis markers: soluble type fractalkine, hyaluronic acid, monc

全部

图片

视频

新闻

购物

地图

图书

找到约 1,230 条结果

时间不限

过去 1 小时内

过去 24 小时内

过去 1 周内

过去 1 个月内

过去 1 年内

所有结果

精确匹配

### [TGFB1 transforming growth factor beta 1 \[ \(human\)\]](#)

[www.ncbi.nlm.nih.gov/gene/7040](http://www.ncbi.nlm.nih.gov/gene/7040)

This gene encodes a member of the **transforming growth factor beta** (TGFB) ... the elevated **serum TGF-beta1** and IL-21 levels could contribute to enhanced ... in the peripheral blood may be used as a molecular **marker** for the **diagnosis** of HCC. .... High TGFbeta expression promotes **pancreatic ductal adenocarcinoma** ...

### [VEGFA vascular endothelial growth factor A \[ \(human\)\]](#)

[www.ncbi.nlm.nih.gov/gene/7422](http://www.ncbi.nlm.nih.gov/gene/7422)

This gene is a member of the PDGF/VEGF **growth factor** family. ... Elevated levels of this **protein** are found in **patients** with PQEMS syndrome. .... **1** and VEGF-A is a potential prognostic **marker** for gallbladder **carcinoma patients**. .... niche and angiogenic **factors**, particularly VEGFA, during **pancreatic beta** cell development.

### [CD44 CD44 molecule \(Indian blood group\) \[ \(human\)\]](#)

[www.ncbi.nlm.nih.gov/gene/960](http://www.ncbi.nlm.nih.gov/gene/960)

24 Jan 2016 ... The **protein** encoded by this gene is a cell-surface glycoprotein involved in ... It is a receptor for **hyaluronic acid** (HA) and can also interact with other ... of cells that highly express CD44 as an **early marker** for neoplastic stem cell .... c-Met and CD44v6 receptors have roles in autocrine **TGF-beta1** signaling in ...

Utility of different serum fibrosis markers in diagnosis patients with chronic pa

全部

图片

视频

新闻

购物

地图

图书

找到约 3,570,000 条结果

时间不限

过去 1 小时内

过去 24 小时内

过去 1 周内

过去 1 个月内

过去 1 年内

所有结果

精确匹配

[Chronic Pancreatitis. Hereditary pancreas disorders ... - Patient](#)  
[patient.info/doctor/chronic-pancreatitis-pro](#) ▼

Chronic pancreatitis is the result of chronic inflammation of the pancreas which ... in patients with recurrent attacks of acute pancreatitis associated with alcohol use .... On the other hand, small duct pancreatitis occurs more in women and is not .... Serum nutritional markers are being evaluated as potential diagnostic factors.

[The clinical utility of serum CA 19-9 in the diagnosis, prognosis and ...](#)  
[www.ncbi.nlm.nih.gov/pmc/articles/PMC3397644/](#) ▼

2 Mar 2011 ... Key Words : Pancreatic cancer, tumor markers, CA 19-9, diagnosis, ... the tissue and sera of patients with other gastrointestinal tumors including esophageal, ... such as acute and chronic pancreatitis, intraductal pancreatic mucinous ... CA 19- 9 serum levels were elevated in 61% of benign causes and 86% ...

[The Clinical Utility of CA 19-9 in Pancreatic Adenocarcinoma - NCBI](#)  
[www.ncbi.nlm.nih.gov/pmc/articles/PMC4419808/](#) ▼

See other articles in PMC that cite the published article. ... CA 19-9 is also not specific to pancreatic cancer as it can be elevated in extra-pancreatic ... (89%) included chronic pancreatitis, the most common confounding diagnosis for patients .... CA 19-9 is a useful serum marker to predict tumor stage and resectability, overall ...



[全部](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 3,490,000 条结果 (用时 0.88 秒)

## Google 学术: Utility of different serum fibrosis markers in diagnosis patients with chronic pancreatitis and pancreatic adenocarcinoma

... pancreatitis and in distinguishing it from pancreatic ... - Ghazale - 被引用次数: 441

... autoimmune pancreatitis from pancreatic cancer - Chari - 被引用次数: 239

... diagnostic criteria for autoimmune pancreatitis: ... - Shimosegawa - 被引用次数: 616

## Chronic Pancreatitis. Hereditary pancreas disorders information | Patient patient.info/doctor/chronic-pancreatitis-pro ▾ 翻译此页

Chronic pancreatitis is the result of chronic inflammation of the pancreas which ... in patients with recurrent attacks of acute pancreatitis associated with alcohol use. ... On the other hand, small duct pancreatitis occurs more in women and is not .... Serum nutritional markers are being evaluated as potential diagnostic factors.

## Hybrid kappa\lambda antibody is a new serological marker to ...

www.nature.com > Scientific Reports > Articles > Article - 翻译此页

4 天前 - Autoimmune pancreatitis (AIP) is a form of chronic pancreatitis ... infiltration and fibrosis in the pancreas are the gold standard for the diagnosis of AIP. ... of AIP is pancreatic cancer, we also aimed to evaluate the use of serum hybrid κλ ... High prevalence of serum hybrid κλ antibodies in patients with AIP in ...

## The clinical utility of serum CA 19-9 in the diagnosis, prognosis and ...

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) ▾ 翻译此页

作者: UK Ballehaninna - 2012 - 被引用次数: 154 - 相关文章

2011年3月2日 - Key Words : Pancreatic cancer, tumor markers, CA 19-9, diagnosis, ... the tissue and sera of patients with other gastrointestinal tumors including esophageal, ..... such as acute and chronic