

## Match Overview

1	Crossref 50 words Kang, M.J., "Venous thromboembolism (VTE) in patients with advanced gastric cancer: An Asian experience", <i>Europea</i>	2%
2	Crossref 18 words Sud, R., "Cancer-associated thrombosis: risk factors, candidate biomarkers and a risk model", <i>Thrombosis Research</i>	1%
3	Internet 18 words crawled on 17-May-2016 <a href="http://meetinglibrary.asco.org">meetinglibrary.asco.org</a>	1%
4	Crossref 13 words "Epidemiology of venous thromboembolism in 9489 patients with malignant glioma", <i>Journal of Neurosurgery</i> , 04/200	<1%
5	Crossref 13 words Kourlaba, Georgia, John Relakis, Charalambos Mylonas, Vasiliki Kapaki, Stathis Kontodimas, Madbrit V. Holm, and Ni	<1%
6	Internet 13 words crawled on 01-Nov-2016 <a href="http://e-sciencecentral.org">e-sciencecentral.org</a>	<1%

Name of journal: *World Journal of Gastrointestinal Oncology*

ESPS Manuscript NO: 31760

Manuscript Type: Original Article

*Prospective Study*

Incidence of venous thromboembolism and the role of D-dimer as predictive marker in patients with advanced gastric cancer receiving chemotherapy: A prospective study

Kwonoh Park, Baek-Yeol Ryoo, Min-Hee Ryu, Sook Ryun Park, Myoung Joo Kang, Jeong Hye Kim, Seungbong Han, Yoon-Koo Kang

Abstract

Incidence of venous thromboembolism and the role of D-dimer as predictive marl

全部 图片 视频 新闻 地图 图书

找到约 4,580 条结果

时间不限

过去 1 小时内  
过去 24 小时内  
过去 1 周内  
过去 1 个月内  
过去 1 年内

所有结果

精确匹配

### [The Role of Thromboprophylaxis in Cancer Patients: Emerging Data](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3088308/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3088308/>

Risk for VTE is influenced by patient-, cancer- and treatment-specific factors. ... of patients with metastatic gastric and gastro-esophageal junction cancers. ... In a prospective study of cancer patients receiving chemotherapy, rates of VTE ... role of D-dimer levels prior to chemotherapy or major surgery in predicting risk for VTE ...

### [Treating venous thromboembolism in patients with cancer - NCBI - NIH](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4435973/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4435973/>

In the hospitalized setting, the rate of VTE in cancer patients is twice that of non- cancer ... for predicting chemotherapy-associated VTE in ambulatory cancer patients was ... laboratory markers, soluble P-selectin (sP-selectin) and D-dimer ( Table 1). .... These varying results underscore the importance of prospective studies ...

### [Biomarkers for prediction of venous thromboembolism in cancer ...](http://www.bloodjournal.org/content/122/12/2011)

[www.bloodjournal.org/content/122/12/2011](http://www.bloodjournal.org/content/122/12/2011)

The incidence among different groups of cancer patients varies considerably ... The role of high platelet count in the occurrence of VTE is supported by several studies. ... Prospective studies investigating potential predictive biomarkers of VTE in ... d-Dimer may even be a valuable biomarker for prediction of recurrent VTE in ...

### [Biomarkers predictive of venous thromboembolism in patients with ...](http://neuro-oncology.oxfordjournals.org/content/16/12/1645.full)

[neuro-oncology.oxfordjournals.org/content/16/12/1645.full](http://neuro-oncology.oxfordjournals.org/content/16/12/1645.full)

1 Jul 2014 ... VTE incidence rates in HGG patients differ widely, but rates between 7% and 24 % ... In previous prospective studies including patients with various .... Initially, D- dimer was a strong VTE predictor, but it did not prove to be ..... Semuloparin for thromboprophylaxis in patients receiving chemotherapy for cancer.

### [D-Dimer Can Serve as a Prognostic and Predictive Biomarker f...](#)



[全部](#)[新闻](#)[图片](#)[购物](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 5,190 条结果 (用时 0.96 秒)

## The Role of Thromboprophylaxis in Cancer Patients: Emerging Data

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: LA Menapace - 2010 - 被引用次数: 16 - 相关文章

Risk for VTE is influenced by patient-, cancer- and treatment-specific factors. ... of patients with metastatic gastric and gastro-esophageal junction cancers, ... In a prospective study of cancer patients receiving chemotherapy, rates of VTE ... role of D-dimer levels prior to chemotherapy or major surgery in predicting risk for VTE ...

## Treating venous thromboembolism in patients with cancer - NCBI

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: C Piatek - 2012 - 被引用次数: 16 - 相关文章

In the hospitalized setting, the rate of VTE in cancer patients is twice that of non-cancer ... for predicting chemotherapy-associated VTE in ambulatory cancer patients was ... laboratory markers, soluble P-selectin (sP-selectin) and D-dimer (Table 1). .... These varying results underscore the importance of prospective studies ...

## Venous thromboembolism and lung cancer: a review

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: C Vitale - 2015 - 被引用次数: 5 - 相关文章

2015年9月15日 - The incidence of VTE in lung cancer patients has been described in several studies. ... Several studies have focused on the role of Tissue Factor (TF) in the .... Although high D-dimer was described as a predictor of VTE in earlier .... in patients with NSCLC at all stages who receive chemotherapy, but not ...

## Assessing Risk of Venous Thromboembolism in the Patient With Cancer

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: AA Khanna - 2009 - 被引用次数: 264 - 相关文章