



34624-Review.doc

Quotes Excluded
Bibliography Excluded

4%
SIMILAR

Name of Journal: *World Journal of Gastrointestinal Oncology*

Manuscript NO: 34624

Manuscript Type: Original Article

Prospective Study

Polyethylene glycol microspheres loaded with irinotecan for arterially directed embolic therapy of metastatic liver cancer

Giammaria Fiorentini

Abstract

AIM: to study tumor response, and tolerability of arterially directed embolic therapy (ADET) with polyethylene glycol embolics loaded with irinotecan for the treatment of colorectal cancer liver metastases (CRC-LM). Secondary objectives were to monitor quality of life, time to progression and survival of patients.

METHODS: patients were included in the study if they were affected by CRC-LM,

Match Overview

1	Internet 32 words crawled on 19-Dec-2015 office.wjgnet.com	1%
2	Internet 26 words crawled on 06-Apr-2017 ar.iiarjournals.org	1%
3	Internet 16 words crawled on 25-Apr-2013 www.131.org.cn	1%
4	Internet 12 words crawled on 31-Mar-2017 www.codental.uobaghdad.edu.iq	<1%
5	Internet 10 words crawled on 08-Apr-2014 www.journalsleep.org	<1%
6	Crossref 10 words on behalf of the United Kingdom Children's Cancer Study Group (UKCCSG) Brain Tumor Committee. "Outcome for patient	<1%

找到约 3,980 条结果 (用时 0.74 秒)

Trans-arterial chemoembolization of metastatic colorectal carcinoma ...

<https://www.ncbi.nlm.nih.gov/pubmed/22199334> - 翻译此页

作者: C Aliberti - 2011 - 被引用次数: 52 - 相关文章

Trans-arterial **chemoembolization of metastatic** colorectal carcinoma to the liver adopting DC Bead®, drug-eluting bead **loaded with irinotecan**: results of a ... locoregional therapy for the treatment of primary **hepatic tumors** and liver **metastases**. ... bead, a spherical **embolic** device capable of being **loaded with irinotecan**.

缺少字词: polyethylene glycol

Transcatheter embolization therapy in liver cancer: an update of ...

[https://www.ncbi.nlm.nih.gov/NCBI/Literature/PubMedCentral\(PMC\)](https://www.ncbi.nlm.nih.gov/NCBI/Literature/PubMedCentral(PMC)) - 翻译此页

作者: YXJ Wang - 2015 - 被引用次数: 33 - 相关文章

Chemoembolization and radioembolization are at the core of the treatment of liver ... The key clinical trials of transcatheter arterial therapy for **liver cancer** are summarized. ... Liver **metastases** can be found in 40% to 70% of patients with colorectal Poly (**ethylene glycol**) methacrylate (PEGMA) hydrolyzable microspheres ...

Intra-arterial Hepatic Chemoembolization (TACE) of Liver Metastases ...

iv.iiarjournals.org/content/23/1/131.abstract - 翻译此页

作者: G FIORENTINI - 2009 - 被引用次数: 50 - 相关文章

Intra-arterial Hepatic **Chemoembolization (TACE)** of Liver **Metastases** from ... DC Beads are new **embolic** products that can be **loaded with irinotecan (IRI)**. ... Hepatic Arterial Infusion of **Polyethylene Glycol** Drug-eluting Beads for Primary and **Metastatic Liver Cancer** Therapy Anticancer Res July 1, 2016 36 (7) 3515-3521.

LifePearl® - Microspheres - Terumo Europe

www.terumo-europe.com/en-emea/...tace/lifepearl-microspheres ▼ 翻译此页

LifePearl are **PEG (polyethylene glycol)** embolization microspheres that can be **loaded** with ... vascular

[全部](#)[图片](#)[购物](#)[新闻](#)[视频](#)[更多](#)[设置](#)[工具](#)

找到约 30,300 条结果 (用时 0.70 秒)

Google 学术 : Polyethylene glycol microspheres loaded with irinotecan for arterially directed embolic therapy of metastatic liver cancer

... and leucovorin for metastatic colorectal cancer - Saltz - 被引用次数 : 3253

... (TACE) for colorectal liver metastases—current status ... - Massmann - 被引用次数 : 8

... from hydrophilic resorbable embolization microspheres ... - Bédouet - 被引用次数 : 7

Trans-Arterial Chemoembolization of Metastatic Colorectal Carcinoma ...

ar.iiarjournals.org/content/31/12/4581.full - 翻译此页

作者 : C ALIBERTI - 2011 - 被引用次数 : 55 - 相关文章

Metastatic colorectal cancer (CRC) is one of the leading causes of ... Surgical resection is the standard treatment in patients with resectable liver disease but ... DC Bead, drug-eluting bead, a spherical embolic device capable of loading with ... Eighty-one DEBIRI with 2 ml of beads loaded with 100 mg of irinotecan and 104 ...

Non-operative therapies for colorectal liver metastases - NCBI - NIH

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者 : JL Noshier - 2015 - 被引用次数 : 12 - 相关文章

The armamentarium for liver-directed therapy includes ablative therapies, ... Keywords: Colon cancer, liver metastases, embolization, radiation, ablation ... Drug eluting beads loaded with irinotecan (DEBIRI) provide an alternative for ... about 80% of liver tumors, the Yttrium-90 microspheres preferentially flow to the tumors.

缺少字词 : polyethylene glycol

Intra-Arterial Therapies for Metastatic Colorectal Cancer - NCBI - NIH

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者 : DS Wang - 2013 - 被引用次数 : 11 - 相关文章

Intra-arterial therapies for unresectable hepatic metastases from colorectal cancer ... There therefore remains a need for liver-directed locoregional therapies for ... or drug-eluting beads loaded with

Polyethylene glycol microspheres loaded with irinotecan for arterially directed em

全部

图片

视频

新闻

地图

图书

找到约 30,900 条结果

时间不限

过去 1 小时内

过去 24 小时内

过去 1 周内

过去 1 个月内

过去 1 年内

所有结果

精确匹配

[Google 学术: Polyethylene glycol microspheres loaded with irinotecan for arterially directed embolic therapy of metastatic liver cancer](#)

[... and leucovorin for metastatic colorectal cancer](#) - Saltz - 被引用次数: 3216

[... \(TACE\) for colorectal liver metastases—current status](#) ... - Massmann - 被引用次数: 8

[... from hydrophilic resorbable embolization microspheres](#) ... - Bédouet - 被引用次数: 7

[Liver-directed therapies in metastatic colorectal cancer - NCBI - NIH](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4173041/>

Hepatic resection is the treatment of choice for colorectal liver metastases ... portal vein embolization (PVE), two stage hepatectomy, ablation and hepatic ... with chemotherapy solution, or as drug-eluting beads loaded with irinotecan ... RE can be performed with microspheres labeled with the β emitter yttrium-90 (^{90}Y).

[Non-operative therapies for colorectal liver metastases - NCBI - NIH](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4311101/>

The armamentarium for liver-directed therapy includes ablative therapies, ... Keywords: Colon cancer, liver metastases, embolization, radiation, ablation Drug eluting beads loaded with irinotecan (DEBIRI) provide an alternative for ... about 80% of liver tumors, the Yttrium-90 microspheres preferentially flow to the tumors.

[Chemoembolization Adopting Polyethylene Glycol Drug-Eluting ...](#)

www.ajronline.org/doi/full/10.2214/AJR.16.17477?src=recsys...0

A new generation of loadable embolics made of polyethylene glycol (PEG) has been ... These microspheres are indicated for embolization of blood vessels Therefore, for patients with unresectable primary liver cancer, liver-directed therapies are Intra-arterial infusion of irinotecan-loaded drug-eluting beads (DEBIRI) ...

[Liver-Directed Therapy for Hepatic Malignancies | Gotoper.com](#)