



6

**Name of Journal:** *World Journal of Gastrointestinal Endoscopy***ESPS Manuscript NO:** 26850**Manuscript Type:** ORIGINAL ARTICLE**Retrospective Study****2**  
**Bleeding risk with clopidogrel and percutaneous endoscopic gastrostomy****Umair Sohail, Chela Harleen, Amin O Mahdi, Murtaza Arif, Douglas L Nguyen, Matthew L Bechtold****Abstract****2**  
**AIM:** To compare bleeding within 48 hours in patients undergoing PEG with or without clopidogrel.**METHODS:** After IRB approval, a retrospective study involving a single center was conducted on adult patients having PEG (1/08-1/14). Patients were divided into two groups: Clopidogrel group consisting of those patients taking clopidogrel within 5 days of PEG and the non-clopidogrel group including those patients not taking clopidogrel within 5 days of the PEG.**RESULTS:** 319 PEG patients were found. 168 males and 151 females with mean BMI  $28.47 \pm 9.75 \text{ kg/m}^2$  and mean age  $65.03 \pm 16.11$  years were identified. 33 patients were**Match Overview**

- |          |   |    |
|----------|---|----|
| <b>1</b> | <b>CrossCheck</b> 109 words<br>Sohail, Umair, Sameer Siddique, Harleen K Chela, Samuel J ersak, Zainab Arif, Matthew L Bechtold, and Murtaza Arif. "56  | 6% |
| <b>2</b> | <b>CrossCheck</b> 49 words<br>"Endoscopy". The American Journal of Gastroenterology, 20 15.   | 3% |
| <b>3</b> | <b>CrossCheck</b> 22 words<br>"Scientific Program for 30th World Congress of Endourology & SWL", Journal of Endourology, 2012.                          | 1% |
| <b>4</b> | <b>CrossCheck</b> 19 words<br>"Abstracts and Index of Authors", Pharmacotherapy The Jour nal of Human Pharmacology and Drug Therapy, 2013.              | 1% |
| <b>5</b> | <b>CrossCheck</b> 16 words<br>Vainrub, S., A. E. Patanwala, R. Cosgrove, R. Poston, and P. E. Nolan. "Bleeding Outcomes in Patients Given Clopidogre... | 1% |
| <b>6</b> | <b>Internet</b> 11 words<br>crawled on 22-Jun-2015<br><a href="http://www.wjgnet.com">www.wjgnet.com</a>  | 1% |



全部 图片 视频 新闻 更多 ▾ 搜索工具

已启用

找到约 22,400 条结果 (用时 0.64 秒)

### 560 Risk of Early Post-Procedure Bleeding With ...

[www.giejournal.org/article/S0016-5107\(15\)02168-9/abstract](http://www.giejournal.org/article/S0016-5107(15)02168-9/abstract) ▾ 翻译此页

作者: U Sohail - 2015

560 Risk of Early Post-Procedure Bleeding With Percutaneous Endoscopic Gastrostomy (PEG) Tube Placement in Patients Using Clopidogrel. Umair Sohail.

### 560 Risk of Early Post-Procedure Bleeding With ...

[https://www.researchgate.net/.../276138103\\_560\\_Risk\\_of\\_Ear...](https://www.researchgate.net/.../276138103_560_Risk_of_Ear...) ▾ 翻译此页

560 Risk of Early Post-Procedure Bleeding With Percutaneous Endoscopic Gastrostomy (PEG) Tube Placement in Patients Using Clopidogrel on ResearchGate, ...

### Risk of Bleeding After Percutaneous Endoscopic ...

[https://www.researchgate.net/.../51851133\\_Risk\\_of\\_Bleeding\\_...](https://www.researchgate.net/.../51851133_Risk_of_Bleeding_...) ▾ 翻译此页

Patients who undergo percutaneous endoscopic gastrostomy (PEG) placement ... and to determine if clopidogrel increases the risk of bleeding following PEG.... as a high-risk endoscopic procedure, bleeding with PEG placement was rare, ... on heparin infusion undergoing PEG, delaying the procedure, alternative use of ...

### Bleeding after percutaneous endoscopic gastrostomy is ...

[www.ncbi.nlm.nih.gov/pubmed/21704806](http://www.ncbi.nlm.nih.gov/pubmed/21704806) ▾ 翻译此页

作者: JA Richter - 2011 - 被引用次数: 33 - 相关文章

Bleeding after percutaneous endoscopic gastrostomy is linked to serotonin reuptake inhibitors, not aspirin or clopidogrel. ... PEG. Serotonin reuptake inhibitors (SRIs) have been associated with an increased risk of GI bleeding. ... aspirin, clopidogrel, or SRI use and bleeding in patients who underwent PEG tube placement.

### Risk factors and complications following percutaneous ...

[www.ncbi.nlm.nih.gov/.../PubMed\\_Central\\_\(PMC\)\\_...](http://www.ncbi.nlm.nih.gov/.../PubMed_Central_(PMC)_...) ▾ 翻译此页



全部 图片 视频 新闻 更多 ▾ 搜索工具

找到约 33,100 条结果 (用时 0.60 秒)

### Google 学术: Bleeding risk with clopidogrel and percutaneous endoscopic gastrostomy

... percutaneous endoscopic gastrostomy: a case series ... - Sabine Richter - 被引用次数: 54

Bleeding after percutaneous endoscopic gastrostomy ... - Richter - 被引用次数: 33

... bleeding after percutaneous endoscopic gastrostomy ( ... - Singh - 被引用次数: 21

### Bleeding after percutaneous endoscopic gastrostomy is linked ... - NCBI

[www.ncbi.nlm.nih.gov/pubmed/21704806](http://www.ncbi.nlm.nih.gov/pubmed/21704806) ▾ 翻译此页

作者: JA Richter - 2011 - 被引用次数: 33 - 相关文章

Bleeding after percutaneous endoscopic gastrostomy is linked to serotonin reuptake inhibitors, not aspirin or clopidogrel. Richter JA(1), Patrie ... Serotonin reuptake inhibitors (SRIs) have been associated with an increased risk of GI bleeding.

### Risk of bleeding after percutaneous endoscopic gastrostomy ... - NCBI

[www.ncbi.nlm.nih.gov/pubmed/22138961](http://www.ncbi.nlm.nih.gov/pubmed/22138961) ▾ 翻译此页

作者: D Singh - 2012 - 被引用次数: 21 - 相关文章

2011年12月3日 - Risk of bleeding after percutaneous endoscopic gastrostomy (PEG). ... to determine if clopidogrel increases the risk of bleeding following PEG.

### Risk of bleeding in patients undergoing percutaneous endoscopic ...

[www.ncbi.nlm.nih.gov/pubmed/25733036](http://www.ncbi.nlm.nih.gov/pubmed/25733036) ▾ 翻译此页

作者: AJ Lucendo - 2015 - 被引用次数: 6 - 相关文章

Risk of bleeding in patients undergoing percutaneous endoscopic ... BACKGROUND AND AIM: Patients undergoing percutaneous endoscopic gastrostomy (PEG) ... aspirin and/or clopidogrel treatment increases the risk of bleeding following ...

全部 图片 视频 新闻 更多 ▾ 搜索工具

找到约 32,100 条结果 (用时 0.60 秒)

### Google 学术: Bleeding risk with clopidogrel and percutaneous endoscopic gastrostomy

Bleeding after percutaneous endoscopic gastrostomy ... - Richter - 被引用次数: 36

... percutaneous endoscopic gastrostomy: a case series ... - Richter-Schrag - 被引用次数: 58

... bleeding after percutaneous endoscopic gastrostomy ( ... - Singh - 被引用次数: 24

### Bleeding after percutaneous endoscopic gastrostomy is linked ... - NCBI

[www.ncbi.nlm.nih.gov/pubmed/21704806](http://www.ncbi.nlm.nih.gov/pubmed/21704806) ▾ 翻译此页

作者: JA Richter - 2011 - 被引用次数: 33 - 相关文章

Bleeding after percutaneous endoscopic gastrostomy is linked to serotonin reuptake inhibitors, not aspirin or clopidogrel. Richter JA(1), Patrie ... Serotonin reuptake inhibitors (SRIs) have been associated with an increased risk of GI bleeding.

### Risk of bleeding after percutaneous endoscopic gastrostomy ... - NCBI

[www.ncbi.nlm.nih.gov/pubmed/22138961](http://www.ncbi.nlm.nih.gov/pubmed/22138961) ▾ 翻译此页

作者: D Singh - 2012 - 被引用次数: 22 - 相关文章

2011年12月3日 - Risk of bleeding after percutaneous endoscopic gastrostomy (PEG). ... to determine if clopidogrel increases the risk of bleeding following PEG.

### Risk of bleeding in patients undergoing percutaneous endoscopic ...

[www.ncbi.nlm.nih.gov/pubmed/25733036](http://www.ncbi.nlm.nih.gov/pubmed/25733036) ▾ 翻译此页

作者: AJ Lucendo - 2015 - 被引用次数: 6 - 相关文章

Risk of bleeding in patients undergoing percutaneous endoscopic ... BACKGROUND AND AIM: Patients undergoing percutaneous endoscopic gastrostomy (PEG) ... aspirin and/or clopidogrel treatment increases the risk of bleeding following ...