23351-Review

- Inte



Name of Journal: World Journal of Gastrointestinal Endoscopy

ESPS Manuscript NO: 23351

Manuscript Type: Case Report

Diagnosis of a submucosal mass at the staple line after sigmoid colon cancer resection by endoscopic cutting-mucosa biopsy

Mitsuaki Morimoto, Koji Koinuma, Alan K Lefor, Hisanaga Horie, Homare Ito, Naohiro Sata, Yoshikazu Hayashi, Keijiro Sunada, Hironori Yamamoto

A 10 years ald man undamitant languagemic piamaid calan magazina for carrow

Abstract

CrossCheck 65 words 5% "COLORECTAL CANCER PREVENTION", The American J ournal of Gastroenterology, 09/2008 CrossCheck 24 words 2% Berho, Mariana, Steven D. Wexner, Ana-Maria Botero-Anu g Dori Pelled, and James W. Fleshman. "Histopathologic A CrossCheck 15 words 1% Han, Yoo Min, Jong Yeul Lee, Il Ju Choi, Chan Gyoo Kim, Soo-Jeong Cho, Jun Ho Lee, Hyun Beom Kim, and Ji Min C CrossCheck 15 words 1% Morris E. Franklin. "The Use of Bioabsorbable Staple Line Reinforcement for Circular Stapler (BSG 222Seamquard ... CrossCheck 15 words 1% F. Manno. "An uncommon finding; a retrorectal Schwanno ma". Colorectal Disease, 5/17/2007 Internet 11 words 1% crawled on 22-Jun-2015 www.wignet.com

Match Overview



Diagnosis of a submucosal mass at the staple line after rectal cancer resect






网页 图片 视频 新闻 购物 更多▼ 搜索工具

找到约 11,200 条结果 (用时 0.78 秒)

Surgical Treatment of Gastric Gastrointestinal Stromal Tumor www.ncbi,nlm.nih.gov > ... → PubMed Central (PMC) ▼ 翻译此页作者: SH Kong - 2013 - 被引用次数: 21 - 相关文章 2013年3月31日 - If the submucosal tumor is suspected to be a GIST and is larger than 2~3 cm on ... Endoscopic biopsy usually yields only normal mucosa, but these biopsies ... a tyrosine kinase inhibitor as the first line treatment such as those patients with After cutting the gastric wall using either energy-based devices or ... 缺少字词: staple

Endoscopic mucosal resection and endoscopic submucosal ...
www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▼ 翻译此页
作者: KK Wang - 2010 - 被引用次数: 45 - 相关文章
Endoscopic mucosal resection (EMR) and endoscopic submucosal dissection ... As opposed to endoscopic biopsies that remove on average 1 mm × 3 mm area of ... ESD has been used after chemoradiation to complete removal of tumor in very ... In the treatment of neoplasia, lesions that have high risk of metastasis are not ...
缺少字词: staple

Endoscopic Mucosal Resection: Overview, Periprocedural ... emedicine.medscape.com/article/1891659-overview ▼ 翻译此页 2015年8月6日 - Introduction Endoscopic mucosal resection (EMR) is a technique used ...







全部 图片 视频 新闻 购物 更多 搜索工具

找到约 10,100 条结果 (用时 0.61 秒)

Colectomy - 日本オンラインショッピング

www.yoursurgery.com/ProcedureDetails.cfm?Proc=17 ▼ 翻译此页

Surgery of colon and rectum is done for various reasons including cancer, diverticulitis, ... the ascending colon, transverse colon, descending colon and sigmoid colon. ...

Submucosa - The area between the mucosa and circular muscle layer and is ... This

technique may detect a lesion but a biopsy cannot be obtained.

缺少字词: endoscopie

Malignant colorectal polyps

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▼ 翻译此页

作者: L Bujanda - 2010 - 被引用次数: 71 - 相关文章

2010年7月7日 - More than 25% of advanced polyps[2] and colon cancers are located in

... When the carcinoma spreads to the submucosa, the polyp is considered the distance between the invasive tumor and the cauterized biopsy margin < 2 ... ESD is

better than conventional endoscopic mucosal resection because it ...

缺少字词: staple

[PDF] Removing large or sessile colonic polyps

www.worldendo.org/.../omed_hid_removing_large_or_sessile_... ▼ 翻译此页 Endoscopic mucosal resection (EMR) has become the standard technique ... for instance in the sigmoid color in the presence of diverticular disease, then submucosal phigns in the presence of diverticular disease, then submucosal phigns in the presence of diverticular disease.



Diagnosis of a submucosal mass at the staple line after sigmoid colon cance



全部

图片 视频

新闻

更多▼

搜索工具

找到约 9,430 条结果 (用时 0.71 秒)

Colectomy - http://www.yoursurgery.com

www.yoursurgery.com/ProcedureDetails.cfm?Proc=17 ▼ 翻译此页

Surgery of colon and rectum is done for various reasons including cancer, diverticulitis, ... the ascending colon, transverse colon, descending colon and sigmoid colon. ...

Submucosa - The area between the mucosa and circular muscle layer and is ... This technique may detect a lesion but a biopsy cannot be obtained. 缺少字词: endoscopic

Malignant colorectal polyps

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▼ 翻译此页

作者: L Bujanda - 2010 - 被引用次数: 70 - 相关文章

2010年7月7日 - More than 25% of advanced polyps[2] and colon cancers are located in

.... When the carcinoma spreads to the submucosa, the polyp is considered the distance between the invasive tumor and the cauterized biopsy margin < 2 ... ESD is better than conventional endoscopic mucosal resection because it ...

缺少字词: staple

[PDF] Removing large or sessile colonic polyps

www.worldendo.org/.../omed_hid_removing_large_or_sessile_... ▼ 翻译此页 Endoscopic mucosal resection (EMR) has become the standard technique ... for instance in the sigmoid colon in the presence of diverticular disease, then submucosal ... biopsies should be taken and tattoos placed around the lesion for surgical ... An early repeat colonoscopy is necessary 3 months after piecemeal EMR to ... 缺少字词: staple

[PDF] How to approach and manage difficult polyps www.gastrocol.com/file/Revista/en_v27n4a06.pdf ▼ 翻译此页