

Match Overview



1	Internet 28 words crawled on 04-Mar-2016 www.wjgnet.com	1%
2	Internet 10 words crawled on 28-Apr-2015 www.hepatogastroenterology.org	<1%
3	CrossCheck 8 words H. Choi. "Endoscopic stone removal using a scissors papil lotome followed by papillary balloon dilation in patients w ...	<1%
4	CrossCheck 7 words Sakai, Yuji. "New cannulation method for pancreatic duct ... annulation-bile duct guidewire-indwelling method", World J	<1%

1
Name of Journal: *World Journal of Gastrointestinal Endoscopy*

ESPS Manuscript NO: 27770

Manuscript Type: Original Article

Retrospective Study

Utility of endoscopic retrograde cholangiopancreatography on biliopancreatic diseases in the patients with Billroth II-reconstructed stomach

2
Yuji Sakai, Toshio Tsuyuguchi, Rintaro Mikata, Harutoshi Sugiyama, Shin Yasui, Masaru Miyazaki, Osamu Yokosuka

Abstract

AIM: To examine the utility of endoscopic retrograde cholangiopancreatography (ERCP) on biliopancreatic diseases in the patients with Billroth II-reconstructed stomach.

METHODS: On 26 cases of biliopancreatic diseases in the patients with Billroth II-reconstructed stomach, the ERCP was conducted using a straight view scope or a retrograde oblique-viewing endoscope. All the cases were the patients aiming at selective insertion into the bile duct. One patient aimed at diagnosis, and 25 patients aimed at treatment. The cases in which the endoscope reaches the duodenal papilla and anastomosis and insertion into the bile duct became possible were considered successful.

RESULTS: The rate of reaching the duodenal papilla and anastomosis was 84.7%.

[全部](#)[图片](#)[视频](#)[新闻](#)[更多 ▾](#)[搜索工具](#)

找到约 802 条结果 (用时 0.68 秒)

(ERCP) after Billroth II reconstruction - UpToDate

www.uptodate.com/.../endoscopic-retrograde-cholangiopancreatography-erc... ▾ [翻译此页](#)

2016年1月7日 - Partial gastrectomy with **reconstruction** can be performed to treat complications of peptic ulcer **disease** (such as perforation, bleeding, ... remains a commonly used **reconstruction** after distal **stomach** resection. ... The endoscopist performing **ERCP** in a **Billroth II patient** has to consider and ... Terms of **Use**.

Endoscopic retrograde cholangiopancreatography in patients ... - NCBI

www.ncbi.nlm.nih.gov ▸ [NCBI](#) ▸ [Literature](#) ▸ [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: TG Moreels - 2014 - 被引用次数: 5 - [相关文章](#)

2014年8月16日 - **ERCP** in **patients** with altered anatomy faces three important challenges ... **ERCP** in **patients** with altered anatomy and difficult-to-access **biliopancreatic** ... The **use** of a semi-rigid overtube in a case of DAE inhibits loop formation of ... Similarly to the **Billroth II reconstruction**, the angulation towards the afferent ...

Endoscopic retrograde cholangiopancreatography in patients ... - NCBI

www.ncbi.nlm.nih.gov ▸ [NCBI](#) ▸ [Literature](#) ▸ [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: JB Samarasena - 2012 - 被引用次数: 2 - [相关文章](#)

2012年4月1日 - Historically, **Billroth II** gastrectomy was the predominant surgically ... Roux-en-Y **reconstruction** is also commonly performed after ... and then advanced up the **biliopancreatic** (afferent) limb until the biliary or pancreatic orifice is reached..... The **use** of a pediatric colonoscope to perform **ERCP** in **patients** with ...

[PDF] ERCP in patients with prior Billroth II gastrectomy: report of 30 ... - A...

www.webaigo.it/download/Endoscopy%20July%202015.pdf ▾ [翻译此页](#)

作者: V Bove - 2015 - 被引用次数: 7 - [相关文章](#)

2015年3月2日 - ment several biliary and pancreatic **diseases**. **ERCP** may also ... with **Billroth II reconstruction** who underwent **ERCP** between ... of the desired **biliopancreatic** duct was 93.8%. (580/618). ... **ERCP** in **patients** with previous **Billroth II** gastrectomy was always ... **Use** fluoroscopy to assess correct loop intubation.

找到约 625 条结果 (用时 0.79 秒)

(ERCP) after Billroth II reconstruction - UpToDate

www.uptodate.com/.../endoscopic-retrograde-cholangiopancreatography-erc... ▾ 翻译此页

2016年1月7日 - Partial gastrectomy with reconstruction can be performed to treat complications of peptic ulcer disease (such as perforation, bleeding, ... remains a commonly used reconstruction after distal stomach resection. ... The endoscopist performing ERCP in a Billroth II patient has to consider and ... Terms of Use.

[PDF] ERCP in patients with prior Billroth II gastrectomy: report of 30 ... - Al...

www.webaigo.it/download/Endoscopy%20July%202015.pdf ▾ 翻译此页

作者: V Bove - 2015 - 被引用次数: 8 - 相关文章

2015年3月2日 - ment several biliary and pancreatic diseases. ERCP may also ... with Billroth II reconstruction who underwent ERCP between ... of the desired biliopancreatic duct was 93.8% (580/618). ... ERCP in patients with previous Billroth II gastrectomy was always ... Use fluoroscopy to assess correct loop intubation.

Endoscopic retrograde cholangiopancreatography in patients ... - NCBI

[https://www.ncbi.nlm.nih.gov/NCBI/Literature/PubMedCentral\(PMC\)](https://www.ncbi.nlm.nih.gov/NCBI/Literature/PubMedCentral(PMC)) ▾ 翻译此页

作者: TG Moreels - 2014 - 被引用次数: 7 - 相关文章

2014年8月16日 - ERCP in patients with altered anatomy faces three important challenges ... ERCP in patients with altered anatomy and difficult-to-access biliopancreatic ... The use of a semi-rigid overtube in a case of DAE inhibits loop formation of ... Similarly to the Billroth II reconstruction, the angulation towards the afferent ...

Endoscopic retrograde cholangiopancreatography in patients ... - NCBI

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3655346/> ▾ 翻译此页

作者: JB Samarasena - 2012 - 被引用次数: 2 - 相关文章

2012年4月1日 - Historically, Billroth II gastrectomy was the predominant surgically ... Roux-en-Y reconstruction is also commonly performed after ... and then advanced up the biliopancreatic (afferent) limb until the biliary or pancreatic orifice is reached. ... The use of a pediatric colonoscope to perform ERCP in patients with ...

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 696 条结果 (用时 0.55 秒)

(ERCP) after Billroth II reconstruction - UpToDate

www.uptodate.com/.../endoscopic-retrograde-cholangiopancreatography-erc... ▼ 翻译此页

2016年12月15日 - Partial gastrectomy with **reconstruction** can be performed to treat complications of peptic ulcer **disease** (such as perforation, bleeding, ... remains a commonly used **reconstruction** after distal **stomach** resection. ... The endoscopist performing **ERCP** in a **Billroth II patient** has to consider and Terms of **Use**.

[PDF] ERCP in patients with prior Billroth II gastrectomy: report of 30 ... - Al...

www.webaigo.it/download/Endoscopy%20July%202015.pdf ▼ 翻译此页

作者: V Bove - 2015 - 被引用次数: 9 - 相关文章

2015年3月2日 - ment several biliary and pancreatic **diseases**. **ERCP** may also ... with **Billroth II reconstruction** who underwent **ERCP** between ... of the desired **biliopancreatic** duct was 93.8%. (580/618). ... **ERCP** in **patients** with previous **Billroth II** gastrectomy was always ... **Use** fluoroscopy to assess correct loop intubation.

Endoscopic retrograde cholangiopancreatography in patients ... - NCBI

[https://www.ncbi.nlm.nih.gov/NCBI/Literature/PubMedCentral \(PMC\)](https://www.ncbi.nlm.nih.gov/NCBI/Literature/PubMedCentral/PMC/) - 翻译此页

作者: TG Moreels - 2014 - 被引用次数: 11 - 相关文章

2014年8月16日 - **ERCP** in **patients** with altered anatomy faces three important challenges ... **ERCP** in **patients** with altered anatomy and difficult-to-access **biliopancreatic** ... The **use** of a semi-rigid overtube in a case of DAE inhibits loop formation of ... Similarly to the **Billroth II reconstruction**, the angulation towards the afferent ...

Endoscopic Retrograde Cholangiopancreatography in Post ... - NCBI

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5152788/> - 翻译此页

作者: CH Park - 2016