

Name of Journal: *World Journal of Hepatology*

Manuscript NO: 36628

Manuscript Type: CASE REPORT

Contrast uptake in primary hepatic angiosarcoma on Gd-EOB-DTPA-enhanced magnetic resonance imaging in the hepatobiliary phase

Hayashi *et al.* Gd-EOB-DTPA-enhanced MRI for angiosarcoma

Manabu Hayashi, Satoshi Kawana, Hirofumi Sekino, Kazumichi Abe, Naoki Matsuoka, Masahito Kashiwagi, Ken Okai, Yukiko Kanno, Atsushi Takahashi, Hiroshi Ito, Yuko Hashimoto, Hiromasa Ohira

Match Overview

1	Internet 67 words crawled on 01-Jan-2013 radiology.rsna.org	3%
2	Crossref 36 words Satoru Joshiita: "A case of well-differentiated hepatic angiosarcoma with prominent vascularization in the hepatobiliary phase of Gd-EOB-DTPA-enhanced MR imaging"	2%
3	Crossref 32 words Ryota Sagami, Hidefumi Nishikiori, Shôji Yamada, Toshiaki Yamamoto, Tôru	1%
4	Crossref 26 words Yasuyuki Shimoyama: "Hereditary hemochromatosis associated with hepatic angiomyomatosis"	1%
5	Crossref 18 words JANET R. HARRISON: "Hepatic Angiosarcoma - An Unusual Cause of Cancer"	1%
6	Internet 15 words crawled on 31-Aug-2017 diva-portal.org	1%
7	Internet 14 words crawled on 02-Sep-2016 mdanderson.influent.utsystem.edu	1%
8	Crossref 14 words Honda, M.: "Development of hepatic angiomyomatosis associated with familial hemochromatosis"	1%
9	Crossref 13 words Akinobu Osame: "Multinodular pseudoluminal angiomyomatosis of the liver associated with hemochromatosis"	1%
10	Crossref 13 words Sachiyo Kogita: "Gd-EOB-DTPA-enhanced MR imaging of hepatic angiomyomatosis"	1%
11	Crossref 13 words Wang, Liang, Ke Lv, Xiao-Yan Chang, Wei Wei, Zhi-Ming Yang, Wei Wei, Liang Yu	1%
12	Crossref 12 words Functional Imaging in Oncology, 2014.	1%
13	Crossref 12 words Yamashita, Taro, Azusa Kitao, Osamu	1%



Contrast uptake in primary hepatic angiosarcoma on Gd-EOB-DTPA-enhanc



学术搜索

找到约 37 条结果 (用时 0.22 秒)

我的著作引用情况

文章

我的图书馆

时间不限

2017以来

2016以来

2013以来

自定义范围...

按相关性排序

按日期排序

不限语言

中文网页

简体中文网页

 包括专利 包含引用 创建快讯

小提示：只搜索中文(简体)结果，可在 学术搜索设置 指定搜索语言

[\[HTML\] Focal hepatic lesions in Gd-EOB-DTPA enhanced MRI: the atlas](#)

JT Campos, CB Sirlin, JY Choi - Insights into imaging, 2012 - Springer

.. a Pre-contrast, b, c arterial phase, d PV phase, e 3 min, f ... confluent hepatic fibrosis demonstrates hypointense areas due to decreased or absent hepatocellular uptake of Gd ... Cavernous haemangiomas represent the most common primary liver tumour and consist of multiple large ...

被引用次数: 46 相关文章 所有 12 个版本 引用 保存

[\[HTML\] springer.com](#)[MR imaging of benign hepatic tumors](#)

T Motohara, RC Semelka... - Magnetic Resonance ... 2002 - mri.theclinics.com

.. with slight heterogeneity on A, moderately low signal intensity on B, enhances in an intense diffuse heterogeneous fashion on C, and fades to slightly higher than background liver on D ... (From

Semelka RC: Hepatic angiomyolipoma with minimal fat content: MR demonstration ...

被引用次数: 42 相关文章 所有 4 个版本 引用 保存

[The natural history of a hepatic angiosarcoma that was difficult to differentiate from cavernous hemangioma](#)[\[PDF\] jst.go.jp](#)

A Okano, H Sonoyama, Y Masano, T Taniguchi... - Internal ... 2012 - jstage.jst.go.jp

.. Many cases have been reported in which angiosarcomas of various organs show high FDG uptake on PET ... Figure 5. Plain (A) and arterial phase contrast-enhanced (B) CT images showed the presence of a 15-mm ... Primary hepatic angiosarcoma: findings at CT and MR imaging ...

被引用次数: 19 相关文章 所有 6 个版本 引用 保存

[Hepatocellular Carcinoma and Other Hepatic Malignancies](#)

CG Roth, DG Mitchell - Radiologic Clinics, 2014 - radiologic.theclinics.com

.. Hepatocellular Carcinoma and Other Hepatic Malignancies, MR Imaging, Christopher ... pattern. (D) The 20 minute-delayed hepatobiliary T1-weighted image shows lack of contrast uptake and nonhepatocellular composition. ...

被引用次数: 19 相关文章 所有 4 个版本 引用 保存

[MRI evaluation of the liver](#)

KL Beavers, RC Semelka - Seminars in liver disease, 2001 - thieme-connect.com

.. with liver during hepatic venous phase, and delayed observation of contrast uptake by the ... is the second most common primary liver malignancy, representing 10% of primary liver tumors ... arterial dominant-phase image demonstrates normal liver anatomy, with contrast present in ...

被引用次数: 21 相关文章 所有 5 个版本 引用 保存

[CT and MR imaging findings of hepatic epithelioid hemangioendotheliomas:](#)



文章

找到约 37 条结果 (用时 0.07 秒)

时间不限

2017以来

2016以来

2013以来

自定义范围...

按相关性排序

按日期排序

不限语言

中文网页

简体中文网页

 包括专利 包含引用 创建快讯

小提示：只搜索中文(简体)结果，可在 学术搜索设置 指定搜索语言

[HTML] Focal hepatic lesions in Gd-EOB-DTPA enhanced MRI: the atlas

[HTML] [springer.com](#)JT Campos, CB Sirlin, JY Choi - *Insights into imaging*, 2012 - Springer

... a Pre-contrast, b, c arterial phase, d PV phase, e 3 min, f ... confluent hepatic fibrosis demonstrates hypointense areas due to decreased or absent hepatocellular uptake of Gd ... Cavernous haemangiomas represent the most common primary liver tumour and consist of multiple large ...

☆ 99 被引用次数: 46 相关文章 所有 12 个版本

The natural history of a hepatic angiosarcoma that was difficult to differentiate from cavernous hemangioma

[PDF] [jst.go.jp](#)A Okano, H Sonoyama, Y Masano, T Taniguchi... - *Internal ...*, 2012 - jstage.jst.go.jp

... have been reported in which angiosarcomas of various organs show high FDG uptake on PET ... Figure 5. Plain (A) and arterial phase contrast-enhanced (B) CT images showed the presence of a 15 ... Molina E, Hernandez A. Clinical manifestations of primary hepatic angiosarcoma ...

☆ 99 被引用次数: 22 相关文章 所有 6 个版本

Hepatocellular Carcinoma and Other Hepatic Malignancies

CG Roth, DG Mitchell - *Radiologic Clinics*, 2014 - radiologic.theclinics.com

... Thomas Jefferson University, 1094 Main Building, 132 South 10th Street, Philadelphia, PA 19107, USA., MD. Hepatocellular Carcinoma and Other Hepatic Malignancies ... (A) Axial in-phase T1-weighted image shows a faintly hypointense lesion near the liver dome (arrow) ...

☆ 99 被引用次数: 20 相关文章 所有 4 个版本

MR imaging of benign hepatic tumors

T Motohara, RC Semelka... - *Magnetic Resonance ...*, 2002 - mri.theclinics.com

... on A, moderately low signal intensity on B, enhances in an intense diffuse heterogeneous fashion on C, and fades to slightly higher than background liver on D. This ... (From Semelka RC. Hepatic angiomyolipoma with ... Magn Reson Imaging 14:687-689, 1996, with permission.) ...

[全部](#)[图片](#)[新闻](#)[地图](#)[更多](#)[设置](#)[工具](#)

找到约 2,420 条结果 (用时 0.64 秒)

Focal hepatic lesions in Gd-EOB-DTPA enhanced MRI: the atlas

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3443279/> ▾ 翻译此页

作者: JT Campos - 2012 - 被引用次数: 47 - 相关文章

2012年6月15日 - Focal hepatic lesions in Gd-EOB-DTPA enhanced MRI: the atlas ... Gd-EOB-DTPA is a contrast agent with combined properties of a conventional ... Gd-EOB-DTPA has greater hepatocellular uptake and biliary excretion than ... Moreover, uptake by functioning hepatocytes in the delayed phase provides ...

Blumgart's Surgery of the Liver, Pancreas and Biliary Tract E-Book

<https://books.google.ae/books?isbn=0323340873> ▾ 翻译此页

William R. Jarnagin - 2016 - Medical

Magnetic resonance cholangiopancreatography and contrastenhanced magnetic ... Imaging features of intrahepatic cholangiocarcinoma in GdEOBDTPAenhanced MRI ... Primary hepatic angiosarcoma: multiinstitutional comprehensive cancer centre ... V. Resonance absorption by nuclear magnetic moments in a solid.

[PDF] A Pictorial Review of Gd-EOB-enhanced MR Imaging of ... - netkey.at

pdf.posterng.netkey.at/download/index.php?module=get_pdf_by... ▾ 翻译此页

Various Benign and Malignant Focal Liver Lesions with. Multimodality ... Keywords: Liver, MR, Contrast agent-intravenous, Education and training ... from uptake of Gd-EOB-DTPA in hepatobiliary phase. 2 ... phase on Gd-EOB-DTPA-enhanced MRI Most common primary malignant hepatic tumor. 2 ... Angiosarcoma. 1.

Gd-EOB-DTPA-Enhanced MRI for Detection of Liver Metastases from ...

<https://www.hindawi.com/journals/ijh/2013/572307/> ▾ 翻译此页

作者: KJ Lafaro - 2013 - 被引用次数: 18 - 相关文章