



15,400 Results Any time ▾

Including results for characterization of patients with both alcoholic and nonalcoholic fatty liver disease **nafl** in a large united states cohort.

Do you want results only for Characterization of Patients with Both Alcoholic and Nonalcoholic Fatty Liver Disease (BAFLD) in a Large United States Cohort ?

Dyslipidemia in Patients with Nonalcoholic Fatty Liver Disease

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3654545>

Nonalcoholic fatty liver disease (NAFLD) is one of the most common causes of chronic liver disease in the Western world. 1,2 Its incidence is increasing in both adults and children, and it is expected to become the most common cause for liver transplantation in the United States by 2020. 3 It encompasses a spectrum of liver disease, ranging from nonalcoholic fatty liver (NAFL) to ...

Cited by: 127 Author: Hemant Chatrath, Raj Vuppalanchi, Naga ...

Publish Year: 2012

Clinical differences between alcoholic liver disease and ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4093692>

Jul 14, 2014 · Core tip: Although alcoholic liver disease (ALD) and nonalcoholic fatty liver disease (NAFLD) have similar pathological spectra, from simple hepatic steatosis to steatohepatitis and liver cirrhosis, the epidemiological and clinical characteristics of these two diseases differ. Comparative analyses of the factors associated with disease susceptibility and progression and the predictors and ...

Cited by: 21 Author: Nobuyuki Toshikuni, Mikihiro Tsutsumi, To...

Publish Year: 2014

[PDF] Alcoholic and Nonalcoholic Fatty Liver Disease and ...

[https://www.cghjournal.org/article/S1542-3565\(19\)30514-2/pdf](https://www.cghjournal.org/article/S1542-3565(19)30514-2/pdf)

10 Background: There are limited data to compare the associations of nonalcoholic fatty liver 11 disease (NAFLD) and alcohol-associated fatty liver disease (AFLD) with risk of incident 12 liver and cardiovascular diseases 13 Findings: In a large cohort study, we found an increased risk of liver disease ...

Prevalence of non-alcoholic fatty liver disease and risk ...

<https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0173499>


Mar 27, 2017 · In the United States, non-alcoholic fatty liver disease (NAFLD) is the most common liver disease and associated with higher mortality according to data from earlier National Health and Nutrition Examination Survey (NHANES) 1988–1994. Our goal was to determine the NAFLD prevalence in the recent 1999–2012 NHANES, risk factors for advanced fibrosis (stage 3–4) and mortality.

Cited by: 57 Author: Michael H. Le, Pardha Devaki, Nghiem B....

Publish Year: 2017

Non-Alcoholic fatty liver disease

Medical Condition



A condition in which fat accumulates in the liver in people who drink little or no alcohol.

- Very common (More than 3 million cases per year in US)
- Diagnosis requires lab test or imaging
- Treatments can help manage condition, no known cure
- Can last several years or be lifelong

The cause of non-alcoholic fatty liver disease is not clearly understood. It is associated with obesity, insulin resistance, hyperglycemia, and high levels of fat in blood. The condition does not usually cause symptoms. In some cases, fatigue, enlarged liver, and pain in the upper right abdomen is seen. Non-alcoholic fatty liver disease has no specific treatment. It involves treating underlying conditions such as obesity.

Symptoms

Usually asymptomatic. Symptoms may include:

Name of Journal: *World Journal of Hepatology*

Manuscript NO: 48729

Manuscript Type: ORIGINAL ARTICLE

1
Characterization of patients with both alcoholic and nonalcoholic fatty liver disease in a large United States cohort

George Khoudari, Amandeep Singh, Mazen Nouredin, Danielle Fritze, Rocio Lopez, MS, Imad Asaad, Eric Lawitz E, Fred Poordad, Kris V Kowdley, Naim Alkhour

Abstract

BACKGROUND

Nonalcoholic fatty liver disease (NAFLD) is the hepatic manifestation of the metabolic syndrome (MetS) and is characterized by steatosis in the absence of significant alcohol consumption. However, MetS and significant alcohol intake coexist in certain individuals

Match Overview

1	Crossref 108 words N. Alkhour, A. Singh, A. Vigni, M. Nouredin, R. Lopez, K. Watt, E. Lawitz, F. Poordad, K.V. Kowdley. "Characterization of	3%
2	Crossref 47 words Marie Boyle, Steven Masson, Quentin M. Anstee. "The bidirectional impacts of alcohol consumption and the metabo ...	2%
3	Internet 22 words crawled on 01-Feb-2015 onlinelibrary.wiley.com	1%
4	Internet 19 words crawled on 17-Apr-2019 www.aldrodriguezriverfoundation.com	1%



国内版

国际版

Characterization of patients with both alcoholic and nonalcoholic fatty liver diseases



All

Images

Videos

关闭取词

33,000 Results

Any time

Choline intake in a large cohort of patients with ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3302364>

Feb 15, 2012 · Choline intake in a large cohort of patients with nonalcoholic fatty liver disease 1, 2, 3 Anthony L Guerrero , Ryan M Colvin , Amy K Schwartz , Jean P Molleston , Karen F Murray , AnnaMae Diehl , Parvathi Mohan , Jeffrey B Schwimmer , Joel E Lavine , Michael S Torbenson , ...

Cited by: 85

Author: Anthony L Guerrero, Ryan M Colvin, Am...

Publish Year: 2012

Characterization of patients with Both Alcoholic and ...

https://www.researchgate.net/publication/324699957_Characterization_of_patients_with...

Request PDF on ResearchGate | On Apr 1, 2018, N. Alkhoury and others published Characterization of patients with Both Alcoholic and Nonalcoholic Fatty Liver Disease (BAFLD) in a large United ...

Treatment options for nonalcoholic fatty liver disease

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3002571>

Background. Nonalcoholic fatty liver disease (NAFLD) is one of the most common forms of liver disease across the world. NAFLD represents a spectrum ranging from bland steatosis to nonalcoholic steatohepatitis (NASH), and is closely associated with obesity and metabolic syndrome [Ludwig et al. 1980]. Owing to the increasing prevalence of NAFLD and the potential for NASH to progress to ...

Cited by: 108

Author: Brian Lam, Zobair M. Younossi

Publish Year: 2010

Prevalence of non-alcoholic fatty liver disease and risk ...

journals.plos.org/plosone/article?id=10.1371/journal.pone.0173499

In the United States, non-alcoholic fatty liver disease (NAFLD) is the most common liver disease and associated with higher mortality according to data from earlier National Health and Nutrition Examination Survey (NHANES) 1988–1994. Our goal was to determine the NAFLD prevalence in the recent 1999–2012 NHANES, risk factors for advanced fibrosis (stage 3–4) and mortality.

Cited by: 70

Author: Michael H. Le, Pardha Devaki, Nghiem B....

Publish Year: 2017


[PDF] Components of Metabolic Syndrome and the Rising ...

[https://www.gastrojournal.org/article/S0016-5085\(11\)64092-4/pdf](https://www.gastrojournal.org/article/S0016-5085(11)64092-4/pdf)

Components of Metabolic Syndrome and the Rising Prevalence of Non-Alcoholic Fatty Liver Disease (NAFLD) and Non-Alcoholic Steatohepatitis (NASH) in the United States Maria Stepanova, Mariam

Non-Alcoholic Fatty Liver Disease

Medical Condition



A condition in which fat accumulates in the liver in people who drink little or no alcohol.

- Very common (More than 3 million cases per year in US)
- Requires lab test or imaging
- Treatments can help manage condition, no known cure
- Can last several years or be lifelong

The cause of non-alcoholic fatty liver disease is not clearly understood. It is associated with obesity, insulin resistance, hyperglycemia, and high levels of fat in blood. The condition does not usually cause symptoms. In some cases, fatigue, enlarged liver, and pain in the upper right abdomen is seen. Non-alcoholic fatty liver disease has no specific treatment. It involves treating underlying conditions such as obesity.

Symptoms

Usually asymptomatic. Symptoms may include:

- Fatigue
- Enlarged liver
- Pain the upper right abdomen

找到约 489,000 条结果 (用时 0.72 秒)

小提示: 仅限搜索简体中文结果。您可以在[设置](#)中指定搜索语言

Google 学术: [Characterization of patients with both alcoholic and nonalcoholic fatty liver disease in a large United States cohort](#)

[Obesity and non-alcoholic fatty liver disease: Disparate ...](#) - Wong - 被引用次数: 79

[... severity of non-alcoholic fatty liver disease in a large ...](#) - Armstrong - 被引用次数: 227

[The natural history of nonalcoholic fatty liver disease: a ...](#) - Adams - 被引用次数: 2589

[Clinical differences between alcoholic liver disease and ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4093692> [翻译此页](#)

作者: N Toshikuni - 2014 - 被引用次数: 43 - [相关文章](#)

2014年7月14日 - [Alcoholic liver disease \(ALD\)](#) and [nonalcoholic fatty liver disease \(NAFLD\)](#) are ... the epidemiological and clinical [characteristics](#) of these [two](#) diseases differ. ... A nationwide study in the [United States](#) that assessed [patients](#) discharged from A [large cohort](#) study found that the AUROCs of the CAP for the ...

其他用户还问了以下问题

What is Afld? 

How do I know if my alcohol is fatty liver? 

Is fatty liver disease the same as cirrhosis? 

How is alcoholic hepatitis different from cirrhosis? 

[反馈](#)

[Significant cohort of non-alcoholic fatty liver disease with](#)