

Microsoft Bing

国内版 国际版

Association of Non-alcoholic Fatty Liver Disease (NAFLD) and COVID-19

Sign in

ALL IMAGES VIDEOS

148,000 Results Any time

COVID-19 and non-alcoholic fatty liver disease: Two ...

<https://pubmed.ncbi.nlm.nih.gov/32589264>

NAFLD can play a role in the outcome of COVID-19 illness due to frequent association with comorbidities. Initial evidences suggest that increased liver fibrosis in NAFLD might affect COVID-19...

Cited by: 14 Author: Piero Portincasa, Marcin Krawczyk, Marc...

Publish Year: 2020

NAFLD and COVID-19: a Pooled Analysis

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7646222>

Nonalcoholic fatty liver disease is associated with a higher risk of symptomatic, severe, and progressive COVID-19. The association is significant even after adjusting for an important confounding...

Cited by: 4 Author: Sonali Sachdeva, Harshwardhan Khandait...

Publish Year: 2020

Association of Non-alcoholic Fatty Liver Disease with ...

<https://journals.plos.org/plosmedicine/article?id=10.1371/journal.pmed.1001680>

Jul 22, 2014 The high morbidity, mortality, and health care costs associated with CKD have led investigators to seek novel modifiable risk factors. Non-alcoholic fatty liver disease (NAFLD), the...

Search Tools

Turn off Hover Translation (关闭弹窗)

激活 Windows

转到“设置”以激活 Windows。

COVID-19 and non-alcoholic fatty liver disease: Two ...

<https://pubmed.ncbi.nlm.nih.gov/32589264>

NAFLD can play a role in the outcome of COVID-19 illness due to frequent association with comorbidities. Initial evidences suggest that increased liver fibrosis in NAFLD might affect COVID-19 outcome. In addition, long-term monitoring of post-COVID-19 ...

Cited by: 24

Author: Piero Portincasa, Marcin Krawczyk, Marcin ...

Publish Year: 2020

NAFLD and COVID-19: a Pooled Analysis

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7646222>

Nonalcoholic fatty liver disease is associated with a higher risk of symptomatic, severe, and progressive COVID-19. The association is significant even after adjusting for an important confounding factor, obesity. Hence, it can be concluded that the hepatic manifestation of COVID-19 is independently linked to the severity of coronavirus disease.

Cited by: 8

Author: Sonali Sachdeva, Harshwardhan Khandait, ...

Publish Year: 2020

Non-alcoholic fatty liver disease and clinical outcomes in ...

<https://europepmc.org/article/PMC/PMC8011308> ▾

Mar 31, 2021 · Non-alcoholic fatty liver disease (NAFLD) patients represent a vulnerable population that may be susceptible to more severe COVID-19. Moreover, not only the underlying NAFLD may influence the progression of COVID-19, but the COVID-19 may affect the clinical course of NAFLD as well.

Author: Ambrish Singh, Salman Hussain, Benny ... Publish Year: 2021

Name of Journal: *World Journal of Hepatology*

Manuscript NO: 67945

Manuscript Type: MINIREVIEWS

Association of Non-alcoholic Fatty Liver Disease (NAFLD) and COVID-19: A literature review of current evidence

NAFLD and COVID-19

Abstract

The Coronavirus Disease-19 (COVID-19) pandemic has swept through nations, crippled economies and caused millions of deaths worldwide. Many people diagnosed with Covid-19 infections are often found to develop liver injury, which, in a small portion of patients, progresses to severe liver disease. Liver injury in the form of elevated transaminases, hyperbilirubinemia and alterations in serum albumin has been observed to be higher in patients with severe forms of the disease. Those who already have insult to the liver from chronic diseases such as Non-alcoholic Fatty Liver Disease (NAFLD)

Match Overview

1	Crossref 48 words Rui Huang, Li Zhu, Jian Wang, Leyang Xue et al. "Clinical ... eatures of Patients With COVID-19 With Nonalcoholic Fatty	2%
2	Crossref 41 words Roberta Portone, Benjamin H. Mullish, Sujit K. Mukherjee, R ooshi Nathwani et al. "In-hospital mortality is associated v ...	1%
3	Internet 41 words crawled on 14-Mar-2021 neuro.unboundmedicine.com	1%
4	Crossref 38 words Sonali Sachdeva, Harshwardhan Khandait, Jonathan Kopel, Mark M. Aloysius, Rupak Desai, Hemant Goyal. "NAFLD v ...	1%
5	Internet 32 words crawled on 15-Feb-2021 www.unboundmedicine.com	1%
6	Crossref 28 words Kamran Mushtaq, Muhammad Umair Khan, Fatima Iqbal, D eema Hussam Alsoub et al. "NAFLD is a predictor of liver in	1%
7	Crossref 25 words Philip O'Gorman, Suzanne Norris. "Exercising in the COV ... D-19 era: implications in non-alcoholic fatty liver disease (N	1%
8	Internet 22 words crawled on 09-Dec-2020 www.mysciencework.com	1%

国内版

国际版

Association of non-alcoholic fatty liver disease and COVID-19: A lit



ALL

IMAGES

VIDEOS

39,300 Results

Any time ▾

[NAFLD and COVID-19: a Pooled Analysis](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7646222>

Nonalcoholic fatty liver disease is associated with a higher risk of symptomatic, severe, and progressive COVID-19. The association is significant even after adjusting for an important confounding factor, obesity. Hence, it can be concluded that the hepatic manifestation of COVID-19 is independently linked to the severity of coronavirus disease.

Cited by: 9

Author: Sonali Sachdeva, Harshwardhan Khandait...

Publish Year: 2020

[COVID-19 and non-alcoholic fatty liver disease: Two ...](#)

<https://pubmed.ncbi.nlm.nih.gov/32589264>

NAFLD can play a role in the outcome of COVID-19 illness due to frequent association with comorbidities. Initial evidences suggest that increased liver fibrosis in NAFLD might affect COVID-19 outcome. In addition, long-term monitoring of post-COVID-19 ...

Cited by: 27

Author: Piero Portincasa, Marcin Krawczyk, Marc...

Publish Year: 2020

[Metabolic surgery may protect against admission for COVID ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC8238640>

SARS-CoV-2 (COVID-19) disease causes significant morbidity and mortality through increased inflammation and thrombosis. Nonalcoholic fatty liver disease (NAFLD) and nonalcoholic steatohepatitis (NASH) are states of chronic inflammation and indicate advanced ...

[Non-alcoholic fatty liver disease and clinical outcomes in ...](#)

<https://europepmc.org/article/PMC/PMC8011308> ▾

Mar 31, 2021 · Non-alcoholic fatty liver disease (NAFLD) patients represent a vulnerable population that may be susceptible to more severe COVID-19. Moreover, not only the underlying NAFLD may influence the progression of COVID-19, but the COVID-19 may affect the clinical course of NAFLD ...

Author: Ambrish Singh, Salman Hussain, Ben...

Publish Year: 2021

[Practice Guidelines | AASLD](#)

<https://www.aasld.org/publications/practice-guidelines> ▾