### Discontinuation of oral antivirals in chronic hepatitis B ...

#### https://pubmed.ncbi.nlm.nih.gov/27100145

Any time ▼

The possibility of safe discontinuation of therapy with nucleos (t)ide analogues (NAs) remains one of the most controversial topics in the management of chronic hepatitis B. Therefore, we systematically reviewed the existing data on NA discontinuation in this setting and tried to identify factors affecting the probability of posttherapy remission. A literature search was performed in order to identify all ...

Cited by: 157 Author: George Papatheodoridis, Ioannis Vlachog...

Publish Year: 2016

1.860.000 Results

### Outcome after discontinuing antiviral agents during ...

### https://www.ncbi.nlm.nih.gov/pubmed/23273664

However, there are limited data on the outcome in pregnant women after withdrawal of antiviral agents. OBJECTIVES: We aimed to investigate the outcome of stopping antiviral agents in pregnant women with chronic hepatitis B virus (HBV) infection. STUDY DESIGN: In this single-center, retrospective cohort study, 12 pregnant patients who had ...

Cited by: 19 Author: Hee Yeon Kim, Jong Young Choi, Chung-...

Publish Year: 2013

# Why not to stop antiviral treatment in patients with ... https://pubmed.ncbi.nlm.nih.gov/29427480

At present, there is no consensus among the three major guidelines for the treatment of chronic hepatitis B on whether or not to stop antiviral treatment. One of the main reasons for this controversy is that virological relapse has been well documented in patients with chronic hepatitis B who stop treatment. Relapse rate is particularly high in patients who are HBeAg-negative when treatment begins, with reported relapse rates of up to 70% 36 months ...


Cited by: 7 Author: Sebastián Marciano, Adrián Gadano


Publish Year: 2018

### Antiviral response is not sustained after cessation of ...

### https://www.ncbi.nlm.nih.gov/pubmed/27769101

Despite meeting the requirements for treatment discontinuation, approximately half of the CHB patients treated with lamivudine relapsed. Thus, the antiviral response is not reliably sustained after lamivudine treatment cessation. J. Med. Virol. 89:849-856, 2017. © 2016 Wiley Periodicals, Inc. © 2016 Wiley Periodicals, Inc. PMID: 27769101







### Challenges in the discontinuation of chronic hepatitis B antiviral ag







ALL

**IMAGES** 

VIDEOS

520 000 Results

Any time ▼

### Why not to stop antiviral treatment in patients with ...

### https://www.ncbi.nlm.nih.gov/pubmed/29427480

Why not to stop antiviral treatment in patients with chronic hepatitis B. Marciano S(1), Gadano A(1). Author information: (1)Liver Unit and Department of Research, Hospital Italiano de Buenos Aires, Buenos Aires, Argentina. Treatment of chronic hepatitis B with entecavir or tenofovir leads to viral suppression in almost all patients.

Cited by: 8 Author: Sebastián Marciano, Adrián Gadano

Publish Year: 2018

### Antiviral therapy of chronic hepatitis B: can we clear the ...

### https://www.ncbi.nlm.nih.gov/pubmed/15646643

Antiviral therapy of chronic HBV infection remains a clinical challenge. Once this infection has been-established, the viral genome persists for life, either as an integrated genome or as episomal covalently closed circular DNA (cccDNA).

Cited by: 73 Author: Fabien Zoulim

Publish Year: 2004

# Outcome after discontinuing antiviral agents during ...

### https://www.ncbi.nlm.nih.gov/pubmed/23273664

However, there are limited data on the outcome in pregnant women after withdrawal of antiviral agents. OBJECTIVES: We aimed to investigate the outcome of stopping antiviral agents in pregnant women with chronic hepatitis B virus (HBV) infection. STUDY DESIGN: In this single-center, retrospective cohort study, 12 pregnant patients who had ...

Cited by: 23 Author: Hee Yeon Kim, Jong Young Choi, Chung-...

Publish Year: 2013

## Successful Withdrawal of Antiviral Treatment in Kidney ...

### https://pubmed.ncbi.nlm.nih.gov/24628837

Background: The optimal duration of antiviral therapy for kidney transplant recipients (KTR) with chronic hepatitis B virus (HBV) infection remains unclear. We reported the long-term outcomes after withdrawal of antiviral agent in KTR with chronic HBV infection. Methods: We retrospectively investigated the hepatitis B surface antigen (HBsAg)-positive KTR with antiviral agents between January ...

Cited by: 16 Author: J.-H. Cho, J.-H. Lim, G.-Y. Park, J.-S. Ki...