

74902_Auto_Edited-check.docx

Name of Journal: *World Journal of Clinical Cases*

Manuscript NO: 74902

Manuscript Type: CASE REPORT

Not all liver tumors are alike – an accidentally discovered primary hepatic leiomyosarcoma: A case report

Garrido I *et al.* An accidentally discovered primary hepatic leiomyosarcoma

Isabel Garrido, Patrícia Andrade, João Pacheco, Elisabete Rios, Guilherme Macedo

Abstract

BACKGROUND

Primary hepatic leiomyosarcoma is a very rare entity that originates from smooth muscle. Preoperative diagnosis requires a high degree of suspicion due to atypical clinical presentation and non-specific imaging features.

CASE SUMMARY

We report the case of a 42-years-old man, with no relevant past medical history, accidentally diagnosed with a nodular liver lesion on a routine abdominal ultrasound. Liver function tests and hematology parameters as well as the tumor markers were normal. A contrast-enhanced abdominal computed tomography scan revealed a heterogenous hepatic lesion measuring 40 x 30 mm, adjacent to the left branch of the portal vein and the round ligament. Due to the unclear diagnosis, the patient underwent surgical resection. Histopathological and immunohistochemical examinations confirmed complete (R0) resection of a hepatic leiomyosarcoma. The patient remains without any signs of tumor recurrence for more than two years.

CONCLUSION

We report a rare case of accidentally diagnosed primary hepatic leiomyosarcoma originating from the portal vein or the round ligament. Although this tumor has aggressive metastatic potential, a tumor-free resection margin is essential to improve survival.

Key Words: Primary hepatic leiomyosarcoma; Portal vein; Round ligament; Surgical resection; Case report

Garrido I, Andrade P, Pacheco J, Rios E, Macedo G. Not all liver tumors are alike – an accidentally discovered primary hepatic leiomyosarcoma: A case report. *World J Clin Cases* 2022; In press

Core Tip: ² Sarcomas comprise only 1%-2% of all primary liver malignancies and leiomyosarcoma is even rarer. We report a rare case of primary hepatic leiomyosarcoma accidentally diagnosed and surgically treated. The diagnosis requires a high level of suspicion because the clinical scenario and cross-imaging are not specific. Thus, histological examination is the only way to reach the diagnosis. In our case, the tumor probably originated from the left branch of the portal vein or the round ligament. Radical hepatectomy is the cornerstone of treatment. However, this tumor ³ has aggressive metastatic potential and is usually diagnosed in locally advanced or metastatic disease. Two years after surgical resection, our patient is alive and with no evidence of tumor recurrence, probably because the diagnosis was established at an early stage and the surgery achieved a tumor-free margin.

² INTRODUCTION

Sarcomas comprise only 1%-2% of all primary liver malignancies and leiomyosarcoma is even rarer^[1]. In the vast majority of cases of primary hepatic leiomyosarcoma, the tumor originates in the inferior vena cava. ⁴ Clinical presentation and imaging features are non-specific and can mimic the most frequent primary liver tumors namely

hepatocellular carcinoma and intrahepatic cholangiocarcinoma. This tumor ³ has aggressive metastatic potential and is usually diagnosed in a locally advanced or metastatic disease^[2]. We report a rare case of primary hepatic leiomyosarcoma originating from the portal vein or the round ligament, accidentally diagnosed and surgically treated.

CASE PRESENTATION

Chief complaints

The patient was asymptomatic.

History of present illness

A 42-years-old man was referred for gastroenterology consultation due to a nodular hepatic lesion identified in a routine abdominal ultrasound.

¹

History of past illness

Healthy, with no specific diseases.

Personal and family history

There was no family history of liver disease.

Physical examination

There were no relevant changes on physical examination.

Laboratory examinations

Liver tests were normal. The metabolic, virological and autoimmune study was negative. Alpha-fetoprotein and CA 19-9 were also normal.

Imaging examinations

A contrast-enhanced abdominal computed tomography scan was performed, revealing a nodular hepatic lesion measuring 40 mm × 30 mm, adjacent to the left branch of the

portal vein, in the transition from the right to the left lobe (Figure 1). The tumor was contrast-enhancing, with heterogeneous areas inside.

MULTIDISCIPLINARY EXPERT CONSULTATION

The imaging characteristics did not allow for a categorical distinction between hydatid cyst or other etiology. Therefore, it was decided at the multidisciplinary team meeting to perform surgical resection.

FINAL DIAGNOSIS

Histological examination showed complete resection of malignant mesenchymal neoplasia consisting of spindle cells with nuclei elongated to ovoid, with atypia and marked pleomorphism (Figure 2). In the immunohistochemical study, diffuse expression of actin and desmin was observed in neoplastic cells, in the absence of S100 expression. A chest-abdomen-pelvis computed tomography scan was performed in the postoperative period, with no evidence of residual or metastatic lesions. Moreover, this examination made it clear that it was a primary lesion. Thus, primary hepatic leiomyosarcoma was diagnosed with origin in the portal vein wall or the round ligament.

TREATMENT

In the explored laparoscopy, an exophytic lesion in segment IV was identified, involving the left branch of the portal vein and the round ligament; non-anatomic resection was performed without complications

OUTCOME AND FOLLOW-UP

The patient remained under clinical surveillance. Two years after surgery, he is asymptomatic, with no evidence of tumor recurrence.

DISCUSSION

Primary hepatic leiomyosarcoma is a rare type of liver sarcoma; so far, less than one hundred cases have been reported in the literature. Its diagnosis requires a high level of suspicion because the clinical scenario and cross-imaging are not specific^[3]. In fact, patients are often asymptomatic or their symptoms are non-specific. Furthermore, alpha-fetoprotein and other serological markers are usually normal. Thus, histological examination is the only mean to achieve the diagnosis, as in our case.

Primary hepatic leiomyosarcoma can arise from intrahepatic vascular structures, bile ducts or the round ligament. In our case, the tumor probably originated from the left branch of the portal vein or the round ligament. ⁶ The first description of leiomyosarcoma by Perl in 1871 involved a tumor of the inferior vena cava^[4]. Indeed, leiomyosarcoma of vascular origin often occurs in the inferior vena cava. On the other hand, those arising from portal vein are extremely rare and only six cases have been reported in the literature (Table 1)^[5-10].

² No risk factors have been identified yet, although the increasing incidence of the tumor has been noted among immunosuppressed patients^[11]. Nevertheless, more studies are needed to investigate the underlying pathogenetic mechanisms of this uncommon entity.

¹ Due to the rarity of primary hepatic sarcomas in general and primary hepatic leiomyosarcoma in particular, the standard of care has not been defined. Radical R0 hepatectomy, in the form of wedge resection, segmentectomy, lobectomy or extended hepatectomy, is the cornerstone of successful management of primary hepatic leiomyosarcoma¹². In our case, as the tumor was located in the liver, we were able to perform a surgical resection, obtaining a tumor-free margin. However, this tumor ³ has aggressive metastatic potential and is usually diagnosed in locally advanced or metastatic disease. The role of adjuvant or neoadjuvant chemotherapy and radiotherapy regimens in these circumstances has not yet been established. Similarly, orthotopic liver transplantation remains controversial.

Little is known about the natural history of the disease. However, survival rates are relatively low as the tumor is usually diagnosed in advanced stages. Chi *et al*^[1] reported

a median overall survival of 19 mo (range 0-181 mo) with 1-, 2-, and 5-year survival rates of 61.2%, 41.1%, and 14.5%, respectively. The smaller size of the lesion and tumor-free resection margin were identified as independent predictors of improved survival. Two years after surgical resection, our patient is alive and with no evidence of tumor recurrence, probably because the diagnosis was established at an early stage and the surgery achieved a tumor-free margin.

CONCLUSION

Primary hepatic leiomyosarcoma is a rare malignant disease with a poor prognosis. The authors report a rare case of resected primary hepatic leiomyosarcoma originating from the portal vein or the round ligament. Although there is no standard treatment due to the rarity of this disease, we showed that liver resection with a tumor-free resection margin could be an effective treatment in the early stages. Gastroenterologists should be alert to this unusual entity as early diagnosis requires a high degree of suspicion due to atypical clinical presentation and nonspecific imaging features. We emphasize the need for a global database for these rare tumors to promote a better understanding and standardized treatment of these patients.

REFERENCES

- 1 **Chi M**, Dudek AZ, Wind KP. Primary hepatic leiomyosarcoma in adults: analysis of prognostic factors. *Onkologie* 2012; **35**: 210-214 [PMID: 22488093 DOI: 10.1159/000337416]
- 2 **Shivathirthan N**, Kita J, Iso Y, Hachiya H, Kyunghwa P, Sawada T, Kubota K. Primary hepatic leiomyosarcoma: Case report and literature review. *World J Gastrointest Oncol* 2011; **3**: 148-152 [PMID: 22046492 DOI: 10.4251/wjgo.v3.i10.148]
- 3 **Hamed MO**, Roberts KJ, Merchant W, Lodge JP. Contemporary management and classification of hepatic leiomyosarcoma. *HPB (Oxford)* 2015; **17**: 362-367 [PMID: 25418451 DOI: 10.1111/hpb.12366]

- 4 **Halyo V.** Martin L. Perl (1927-2014). *Nature* 2014; **516**: 330 [PMID: 25519123 DOI: 10.1038/516330a]
- 5 **Wilson SR,** Hine AL. Leiomyosarcoma of the portal vein. *AJR Am J Roentgenol* 1987; **149**: 183-184 [PMID: 3495979 DOI: 10.2214/ajr.149.1.183]
- 6 **Sundaresan M,** Kelly SB, Benjamin IS, Akosa AB. Primary hepatic vascular leiomyosarcoma of probable portal vein origin. *J Clin Pathol* 1990; **43**: 1036 [PMID: 2266177 DOI: 10.1136/jcp.43.12.1036-a]
- 7 **Boudjema K,** Sulpice L, Levi Sandri GB, Meunier B. Portal vein leiomyosarcoma, an unusual cause of jaundice. *Dig Liver Dis* 2014; **46**: 1053-1054 [PMID: 25087678 DOI: 10.1016/j.dld.2014.07.001]
- 8 **Gohrbandt AE,** Hansen T, Ell C, Heinrich SS, Lang H. Portal vein leiomyosarcoma: a case report and review of the literature. *BMC Surg* 2016; **16**: 60 [PMID: 27580598 DOI: 10.1186/s12893-016-0174-x]
- 9 **Gaignard E,** Bergeat D, Stock N, Robin F, Boudjema K, Sulpice L, Rayar M, Meunier B. Portal vein leiomyosarcoma: A rare case of hepatic hilar tumor with review of the literature. *Indian J Cancer* 2019; **56**: 83-85 [PMID: 30950452 DOI: 10.4103/ijc.IJC_199_18]
- 10 **Esposito F,** Lim C, Baranes L, Salloum C, Feray C, Calderaro J, Azoulay D. Primary leiomyosarcoma of the liver: Two new cases and a systematic review. *Ann Hepatobiliary Pancreat Surg* 2020; **24**: 63-67 [PMID: 32181431 DOI: 10.14701/ahbps.2020.24.1.63]
- 11 **Brichard B,** Smets F, Sokal E, Clapuyt P, Vermynen C, Cornu G, Rahier J, Otte JB. Unusual evolution of an Epstein-Barr virus-associated leiomyosarcoma occurring after liver transplantation. *Pediatr Transplant* 2001; **5**: 365-369 [PMID: 11560757 DOI: 10.1034/j.1399-3046.2001.00022.x]

ORIGINALITY REPORT

15%

SIMILARITY INDEX

PRIMARY SOURCES

- | | | |
|----------|--|----------------------|
| 1 | www.wjgnet.com
<small>Internet</small> | 54 words — 4% |
| <hr/> | | |
| 2 | Ming Chi, Arkadiusz Z. Dudek, Kenneth P. Wind.
"Primary Hepatic Leiomyosarcoma in Adults:
Analysis of Prognostic Factors", Onkologie, 2012
<small>Crossref</small> | 45 words — 3% |
| <hr/> | | |
| 3 | surgicalcasereports.springeropen.com
<small>Internet</small> | 42 words — 3% |
| <hr/> | | |
| 4 | www.scilit.net
<small>Internet</small> | 35 words — 3% |
| <hr/> | | |
| 5 | www.science.gov
<small>Internet</small> | 20 words — 1% |
| <hr/> | | |
| 6 | Karim Boudjema, Laurent Sulpice, Giovanni Battista
Levi Sandri, Bernard Meunier. "Portal vein
leiomyosarcoma, an unusual cause of jaundice", Digestive and
Liver Disease, 2014
<small>Crossref</small> | 16 words — 1% |

EXCLUDE QUOTES ON

EXCLUDE MATCHES

< 12 WORDS

EXCLUDE BIBLIOGRAPHY ON